



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

### Usage guidelines


Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.


We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

### About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>





600036247S

Q.105. S. 11



E. BIBL. RADCL.

6  
~~7~~ 3  
13


5  
~~7~~ 4  
17

~~55 13 23~~

18949

e.

$\frac{299}{2}$







**IOH. CHRIST. FABRICII**

**HISTOR. NAT. OECON. ET CAMERAL. PROF. KILONIENS.**

**ACAD. HAFNIENS. NORWEG. BEROLINENS. ET**

**LVND. SOC.**

**SPECIES INSECTORVM**

**EXHIBENTES**


**EORVM**

**DIFFERENTIAS SPECIFICAS, SYNONYMA**

**AVCTORVM, LOCA NATALIA, METAMORPHOSIN**

**A. D. I. E. C. T. I. S**

**OBSERVATIONIBVS, DESCRIPTIONIBVS.**



---

***T O M. II.***

---

***HAMBVRGI et KILONII***

**IMPENSIS CAROL. ERNEST. BOMNII**

**MDCCLXXXI.**

1. The first part of the document discusses the importance of maintaining accurate records of all transactions and activities. It emphasizes that proper record-keeping is essential for transparency and accountability, particularly in financial matters. The text outlines various methods for organizing and storing data, including digital databases and physical filing systems. It also mentions the need for regular audits and reviews to ensure the integrity of the information.

2. The second part of the document focuses on the role of communication in achieving organizational goals. It highlights the importance of clear and concise communication, both internally and externally. The text provides guidelines for effective communication, such as using appropriate language, listening actively, and providing feedback. It also discusses the benefits of open communication and how it can foster a collaborative work environment.

3. The third part of the document addresses the issue of resource management. It discusses the importance of identifying and allocating resources effectively to support the organization's mission. The text provides strategies for managing resources, including budgeting, prioritizing tasks, and delegating responsibilities. It also mentions the need for regular monitoring and evaluation of resource usage to ensure optimal performance.

4. The fourth part of the document discusses the importance of continuous learning and improvement. It emphasizes that organizations should strive to stay up-to-date with the latest trends and technologies in their field. The text provides suggestions for promoting a culture of learning, such as offering training opportunities, encouraging innovation, and seeking feedback from employees and customers. It also mentions the importance of documenting lessons learned and applying them to future projects.

5. The fifth part of the document discusses the importance of maintaining a strong ethical foundation. It emphasizes that organizations should adhere to high standards of ethics and integrity in all their dealings. The text provides guidelines for ethical behavior, such as being honest, transparent, and respectful. It also mentions the importance of having a clear code of ethics and providing training to ensure that all employees understand and follow the principles.

6. The sixth part of the document discusses the importance of maintaining a strong relationship with stakeholders. It emphasizes that organizations should engage with their stakeholders and listen to their concerns. The text provides suggestions for building strong relationships, such as holding regular meetings, providing timely responses, and being open to feedback. It also mentions the importance of having a clear communication plan and identifying key stakeholders.

7. The seventh part of the document discusses the importance of maintaining a strong financial position. It emphasizes that organizations should manage their finances carefully and ensure that they have sufficient resources to support their operations. The text provides guidelines for financial management, such as budgeting, monitoring expenses, and seeking funding opportunities. It also mentions the importance of having a clear financial strategy and regularly reviewing the financial performance.

8. The eighth part of the document discusses the importance of maintaining a strong legal and regulatory compliance. It emphasizes that organizations should ensure that they are compliant with all applicable laws and regulations. The text provides suggestions for maintaining compliance, such as staying up-to-date with legal changes, conducting regular audits, and seeking legal advice when needed. It also mentions the importance of having a clear policy on legal and regulatory compliance.

9. The ninth part of the document discusses the importance of maintaining a strong environmental and social responsibility. It emphasizes that organizations should strive to minimize their environmental impact and promote social responsibility. The text provides suggestions for maintaining environmental and social responsibility, such as reducing waste, conserving resources, and supporting community development. It also mentions the importance of having a clear policy on environmental and social responsibility.

10. The tenth part of the document discusses the importance of maintaining a strong overall organizational performance. It emphasizes that organizations should strive to achieve their goals and objectives through effective management and leadership. The text provides suggestions for maintaining overall performance, such as setting clear goals, monitoring progress, and providing support and encouragement to employees. It also mentions the importance of having a clear vision and mission statement.



# SPECIES INSECTORVM

## PARS II.

---

### CLASSIS VI.

### GLOSSATA.

---

#### 141. PAPILIO.

\* Equites Troes.

1. **P.** E. T. alis caudatis nigris, posticis macula cyanea *Paris*.  
ocelloque purpureo, subtus lunulis septem. *Syst.*  
*Ent.* 442. 1. *Linn. Syst. Nat.* 2. 745. 3. *Mus. Lud.*  
*Vlr.* 184.

*Papilio Paris.* *Clerk. Icon.* tab. 13. fig. 1.

*Papilio Paris.* *Cram. Inf.* 9. tab. 103. fig. A. B.

*Knorr. del.* tab. 6. 3. fig. 1.

*Drury Inf.* 1. tab. 12. fig. 1.

Habitat in Asia frequens.

2. P. E. T. alis caudatis subconcoloribus nigris, posticis *Bianor*.  
lunulis quinque rufis.

*Papilio Bianor.* *Cram. Inf.* 9. tab. 103. fig. C.

Habitat in China.

Vix satis distinctus a P. Paride. Differt tamen defectu  
maculae cyaneae in alis posticis, lunulisque quinque  
rubris et in pagina superiori conspicuis.

*Fabr. Sp. Inf. T. II.*

A

3. P.

2 CLASS. VI. GLOSSATA. Papilio.

*Thefeus*. 3. P. E. T. alis caudatis concoloribus fuscis, posticis lunulis nouem rubris albo punctatis.

*Papilio Thefeus*. *Cram. Inf.* 15. tab. 180. fig. B.

Habitat in Infula Sumatra.

Multo praecedenti minor. Caput et thorax albo punctata. Alae anticae fuscae, basi obscuriores, immaculatae, posticae fuscae apicē obscuriores lunulis nouem rubris atomis albis irroratis.

*Polytes*. 4. P. E. T. alis caudatis concoloribus nigris, posticis macula quintuplici alba lunulisque rubris, corpore nigro. *Syst. Ent.* 443. 2. *Linn. Syst. Nat.* 2. 746. 5. *Mus. Lud. Vlr.* 186.

*Papilio Polytes*. *Clerk. Icon.* tab. 14. fig. 1.

*Papilio Alphenor*. *Cram. Inf.* 8. tab. 90. fig. B.

*Sulz. Hist. Inf.* tab. 12. fig. 3.

*Seb. Mus.* 4. tab. 27. fig. 23. 24.

Habitat in Asiae Chalcas. *D. Koenig.*

*Hector*. 5. P. E. T. alis caudatis concoloribus nigris, anticis fascia alba, posticis maculis rubris. *Syst. Ent.* 443. 4. *Linn. Syst. Nat.* 2. 745. 2. *Mus. Lud. Vlr.* 181.

*Papilio Aristolochiae* alis caudatis nigris concoloribus, posticis fascia maculari alba lunulisque rubris, capite caudaque sanguineis. *Syst. Ent.* 443. 3.

*Papilio Hector*. *Cram. Inf.* 12. tab. 141. fig. A.

*Papilio Hector*. *Clerk. Icon.* tab. 33. fig. 1.

*Papilio indica maxima nigra*. *Raj. Inf.* 134. 137.

*Sulz. Hist. Inf.* tab. 12. fig. 1.

*Seb. Mus.* 4. tab. 28. fig. 23. 24.

Habitat in Indiae Aristolochiis. *D. Koenig.*

Conf. *Papilio Romulus*. *Cram. Inf.* 4. tab. 43. fig. A.  
Vix distinctus videtur.

*Ascanius*. 6. P. E. T. alis caudatis concoloribus atris, fascia comuni alba, posticarum rubro nebulosa.

*Papilio Ascanius*. *Cram. Inf.* 2. tab. 14. fig. A.

Habitat in Brasilia. *Mus. Dom. Teats.*

Affinis *P. Hectori*. Corpus nigrum pectore sanguineo maculato. Fascia communis niuea posticarum rubro inquinata marginem exteriorem haud attingit. Lunulae rubrae in alis posticis.

CLASS. VI. GLOSSATA. Papilio.

3

7. P. E. T. alis caudatis nigris, posticis macula alba lunulaque duplici purpurea. *Syst. Ent.* 443. 5. *Helennus.*  
*Papilio Helennus* E. T. alis caudatis nigris, posticis macula alba, subtus tribus albidis lunulisque septem ferrugineis. *Linn. Syst. Nat.* 2. 745. 4. *Mus. Lud. Vlr.* 185.  
*Papilio Helennus.* *Cram. Inf.* 13. tab. 153. fig. A. B.  
*Papilio Helennus.* *Clerk. Icon.* tab. 13. fig. 2.  
*Ehret. pict.* tab. 10.  
*Edw. Aves* tab. 342.  
*Seb. Mus.* tab. 46. fig. 17. 18.  
Habitat in Asia.  
Ocellus purpureus ad angulum ani et sub ocello lunula rubra.
8. P. E. T. alis caudatis concoloribus atris albo maculatis, posticis lunulis marginalibus rubris. *Syst. Ent.* 443. 6. \* *Antenor.*  
*Papilio Antenor.* *Drury Inf.* 2. tab. 3. fig. 1.  
Habitat in India.
9. P. E. T. alis caudatis nigris, anticis punctis marginalibus pallidis, posticis supra pallido subtus fulvo maculatis. *Syst. Ent.* 444. 7. *Linn. Syst. Nat.* 2. 746. 6. *Mus. Lud. Vlr.* 187. *Troilus.*  
*Papilio Troilus.* *Cram. Inf.* 18. tab. 207. fig. A. B.  
*Drury Inf.* 1. tab. 11. fig. 2. 3. 5.  
*Petiv. Mus.* 51. n. 523.  
Habitat in Indiis.  
Macula ferruginea ad marginem tenuiorem alae posticae et altera ad angulum ani.
10. P. E. T. alis caudatis nigris, subtus basi rubro maculatis, posticis maculis septem rubris subanhularibus. *Syst. Ent.* 444. 8. *Linn. Syst. Nat.* 2. 746. 7. *Mus. Lud. Vlr.* 188. *Deiphobus.*  
*Papilio Deiphobus.* *Clerk. Icon.* tab. 25.  
*Papilio Deiphobus.* *Cram. Inf.* 18. tab. 181. fig. A. B.  
*Papilio Luzojicus* maximus nigrescens, maculis coctineis. *Petiv. Gazoph.* tab. 11. fig. 8.  
*Seb. Mus.* 4. tab. 47. fig. 5. 6.  
*Edw. Av.* tab. 346.

Habitat in Asia.

Lunulae rubrae alae posticae apice flavescent.

*Alphenor*. 11. P. E. T. alis caudatis concoloribus fuscis, basi atris, maculis anticarum rufa, posticarum albis.

*Papilio Antenor*. *Cram. Inf.* 14. tab. 166. fig. A.

Habitat in China.

Alae anticae fuscae basi atrae macula magna rufa, posticae fuscae basi venisque atris maculisque aliquot albis. Margo interior rufus.

*Pelaus*. 12. P. E. T. alis caudatis atris, anticis fascia, posticis lunulis marginalibus albis punctisque duobus rubris.

*Syst. Ent.* 444. 9. \*

*Papilio torquatus*. *Cram. Inf.* 15. tab. 177. fig. A. B. C.

Habitat in India. *Dom. Drury*.

*Polyxenes*. 13. P. E. T. alis caudatis atris, fasciis duabus macularibus flavis, angulo ani fulvo puncto nigro. *Syst. Ent.*

444. 10. \*

Habitat in Americae meridionalis insulis.

Affinis P. Troilo.

*Phorbantha*. 14. P. E. T. alis caudatis nigris coeruleo maculatis, posticarum fascia interrupta subtus alba. *Syst. Ent.*

445. 11. *Linn. Mant.* 1. 535.

*Aubent. miscell.* tab. 43. fig. 1. 2.

Habitat in Caienna.

*Philenor*. 15. P. E. T. alis caudatis nigris, posticis virescenti nitidis, subtus maculis septem rubris subocellaribus. *Syst.*

*Ent.* 445. 12. \*

*Papilio Philenor* E. T. alis caudatis nigris, posticis subtus nitenti cyaneis, ocellis septem concatenatis.

*Linn. Mant.* 1. 535.

*Papilio Astenous*. *Drury Inf.* 1. tab. 11. fig. 1. 4.

*Papilio Astenous*. *Cram. Inf.* 18. tab. 208. fig. A. B.

Habitat in America boreali.

*Pammon*. 16. P. E. T. alis caudatis concoloribus nigris, margine maculatis, posticis fascia maculis septem albis. *Syst. Ent.*

445. 13. *Linn. Syst. Nat.* 2. 746. 8. *Mus. Lud. Vlr.*

189.

*Papilio*



*Papilio Pammon. Clerk. Icon. tab. 14. fig. 2.*

*Papilio Pammon. Cram. Inf. 12. tab. 141. fig. B.*

*Klemann. Inf. 1. tab. 2. fig. 2. 3.*

*Seb. Mus. 4. tab. 27. fig. 21. 22.*

— — — 41. fig. 11. 12.

— — — 34. fig. 23. 24.

*Aubent. miscell. 1. tab. 43. fig. 1.*

Habitat in Asiae Chalcas. *D. Koenig.*

17. P. E. T. alis caudatis nigris, fascia communi alba, posticis subtus basi apiceque rubro maculatis. *Asus.*

Habitat in America meridionali. *Dom. Teats.*

Corpus paruum nigrum. Thorax linea laterali cinerea. Punctum cinereum vtrinque in pectore et linea lateralis subtus in abdomine. Alae anticae concolores, nigrae fascia alba. Posticae nigrae supra fascia alba lunulis tribus coccineis ad angulum ani, et quinque albis ad marginem, subtus fuscae fascia alba, punctis bascos rubris, linea ad marginem tenuiorem lunulisque tribus anguli ani rubris. Lunulae quatuor albae marginales.

18. P. E. T. alis caudatis concoloribus nebulosis, anticis macula flava, posticis macula ani fulva. *Glaucus.*

*Syst. Ent. 445. 14. Linn. Syst. Nat. 2. 746. 9. Mus. Lud. Vlr.*

190.

*Papilio Glaucus. Clerk. Icon. tab. 24. fig. 1.*

*Papilio Glaucus. Cram. Inf. 12. tab. 139. fig. A. B. ?*

Habitat in America.

Alae posticae subtus lunulis rufis flavisque.

19. P. E. T. alis caudatis subconcoloribus nigris basi rufis, posticis macula octuplici alba. *Achates.*

*Papilio Achates. Cram. Inf. 16. tab. 182. fig. A. B.*

*Sulz. Hist. Inf. tab. 12. fig. 2.*

Habitat in China.

Nimis affinis videtur *P. Agenori*, at caudatus. Alae anticae fuscae, striatae basi vtrinque macula sanguinea. Posticae fuscae basi supra macula atra, subtus sanguinea, in disco macula magna alba futuris octofida. Margo posticus lunulis rubris.

*Polydorus*. 20. P. E. T. alis subcaudatis concoloribus nigris, posticis macula alba futuris sexfida lunulisque septem rubris. *Syst. Ent.* 446. 15. *Linn. Syst. Nat.* 2. 746. 10. *Amoen. acad.* 6. 401. 40.

*Papilio Polydorus*. *Clerk. Icon.* tab. 23. fig. 2.

*Papilio Polydorus*. *Cram. Inf.* 11. tab. 128. fig. A. B.

*Kleman. Inf.* 1. tab. 8. fig. 1. 2.

*Seb. Mus.* 4. tab. 28. fig. 21. 22.

Habitat in India orientali.

*Priamus*. 21. P. E. T. alis denticulatis holosericeis, anticis supra viridibus, macula atra, posticis maculis sex nigris. *Syst. Ent.* 446. 16. *Linn. Syst. Nat.* 2. 744. 1. *Musf. Lud. Vlr.* 182. *Amoen. acad.* 5. tab. 3.

*Papilio Priamus*. *Cram. Inf.* 2. tab. 23. fig. A. B.

*Papilio Priamus*. *Clerk. Icon.* tab. 17.

*Papilio amboinensis* viridi et nigro holosericeus, infignis. *Vincent. Musf.* 10.

*Papilio* siue *Atlas amboinensis* alis superioribus holosericeo nigris, infitis viridibus. *Musf. Petropol.* 664.

*Aubent. miscell.* 1. tab. 45.

Habitat in Amboina.

*Rénus*. 22. P. E. T. alis dentatis subconcoloribus nigris, posticis utrinque maculis flavis marginalibus. *Gen. Inf. Mant.* 250. \*

*Papilio Hippolythus*. *Cram. Inf.* 1. tab. 10. fig. A. B.

— — — — — tab. 11. fig. A. B.

*Seb. Musf.* 4. tab. 46. fig. 11. 12. 19. 20.

Habitat in Amboina.

*Memnon*. 23. P. E. T. alis dentatis, omnibus basi subtus rubro notatis. *Linn. Syst. Nat.* 2. 747. 13. *Musf. Lud. Vlr.* 193.

*Papilio Laomedon*. *Cram. Inf.* 5. tab. 50. fig. A. B.

*Papilio Androgeus*. *Cram. Inf.* 8. tab. 91. fig. A. B. ‡

*Papilio Memnon*. *Cram. Inf.* 8. tab. 91. fig. C.

*Sulz. Hist. Inf.* tab. 12. fig. 5.

*Petiv. Gazoph.* tab. 11. fig. 8.

*Raj. Inf.* 135.

*Seb. Musf.* 4. tab. 5. fig. 6.

— — — — — tab. 16. fig. 10. 11.

Habitat

Habitat in China.

Macula rubra ad basin alae anticae.

24. P. E. T. alis dentatis nigris, posticis subtus macula *Protenor*.  
difformi rubra anguli ani.

*Papilio Memnon. Syst. Ent. 446. 17.*

*Papilio Protenor. Cram. Inf. 5. tab. 49. fig. A. B.*

Habitat Surinami.

Alae anticae supra fuscae, immaculatae, posticae atrae  
puncto gemino anguli ani, subtus atrae lunulis mar-  
ginalibus rubris, macula difformiad angulum ani ru-  
bra puncto nigro.

25. P. E. T. alis dentatis nigris basi sanguineis, posticis *Agenor*.  
disco albo, maculis nigris. *Syst. Ent. 446. 18. Linn.*

*Syst. Nat. 2. 747. 14. Mus. Lud. Vlr. 194.*

*Papilio Agenor. Clerk. Icon. tab. 15.*

*Papilio Agenor. Cram. Inf. 3. tab. 32. fig. A. B.*

*Seb. Musf. 4. tab. 46. fig. 11. 12. 15. 16.*

Habitat in China.

26. P. E. T. alis dentatis concoloribus nigris, posticis ma- *Anchises*.  
culis septem coccineo ouatis. *Syst. Ent. 446. 19.*

*Linn. Syst. Nat. 2. 746. 11. Musf. Lud. Vlr. 191.*

*Papilio Anchises. Clerk. Icon. tab. 29. fig. 1.*

*Sulz. Hist. Inf. tab. 12. fig. 4.*

*Seligm. Av. 6. tab. 102.*

*Merian. Sur. tab. 17.*

*Sloan. Iam. 2. 216. tab. 239. fig. 19. 20.*

*Edw. Av. tab. 207.*

*Ehret. pict. tab. 9.*

*Seb. Musf. 4. tab. 7. fig. 27. 28.*

— — tab. 11. fig. 11. 12.

Habitat in Americis Citro.

*Larva* gregaria, flavo tentaculata, spinosa fusca, albo  
annulata.

*Puppa* brunnea, antice quadridentata.

27. P. E. T. alis dentatis nigris, posticis limbo coerule- *Astya*.  
scente, fascia nigra, subtus fulvo maculatis. *Syst. max.*  
*Ent. 447. 20. \**

Habitat in America. *Musf. D. Hunter.*

- Sarpedon.* 28. P. E. T. alis dentatis fuscis fascia viridi, posticis subtus linea baseos maculisque quinque rubris. *Syst. Ent.* 447. 21. *Linn. Syst. Nat.* 2. 747. 15. *Mus. Lud. Vlr.* 196.

*Cram. Inf.* 11. tab. 122. fig. D. E.

*Seb. Mus.* 4. tab. 37. fig. 3. 4. 15. 16.

Habitat in Asia.

- Polydamas.* 29. P. E. T. alis dentatis nigro aeneis fascia maculari flava, posticis subtus lunulis flexuosis rubris. *Syst. Ent.* 447. 22. *Linn. Syst. Nat.* 2. 747. 12. *Mus. Lud. Vlr.* 192.

*Papilio Polydamas.* *Cram. Inf.* 18. tab. 211. fig. D. E.

*Drury Inf.* 1. tab. 17. fig. 1. 2.

*Merian. Inf. Sur.* tab. 31.

*Seb. Mus.* 4. tab. 39. fig. 2. 3.

Habitat in Hibisco mutabili, Citro Americis.

*Larva* albida fusco maculata.

*Puppa* cinerea nigro maculata.

- Androgeus.* 30. P. E. T. alis dentatis nigro aeneis, posticis subtus lunulis rubris cyaneis flavisque. *Gen. Inf. Mant.* 251. \*

*Papilio Androgeus.* *Cram. Inf.* 2. tab. 16. fig. C. D.

*Papilio Peranthus.* *Cram. Inf.* 17. tab. 204. fig. A. B.

*Kleman. Inf.* 1. tab. 8. fig. 1. 2.

Habitat Surinami. *Mus. Dom. Teats.*

Affinis praecedenti.

- Amphimedon.* 31. P. E. T. alis dentatis concoloribus fuscis, anticis albo radiatis, posticis macula quinquefida rubra lunulisque albis.

*Papilio Amphimedon.* *Cram. Inf.* 17. tab. 194. fig. A.

*Seb. Mus.* 4. tab. 16. fig. 6. 7.

Habitat in Amboina.

Corpus magnum fuscum thorace antice strigis sanguineis.

- Aeneas.* 32. P. E. T. alis dentatis nigris, anticis supra macula viridi, posticis macula palmata sanguinea. *Syst. Ent.* 448.

CLASS. VI. GLOSSATA. Papilio. 9

448. 23. Linn. Syst. Nat. 2. 747. 16. Mus. Lud. Vlr.  
197.

*Papilio Vertumnus.* Cram. Inf. 18. tab. 211. fig. A.  
B. C.

*Papilio Lyfander.* Cram. Inf. 3. tab. 29. fig. C. D.

*Papilio Hyppason.* Cram. Inf. 3. tab. 29. fig. E.

*Papilio Euristeus.* Cram. Inf. 3. tab. 29. fig. F.

Roes. Inf. 4. tab. 2. fig. 2.

Seb. Mus. 4. tab. 7. fig. 25. 26.

— — — — — tab. 20. fig. 17. 18.

Habitat in India.

Conf. *Papilio Sefostris.* Cram. Inf. 18. tab. 211. fig. F.

G. Vix satis distinctus.

33. P. E. T. alis dentatis atris, anticis striga alba, posticis *Lyfander*  
macula media alba lunulisque rubris. Gen. Inf.

Mant. 251. \*

Habitat in India. *D. Fothergil.*

34. P. E. T. alis dentatis virefcentibus, posticis margine *Belus*.  
interiori pallido, subtus fuscis lunulis rubris.

*Papilio Belus.* Cram. Inf. 10. tab. 112. fig. A. B. C.

— — — — — tab. 113. fig. A. B. C.

Habitat Surinami.

Alae subtus fuscae. Pectus fuscum punctis flavis. Ab-  
domen supra pallidum, subtus fuscum vtrinque pun-  
ctis albis.

Variat interdum macula media flavescente alae anticae  
et striga punctorum flavescentium alae posticae.

35. P. E. T. alis dentatis subconcoloribus nigris, posticis *Polymne-*  
apice coerulescentibus nigro maculatis. *flor.*

*Papilio Polymnesfor.* Cram. Inf. 5. tab. 53. fig. A. B.

Habitat in Asia.

Alae basi subtus sanguineo maculatae.

36. P. E. T. alis dentatis concoloribus nigris, anticis albo *Pantho-*  
maculatis, posticis maculis albis nigro foetis. *us.*

Ent. 448. 25. Linn. Syst. Nat. 2. 748. 17. Mus. Lud.

Vlr. 195.

*Papilio Panthous.* Clerk. Icon. tab. 18. 19.

*Papilio Panthous.* Cram. Inf. 11. tab. 123. fig. A.

— — — — — tab. 124. fig. A.

10 CLASS. VI. GLOSSATA. Papilio.

*Papilio Remus. Cram. Inf.* 12. tab. 135. fig. A.

— — — — — tab. 136. fig. A.

*Seb. Musf.* 4. tab. 44. fig. 21 — 23.

— — — — — tab. 45. fig. 17 — 20.

Habitat in Indiis.

**Pandarus.** 37. P. E. T. alis subdentatis concoloribus nigricantibus albo maculatis, posticis flauis, ocellis septem nigris.

*Syst. Ent.* 448. 26. *Linn. Syst. Nat.* 2. 748. 18. *Musf.*

*Lud. Vlr.* 198.

Habitat in India.

Ocelli septem nigri pupilla coerulea, vltima duplicata ad angulum ani.

**Athenous.** 38. P. E. T. alis dentatis concoloribus nigris, anticis macula radiata alba, posticis disco flauo. *Syst. Ent.* 448. 27.\*

*Papilio Pompeus. Cram. Inf.* 3. tab. 25. fig. A.

Habitat ad Cap. bon. sp.

Conf. *Papilio Minos. Cram. Inf.* 17. tab. 195. fig. A.

**Helena.** 39. P. E. T. alis dentatis concoloribus atris, posticis disco communi aurato. *Syst. Ent.* 449. 28. *Linn. Syst. Nat.*

2. 748. 19. *Musf. Lud. Vlr.* 199.

*Papilio Helena. Clerk. Icon.* tab. 22. fig. 1.

*Papilio Helena. Cram. Inf.* 12. tab. 140. fig. A. B.

*Merian. Surin.* in tabula titulo praefixa.

*Seb. Musf.* 4. tab. 45. fig. 9. 10. 11. 12.

Habitat in floribus Arcae Americae.

\*\* Equites Achiui.

**Iafius.** 40. P. E. A. alis bicaudatis fuscis postice lutescentibus subtus fascia characteribusque albis. *Syst. Ent.* 449.

29. *Linn. Syst. Nat.* 2. 749. 26.

*Papilio Iafius. Drury Inf.* 1. tab. 1. fig. 1.

*Cram. Inf.* 16. tab. 186. fig. A. B.

*Seb. Musf.* 4. tab. 46. fig. 21. 22.?

Habitat in Barbaria, Africa.

Fascia in alis posticis magis distincta.

**Pyrrhus.** 41. P. E. A. alis bicaudatis fuscis, fascia communi alba, anticarum dimidiata. *Syst. Ent.* 449. 30. *Linn. Syst.*

*Nat.* 2. 749. 25. *Musf. Lud. Vlr.* 205.

*Papilio*

CLASS. VI. GLOSSATA. Papilio. 11

*Papilio Pyrrhus. Clerk. Icon. tab. 25. fig. 2.*

*Papilio Pyrrhus. Cram. Inf. 8. tab. 89. fig. C. D.*

*Papilio Athamas. Drury Inf. 1. tab. 2. fig. 3. 4.*

*Seb. Musf. 4. tab. 47. fig. 2. 3.*

Habitat in Indiis.

42. P. E. A. alis bicaudatis albis, fasciis fulvis nigro mar- *Camillus.*  
ginatis, posticis macula caudali atra.

Habitat in Africa aequinoctiali. *Musf. Dom. Banks.*

Parvus. Alae anticae concolores, albae fasciis duabus  
fulvis, nigro marginatis, tunc duabus abbreviatis,  
tunc fascia tota fulva, demum striis strigisque fuscis.  
Posticae albae fasciis fulvis fuscisque anguloque ani  
nigro punctato. Caudae duae, altera acuta, nigra,  
albo lineata, altera obtusa, fulva punctis duobus atris.  
Alae subtus subconcolores.

43. P. E. A. alis dentato bicaudatis supra nigris coeruleo *Tiridates.*  
maculatis margineque albo punctato.

*Papilio Tiridates. Cram. Inf. 13. tab. 161. fig. A. B.*

Habitat in Amboina.

Alae omnes supra nigrae strigis duobus punctorum coe-  
ruleorum strigaeque marginali e punctis albis. Sub-  
tus fuscae flavo coeruleoque maculatae maculis posti-  
cis alarum posticarum pupilla alba.

44. P. E. A. alis dentato bicaudatis, fulvis, margine nigris, *Castor.*  
subtus fascia maculisque albis nigra foetis. *Gen. Inf.*  
*Mant. 251.\**

*Papilio Castor. Cram. Inf. 4. tab. 37. fig. E. F.*

Habitat in Guinea.

45. P. E. A. alis dentato bicaudatis, fuscis fascia flava, sub- *Pollux.*  
tus fascia maculisque albis nigra foetis. *Gen. Inf.*  
*Mant. 251.\**

*Papilio Pollux. Cram. Inf. 4. tab. 37. fig. C. D.*

Habitat in Guinea.

Nimis praecedenti affinis.

Conf. *Papilio Pelias. Cram. Inf. 1. tab. 3. fig. C. D.*  
vix distinctus videtur.



15 CLASS. VI. GLOSSATA. Papilio.

*Etheocles*. 46. P. E. A. alis bicaudatis, coeruleo fuscis, vtrinque fascia alba, posticis margine viridi.

*Papilio Etheocles*. *Cramer Inf.* 10. tab. 119. fig. D. E.  
Habitat in Africa aequinoctiali.

Alae supra obscure coeruleae fascia alba anticarum interrupta, posticae apice atrae margine viridi. Subtus brunneae fascia alba, basi lineolae coeruleae, apice maculae nigrae. Angulus ani viridis maculis brunneis.

*Fabius*. 47. P. E. A. alis dentato bicaudatis, fuscis, flavo fasciatis, subtus basi nigro vndatis apice punctis fulvis.

Habitat in India. *Mus. Dom. Banks.*

Paruus. Corpus fuscum capite punctis quatuor albis.

Alae omnes supra fuscae fascia flava anticarum maculari. Posticae bicaudatae margineque flavo maculato. Subtus omnes glaucae, basi nigro vndatae, in medio fascia albida et apice striga e punctis fulvis posticarum duplicata.

*Leilus*. 48. P. E. A. alis caudatis concoloribus nigris, fascia strigisque viridibus nitentibus numerosis. *Syst. Ent.* 449. 31. *Linn. Syst. Nat.* 2. 750. 31. *Mus. Lud.* *Vlr.* 206.

*Papilio Leilus*. *Clerk. Icon.* tab. 27. fig. 1.

*Cram. Inf.* 8. tab. 85. fig. C. D.

*Scop. Ann. Hist. nat.* 5. 110. 111.

*Gronov. Zooph.* 724.

*Kleman Inf.* 1. tab. 2. fig. 1.

*Merian. Sur.* tab. 29.

*Knorr delic.* tab. 6. fig. 1.

*Seb. Mus.* 4. tab. 36. fig. 1. 2.

*Aubent. miscell.* 1. tab. 71. fig. 1.?

Habitat in Citro Americæ.

Antennae obtusae, filiformes. Thorax niger lineis lateralibus viridibus.

Variat macula magna aurea alarum posticarum.

*Sloan. Iam.* 2. 215. tab. 239. fig. 11. 12.

*Papilio Sloanens*. *Cram. Inf.* 8. tab. 85. fig. E. F.

*Larva* pilosa viridis, capite coeruleo.

*Puppa* brunnea, maculata.

49. P. E. A. alis caudatis nigris, viridi fasciatis, subtus argentatis strigis fulvis. *Syst. Ent.* 450. 32.\*  
Habitat in America. *Dom. v. Rohr.*

50. P. E. A. alis caudatis albis: limbo nigro, posticis sub- *Brutus.*  
tus fascia fusca.

*Papilio Merope. Cram. Inf.* 13. tab. 151. fig. A. B. at  
alae ecaudatae.

Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*

Magnus. Caput et thorax nigra, albo punctata. Abdomen albidum linea dorsali nigra lateribusque nigro punctatis. Alae anticae concolores albiae margine exteriori et apicis nigro. Punctum flavum in apice nigro. Posticae caudatae albiae margine postico nigro maculato, subtus albiae fusco venosae fasciaque lata subinterrupta in medio margineque postico fuscis.

51. P. E. A. alis caudatis albis, limbo communi nigro, *Doliceon.*  
posticis utrinque punctis marginalibus albis.

*Papilio Doliceon. Cram. Inf.* 2. tab. 17. fig. C. D.

Habitat in America meridionali. *Mus. Dom. Teats.*

Alae concolores, at subtus annulus niger in disco albo alae posticae. Cauda nigra apice alba.

52. P. E. A. alis caudatis nigris, disco coeruleo radiante *Vlysses.*  
posticis subtus ocellis septem. *Syst. Ent.* 450. 33.

*Linn. Syst. Nat.* 2. 748. 21. *Mus. Lud. Vlr.* 201.

*Papilio Vlysses. Clerk. Icon.* tab. 23. fig. 1.

*Papilio Vlysses. Cram. Inf.* 11. tab. 121. fig. A. B.

*Seb. Mus.* 4. tab. 46. fig. 9. 10.

— — — tab. 47. fig. 9. 10.

Habitat in Asia.

53. P. E. A. alis caudatis concoloribus fuscis, fascia lineari *Patro-*  
obliqua alba apicibusque albis. *Syst. Ent.* 450. 34. *clus.*

*Linn. Syst. Nat.* 2. 749. 24. *Mus. Lud. Vlr.* 204.

*Papilio Patroclus. Drury Inf.* 1. tab. 7. fig. 1.

— — — — — tab. 8. fig. 1.

*Papilio Patroclus. Cram. Inf.* 10. tab. 109. fig. A. B.

— — — — — 17. tab. 198. fig. A.

*Clerk.*

14 CLASS. VI. GLOSSATA. Papilio.

*Clerk. Icon.* tab. 37. fig. 1.

*Seb. Mus.* 4. tab. 47. fig. 13-16.

*Aub. miscell.* tab. 17. fig. 1. 2.

Habitat in Indiis.

*Diomedes.* 54. P. E. A. alis caudatis nigris, supra introrsum viridibus, posticis lunulis cyaneis. *Syst. Ent.* 450. 35. *Linn.*

*Syst. Nat.* 2. 749. 23. *Mus. Lud. Vlr.* 203.

*Papilio Diomedes. Cram. Inf.* 11. tab. 122. fig. A.

*Seb. Mus.* 4. tab. 47. fig. 11. 12.

Habitat in Indiis.

*Varanes.* 55. P. E. A. alis caudatis, supra fuscis fuluo nigroque maculatis basi albis.

*Papilio Varanes. Cram. Inf.* 13. tab. 160. fig. D. E.

*Seb. Mus.* 4. tab. 35. fig. 13. 14.

Habitat in India.

Alae omnes supra basi albae, apice fuscae maculis fulvis nigrisque. Subtus basi virentes lineolis rufis, apice subfulvae anticis immaculatis, posticis strigis duabus e punctis coeruleis.

*Protesilaus.* 56. P. E. A. alis caudatis subconcoloribus albis, fasciis fuscis, vnica subtus sanguinea, angulo ani rubro. *Syst. Ent.* 450. 36. *Linn. Syst. Nat.* 2. 752. 39. *Mus. Lud. Vlr.* 209.

*Papilio Protesilaus. Clerk. Icon.* tab. 27. fig. 2.

*Papilio caudatus fuscus striis pallescentibus, linea et maculis sanguineis subtus notatus. Petiv. Mus.* 50 n. 502.

*Cram. Inf.* 17. tab. 202. fig. A. B.

*Sulz. Hist. Inf.* tab. 14. fig. 5.

*Sloan. Iam.* 2. 218. tab. 239. fig. 17. 18.

*Seb. Mus.* 1. tab. 11. fig. 2.

— — 4. tab. 36. fig. 11. 12.

*Catesb. Carol.* 2. tab. 100.

*Merian. Surin.* tab. 43.

*Aubent. miscell.* tab. 44. fig. 1. 2.

Habitat in America meridionali.

*Larva spinosa, nigra.*

*Conf. Papilio Marcellus. Cram. Inf.* 2. tab. 98. fig. F. G.

57. P. E. A. alis caudatis concoloribus flavis, fasciis marginique nigris, caudis albis longitudine alae. *Syst. Antilocus.*  
*Ent.* 451. 37. *Linn. Syst. Nat.* 2. 751. 35. *Mus. Lud.*  
*Vlr.* 207.

*Papilio* caudatus maximus striis vmbrique nigris.  
*Petiv. Mus.* 50. n. 505.

*Catesb. Carol.* 2. tab. 83.

Habitat in America.

58. P. E. A. alis caudatis subconcoloribus flavescentibus, fasciis fuscis geminatis, posticis subtus linea sanguinea. *Syst. Podalirius.*  
*Ent.* 451. 38. *Linn. Syst. Nat.* 2. 751. 36.  
*Mus. Lud. Vlr.* 208.

*Papilio* alis pallide flavis, riuulis transversis nigris secundariis angulo subulato maculaque crocea. *Geoff.*  
*Inf.* 2. 56. 24.

*Papilio* alis amplissimis pallidius flavicantibus. *Raj.*  
*Inf.* 111. 3.

*Papilio* *Sinon.* *Pod. Inf.* 62. tab. 2. fig. 1.

*Cram. Inf.* 13. tab. 152. fig. B.

*Esp. pap.* 1. tab. 1. fig. 2.

*Bergstr. Inf.* 2. tab. 18. fig. 1. 2. 3. 4.

*Merian. Europ.* 163. tab. 44.

*Reaum. Inf.* 1. tab. 11. fig. 3. 4.

*Roes. Inf.* 1. pap. 2. tab. 2.

*Schaeff. elem.* tab. 94. fig. 4.

— — *Icon.* tab. 45. fig. 3. 4.

*Mouff. Pap. diurn.* 3. 99. fig. 3.

*Ionst. Inf.* tab. 5. fig. 5.

Habitat in Europae Brasilia.

*Larva* solitaria, flavescent, fusco punctata, capite pallide virecente.

*Puppa* flavescent, fusco punctata, antice bidentata.

59. P. E. A. alis caudatis nigris viridi fasciatis maculatisque, posticis subtus linea sanguinea. *Syst. Sinon.*  
*Ent.* 452. 39.

*Papilio* *Protesilaus.* *Drury Inf.* 1. tab. 22. fig. 3. 4.

*Papilio* *Policenes.* *Cram. Inf.* 4. tab. 37. fig. A. B.

*Aubent. miscell.* tab. 18. fig. 1. 2.

Habitat in India.

16 CLASS. VI. GLOSSATA. *Papilio*.

- Chiron*. 60. P. E. A. alis caudatis nigris fusco fasciatis, subtus antice albis, strigis ferrugineis. *Syst. Ent.* 452. 40.\*  
*Papilio Murius*. *Cram. Inf.* 17. tab. 200. fig. D. E.  
*Aubent. miscell.* 1. tab. 71. fig. 2. 3. ?  
*Seb. Mus.* 4. tab. 34. fig. 13. 14.  
 — — — tab. 35. fig. 3. 4.  
 Habitat in India. *Dom. Drury*.
- Crithon*. 61. P. E. A. alis caudatis supra fuscis, subtus glaucis, fascia vtrinque alba. *Gen. Inf. Mant.* 252.\*  
 Habitat Surinami. *D. Schulz*.  
 Praecedenti affinis.
- Orsilochus*. 62. P. E. A. alis caudatis, anticis fasciis duabus, posticis vnica albis, subtus basi albis, fascia fulua. *Gen. Inf. Mant.* 252.\*  
*Papilio Cinna*. *Cram. Inf.* 17. tab. 200. fig. F. G.  
 Habitat Surinami. *D. Schulz*.
- Faunus*. 63. P. E. A. alis caudatis hyalinis, margine strigaeque nigris. *Syst. Ent.* 532. 380.\*  
*Papilio Chorineus*. *Cram. Inf.* 5. tab. 59. fig. A.  
 Habitat Surinami. *Mus. Hagense*.
- Licarfis*. 64. P. E. A. alis caudatis atris: fasciis duabus albis, angulo ani rubro bimaculato. *Syst. Ent.* 532. 381.\*  
*Papilio Rhetus*. *Cram. Inf.* 6. tab. 63. fig. 6. ?  
 Habitat Surinami. *D. Schulz*.  
 Conf. *Papilio Periander*. *Cram. Inf.* 16. tab. 188. fig. C.
- Meliboeus*. 65. P. E. A. alis caudatis fuscis, anticis striga, posticis duabus sanguineis, subtus cyaneo nitidissimis. *Gen. Inf. Mant.* 252.\*  
*Papilio Pretus*. *Cram. Inf.* 12. tab. 144. fig. A. B.  
 Habitat Surinami. *D. Schulz*.  
 Conf. *Papilio Tedeia*. *Cram. Inf.* 9. tab. 181. fig. A.
- Turnus*. 66. P. E. A. alis caudatis concoloribus flauis, margine fasciisque abbreviatis nigris, angulo ani fuluo. *Syst. Ent.* 452. 41.

*Papilio Turnus* E. A. alis caudatis concoloribus flauis, prioribus fasciis quinque dimidiatis posticeque nigris. *Linn. Mant.* 1. 536.

*Papilio* diurna prima omnium maxima. *Raj. Inf.* 111.

*Papilio Alcidas*. *Cram. Inf.* 4. tab. 38. fig. A. B.

*Catesb. carol.* tab. 97.

Habitat in America.

Statura *P. Machaonis*, at triplo maior.

67. *P. E. A.* alis caudatis concoloribus flauis, limbo fusco *Machaon*. lunulis flauis, angulo ani fuluo. *Syst. Ent.* 452. 42.

*Linn. Syst. Nat.* 2. 750. 93. *Fn. Sv.* 1031.

*Papilio* alis flauo nigroque variegatis, secundariis angulic subulato maculaque fulua. *Geoff. Inf.* 2. 54. 23.

*Papilio* alis amplissimis, flauicante et nigro coloribus variegatis. *Raj. Inf.* 110. 1.

*Papilio* diurnus maximus secundus. *Merr. pin.* 192.

*Papilio* *Erucae rutaceae*. *Column. scph.* 2. tab. 86.

*Eruca Anethi*. *Frisch. Inf.* 2. tab. 10.

*Papilio* maior caudatus ex nigro et luteo variegatus.

*Petiv. Mus.* 35. 322.

*Scop. carn.* 444.

*Gronov. Zooph.* 725.

*Esp. Papil.* tab. 1. fig. 1.

*Bergstr. Inf.* 2. tab. 17. fig. 3. 4.

*Mouff. Inf.* 99. tab. 968.

*Aldrov. Inf.* 96. n. 5. 6. fig. 1.

*Ionst. Inf.* tab. 5. 7.

*Hoeffn. Inf.* 1. tab. 12.

*Merian. Europ.* tab. 94.

*Reaum. Inf.* 1. tab. 29. fig. 1.

*Edw. Av.* tab. 343.

*Roes. Inf.* 1. pap. 2. tab. 1.

*Wilk. pap.* 47. tab. 1. a. 1.

*Schaeff. Icon.* tab. 45. fig. 1. 2.

*Seb. Mus.* 4. tab. 32. fig. 9. 10.

— — — tab. 59. fig. 12. 13.

Habitat in Europae umbellatis, ruta.

*Larva* solitaria, glabra, tentaculata, nigro viridique annulata rubroque punctata.

*Puppa* flauescens.

*Fabr. Sp. Inf. T. II.*

B

68. P.

*Thersites*. 68. P. E. A. alis caudatis flavis, limbo nigro posticis lunulis flavis. *Syst. Ent.* 453. 43.\*

Habitat in America. *Mus. D. Hunter*.

*Codrus*. 69. P. E. A. alis caudatis supra atris subtus fuscis, anticis vtrinque fascia maculari alba.

*Papilio Codrus*. *Cram. Inf.* 15. tab. 179. fig. A. B.

Habitat in Amboina.

Magnus. Alae omnes supra atrae viridi colore resplendentes ad marginem interiorem virescentes atomis ferrugineis. Subtus fuscae nigro nervosae. Fascia e maculis octo albidis vtrinque in alis anticis.

*Chalcus*. 70. P. E. A. alis caudatis nigris, fasciis duabus macularibus flavis, posticis subtus vitta flava lunulisque rufis. *Syst. Ent.* 453. 44.\*

*Papilio Palamedes*. *Drury Inf.* 1. tab. 19. fig. 12.

*Papilio Palamedes*. *Cram. Inf.* 8. tab. 93. fig. A. B.

Habitat in America. *Mus. D. Hunter*.

*Decius*. 71. P. E. A. alis caudatis fascia communi alba, posticis apice rufis, ocellis punctisque duobus albis.

*Papilio Decius*. *Cram. Inf.* 10. tab. 114. fig. A. B.

Habitat in Guinea.

Alae omnes basi griseae, apice nigrae striga maculari ferruginea. In medio fascia communis alba. Posticae apice ferrugineae, ocellis duobus atris, pupilla coerulea et ad angulum ani puncta duo alba. Subtus basi ferrugineae lunulis coeruleis atro adnatis, apice variegatae posticis ocellis septem duobus maioribus et ultimo pupilla duplici.

*Doreus*. 72. P. E. A. alis caudatis fascia supra viridi subtus alba. *Syst. Ent.* 457. 62.\*

Habitat in Sierra Leon Africae. *Mus. Dom. Banks*.

*Palamedes*. 73. P. E. A. alis caudatis nigris, lunulis marginalibus flavis, anticis fascia flava, posticis maculari rufa. *Syst. Ent.* 454. 45.\*

Habitat in America. *Mus. D. Hunter*.

*Meneſtheus*. 74. P. E. A. alis caudatis nigris, subtus albido striato maculatis, posticis lunulis rufis coeruleſcentibusque. *Syst. Ent.* 454. 46.\*

*Papilio Meneſtheus*. *Drury Inf.* 2. tab. 9. fig. 1. 2.

*Papilio*



CLASS. VI. GLOSSATA. Papilio. 19

*Papilio Menestheus*. *Cram. Inf.* 12. tab. 142. fig. A. B.

Habitat in India.

Maior P. Xutho.

75. P. E. A. alis caudatis nigris albo striato maculatis, posticis subtus ocellis coeruleis fulvisque subfasciatis. *Xuthus*.

*Syst. Ent.* 454. 47. *Linn. Syst. Nat.* 2. 751. 34.

*Papilio Xuthus*. *Cram. Inf.* 7. tab. 73. fig. A. B.

Habitat in India orientali.

76. P. E. A. alis caudatis nigris flavo fasciatis, posticis subtus flavis, fascia nigra, lunulis cyaneis. *Thoas*.

*Syst. Ent.* 454. 48.

*Papilio Thoas* alis caudatis, supra nigris fasciis duabus flavis interruptis, subtus flavis, coerulca. *Linn.*

*Mant.* 1. 536.

*Papilio Thoas*. *Drury Inf.* 1. tab. 22. fig. 1. 2.

*Papilio Thoas*. *Cram. Inf.* 19. tab. 167. fig. A. B.

*Papilio Cresphontes*. *Cram. Inf.* 14. tab. 165. fig. A.

— — — — — tab. 166. fig. B.

*Aubert. miscell.* 1. tab. 69.

*Seb. Mus.* 4. tab. 38. fig. 5. 6. 7.

Habitat in America meridionali.

77. P. E. A. alis caudatis atrae fascia viridi, posticis sub-Cresphontes.  
tus fascia coerulea.

*Papilio Demolion*. *Cram. Inf.* 8. tab. 89. fig. A. B.

Habitat in India orientali.

Præcedente minor. Alae omnes supra atrae fascia viridi anticarum maculari. Lunulae quinque virides ante marginem alarum posticarum. Subtus anticae atrae striis quatuor bascos fasciaque maculari viridibus. Posticae fasciis atris viridibusque alternis mediaque coerulea.

78. P. E. A. alis dentato caudatis nigris fascia flava, posticis subtus lunulis fulvis coeruleis flavisque. *Polycaon*.

*Papilio Polycaon*. *Cram. Inf.* 17. tab. 203. fig. A. B.

*Klemann Inf.* 1. tab. 7. fig. 1. 2.

*Mercari Swr.* tab. 31.

Habitat in Althacis Surinami.

Affinis videtur P. Thoadi. Alae posticae supra strigis duabus lunularum virentium, at obsoletiorum, quae ad angulum ani macula coeruleo-rufa, ad marginem interiorem macula flava terminantur.

- Ajax.* 79. P. E. A. alis obtuse caudatis concoloribus fuscis, fasciis flavescentibus, angulo ani fulvo. *Syst. Ent.* 455. 49. *Linn. Syst. Nat.* 2. 750. 32.  
*Raj. Inf.* 111. 2.  
*Edw. Av.* 34. tab. 34.  
 Habitat in America boreali.
- Orontes.* 80. P. E. A. alis caudatis nigris, fasciis duabus virefcentibus, caudis albis distantibus. *Syst. Ent.* 455. 50. *Linn. Syst. Nat.* 2. 750. 27. *Amoen. acad.* 6. 402. 51.  
*Papilio Orontes. Clerk. Inf.* tab. 26. fig. 1.  
*Cram. Inf.* 7. tab. 83. fig. A. B.  
*Seb. Mus.* 4. tab. 16. fig. 1. 2.  
 — — — tab. 47. fig. 7. 8.  
 Habitat in India.  
 Antennae obtusae, filiformes.
- Agamemnon.* 81. P. E. A. alis caudatis nigris viridi maculatis, posticis subtus lunulis tribus rufis. *Syst. Ent.* 455. 51. *Linn. Syst. Nat.* 2. 748. 22. *Mus. Lud. Vlr.* 202.  
*Papilio Aegisteus. Cram. Inf.* 9. tab. 106. fig. C. D.  
 Habitat in Asia.
- Aegisthus.* 82. P. E. A. alis caudatis fusco luteoque vndulatis, ocellis caudalibus subternis.  
*Phalaena Lunus. Attacus feticornis spirilinguis*, alis caudatis fusco luteoque vndatis, ocellis caudalibus subternis. *Linn. Syst. Nat.* 2. 810. 6. *Mus. Lud. Vlr.* 371. *Clerk. Icon.* tab. 52. fig. 2.  
*Papilio Lunus. Cram. Inf.* 17. tab. 200. fig. A. B. C.  
 Habitat in Indiis.  
 Antennae filiformes vti in Oronte, Leilo, Phidippo.  
 Conf. *Papilio Empedocles. Cram. Inf.* 17. tab. 199. fig. A. B.
- Euryalus.* 83. P. E. A. alis caudatis nigris, posticis ante marginem coeruleis, maculis nigris, pupilla alba.  
*Papilio Euryalus. Cram. Inf.* 7. tab. 74. fig. A. B.  
 Habitat in Amboina.  
 Alae anticae nigrae fascia fusca, posticae ante marginem coerulefcentes maculis septem atris pupilla alba.  
 Subtus

CLASS. VI. GLOSSATA. Papilio. 21

Subtus omnes fusco rubro coeruloque variegatae fascia flava distincta.

84. P. E. A. alis caudatis nigris, anticis fascia fulva, posticis alba submarginali. *Nifus*.  
*Papilio Nifus*. *Cram. Inf.* 13. tab. 150. fig. A. B.  
 Habitat in Amboina.  
 Magnus. Alae omnes supra atrae. Anticae fasciâ e maculis magnis altero latere emarginatis fulvis. Posticae fascia e lunulis cyaneis, tunc striga e punctis albis et tandem fascia submarginali alba. Subtus omnes variegatae.
85. P. E. A. alis subcaudatis fuscis, subtus fasciis albis biocellatis caudaeque ocellis geminis. *Phidippus*.  
*Syst. Ent.* 455. 52. *pus*.  
*Linn. Syst. Nat.* 2. 752. 37. *Amoen. acad.* 6. 402. 52.  
*Papilio Phidippus*. *Cram. Inf.* 6. tab. 69. fig. A. B.  
*Seb. Musf.* 4. tab. 5. fig. 5.  
 Habitat in India.  
 Apex alae ferè in caudam obtusam definit ocellis duobus coecis.
86. P. E. A. alis subcaudatis fuscis apice atris albo maculatis, posticis subtus ocellis duobus. *Aurelius*.  
*Papilio Aurelius*. *Cram. Inf.* 14. tab. 168. fig. A. B.  
 Habitat in India orientali.  
 Statura P. Phidippi. Alae subtus albae fasciis fuscis.
87. P. E. A. alis dentatis nigris flavo maculatis, posticis ocello coeruleo rufoque. *Demoleus*.  
*Syst. Ent.* 455. 53.  
*Papilio Demoleus* alis dentatis fuscis, maculis fasciaeque maculosa flavis, posticis ocellis binis. *Linn. Syst. Nat.* 2. 753. 46. *Musf. Lud. Vlr.* 214.  
*Kleman. Inf.* 1. tab. 1. fig. 2. 3.  
*Ehret. pict.* tab. 5.  
*Seb. Musf.* 4. tab. 37. fig. 17. 18.  
 — — — tab. 44. fig. 6-9.  
 Habitat in Citro Indiae orientalis. *D. Koenig*.
88. P. E. A. alis dentatis supra nigris fascia coerulea, subtus fuscis ocellis tribus quinisue. *Achilles*.  
*Syst. Ent.* 456.  
 54. *Linn. Syst. Nat.* 2. 752. 42. *Musf. Lud. Vlr.* 211.  
 B 3 *Papilio*

*Papilio Achilles.* Clerk. Icon. tab. 24. fig. 2.

*Papilio Achilles.* Cram. Inf. 3. tab. 27. fig. A. B. ♀

*Papilio Achilles.* Cram. Inf. 3. tab. 28. fig. A. ♂

Merian. Sur. tab. 7.

Seb. Musf. 4. tab. 24. fig. 1. 2.

Gronov. Zooph. 719.

Knorr. delic. tab. C. 2. fig. 1. 2.

Habitat in Pfdio Americes.

Larva pilosa, flava.

Puppa obesa viridis.

Variat ocellis tantum quatuor alae posticae.

Conf. *Papilio Helenor.* Cram. Inf. 8. tab. 86. fig. A. B.  
vix satis distinctus.

*Medon.* 89. P. E. A. alis supra nigris, anticis fascia lutea apiceque albo, posticis disco coerulescentibus. *Syst. Ent.* 456.  
55. *Linn. Syst. Nat.* 2. 753. 43. *Amoen. acad.* 6.  
402. 53.

Clerk. Icon. tab. 28. fig. 1.

Drury Inf. 2. tab. 15. fig. 1. 2.

Cram. Inf. 17. tab. 205. fig. B. C.

Habitat in Indiis.

Alae subtus cinerascetes punctis baseos tribus nigris.

*Philotes.* 90. P. E. A. alis dentatis fuscis, posticis ocellis duobus, pupilla nigra punctisque tribus albis. *Syst. Ent.* 456. 56. *Linn. Syst. Nat.* 2. 750. 29. *Musf. Lud.* Vlr. 219.

Drury Inf. 2. tab. 1. fig. 1. 2.

Cram. Inf. 2. tab. 20. fig. A. B. C.

Clerk. Icon. tab. 30. fig. 3.

Sulz. Hist. Inf. tab. 13. fig. 5.

Seb. Musf. 4. tab. 4. fig. 9. 10.

— — — tab. 21. fig. 7. 8.

Habitat in Indiis.

Alae anticae subtus fascia lineari alba lineaque transuersa atra.

*Lucretius.* 91. P. E. A. alis dentatis atris fuluo fasciatis, subtus maculis atris linea coerulea cinctis.

*Papilio Lucretius.* Cram. Inf. 7. tab. 82. fig. E. F.

Habi-

CLASS. VI. GLOSSATA. Papilio, 29

Habitat in Guinea.

Antennae extrorsum crassiores.

98. P. E. A. alis dentatis fuscis, fasciis duabus viridibus *Stelenes*.  
altera posticarum integra. *Syst. Ent.* 456. 57.

*Papilio Stelenes* alis subcaudatis supra fuscis, fascia vi-  
rescente obtusa interrupta. *Linn. Syst. Nat.* 2. 750.  
30. *Mus. Lud. Vlr.* 218.

*Papilio Stelenes*. *Clerk. Icon.* tab. 35. fig. 2.

*Papilio Jamaicensis*, e fusco lutescens, subtus virescens  
croceo mixtus. *Petiv. Gazoph.* 20. tab. 13. fig. 1.

*Sloan. Jam.* 2. 217. tab. 239. fig. 9. 10.

*Cram. Inf.* 7. tab. 79. fig. A. B.

*Aubert. miscell.* tab. 95 fig. 1. 2.

*Merian. Sur.* tab. 11.

Habitat in America meridionali.

Larva viridis stria alba coccineaque, cauda spinis  
duabus.

99. P. E. A. alis dentatis nigris fascia inaurato viridi, sub- *Nireus*.  
tus nigricantibus. *Syst. Ent.* 457. 58. *Linn. Syst.*  
*Nat.* 2. 750. 28. *Mus. Lud. Vlr.* 207.

*Papilio Nireus*. *Clerk. Icon.* tab. 30. fig. 1.

*Papilio Nireus*. *Cram. Inf.* 16. tab. 187. fig. A. B.

*Drury Inf.* 2. tab. 4. fig. 1. 2.

*Seb. Mus.* 4. tab. 9. fig. 21. 22.

— — — tab. 6. fig. 21. 22.

Habitat in Indiis.

Alae posticae subtus striga maculari alba.

94. P. E. A. alis dentatis atris, fascia coerulea nitida, sub- *Amphi*-  
tus antice albis postice cinereis. *Syst. Ent.* 457. 59.\* *machus*.

*Sulz. Hist. Inf.* tab. 14. fig. 23.

Habitat in Indiis. *Mus. Lugdunense*.

95. P. E. A. alis fulcatis subdentatis fuscis, anticis disco *Odius*.  
flavo. *Syst. Ent.* 457. 60.\*

*Papilio Odius*. *Sulz. Hist. Inf.* tab. 13. fig. 2.

Habitat in Indiis. *Mus. Lugdunense*.

24 CLASS. VI. GLOSSATA. *Papilio*

*Amphitrion*. 96. P. E. A. alis dentatis nigris fascia inaequali flava, posticis subtus striga e punctis flavis lunulisque cyaneis.  
*Gen. Inf. Mant.* 253. \*

*Papilio Amphitrion*. *Cram. Inf.* 1. tab. 7. fig. A. B.

*Papilio Gambrius*. *Cram. Inf.* 13. tab. 157. fig. A. B.

*Seb. Mus.* 4. tab. 8. fig. 7. 8.

Habitat in America.

*Demonophon*. 97. P. E. A. alis dentatis supra nigris fascia viridi, subtus griseo marmoratis. *Syst. Ent.* 457. 61. *Linn. Syst. Nat.* 2. 753. 47. *Mus. Lud. Vr.* 215.

*Papilio Peridamas*. *Cram. Inf.* 13. tab. 158. fig. A. B.  
C. D. E.

*Papilio lufuosus*. *Naturf.* 7. tab. 1. fig. 1.

*Roef. Inf.* 4. tab. 6. fig. 1.

*Clerk. Icon.* tab. 29. fig. 2.

*Seb. Mus.* 4. tab. 18. fig. 2. 3.

— — — tab. 26. fig. 11 — 15.

Habitat in Indiis.

*Conf. Papilio Meander*. *Cram. Inf.* 2. tab. 12. fig. A. B.

*Nestor*. 98. P. E. A. alis dentatis supra fuscis albo maculatis discoque coeruleo, subtus ocellis tribus quatuorque. *Syst. Ent.* 458. 63. *Linn. Syst. Nat.* 2. 752. 40.

*Gron. Zooph.* 718.

*Cram. Inf.* 2. tab. 19. fig. A. B.

*Seb. Mus.* 4. tab. 43. fig. 23. 24.

*Merian. Sur.* 9. tab. 9.

Habitat in Punica America.

Numerus ocellorum variat.

*Perseus*. 99. P. E. A. alis dentatis pallide coeruleis apice atris ferrugineo maculatis, subtus undatis ocellis tribus quatuorque.

*Papilio Perseus*. *Cram. Inf.* 6. tab. 71. fig. A. B.

*Seb. Mus.* 4. tab. 18. fig. 15. 16.

— — — tab. 17. fig. 13. 14.

Habitat Surinami.

Nimis affinis P. Nestori.

**CLASS. VI. GLOSSATA. Papilio.** 29

100. P. E. A. alis dentatis fuscis, area communi coerulea *Telema-*  
radiata, singulis subtus ocellis sex. *Syst. Ent.* 458. *chus.*  
64. *Linn. Syst. Nat.* 2. 752. 41.  
*Merian. Sur.* tab. 68.  
Habitat in America meridionali.
101. P. E. A. alis dentatis supra coeruleis nitidissimis, *Menelaus.*  
subtus nebulosis punctis fuscis. *Syst. Ent.* 458. 65.  
*Linn. Syst. Nat.* 2. 748. 20. *Mus. Lud. Vlr.* 200.  
*Clerk. Icon.* tab. 21. fig. 1.  
*Merian. Sur.* tab. 53.  
*Knorr. Delic.* tab. C. 4. fig. 2.  
*Seb. Mus.* 4. tab. 31. fig. 1. 2.  
*Cram. Inf.* 2. tab. 21. fig. A. B.  
Habitat in America meridionali.  
*Larva* spinosa flava, striis pedibusque roseis, capite  
brunneo.  
*Puppa* pallida, cauda cylindrica inflexa.  
*Conf. Papilio Rhetenor. Cram. Inf.* 2. tab. 15. fig.  
A. B.  
*Sulz. Hist. Inf.* tab. 13. fig. 7.
102. P. E. A. alis subrepandis lividiusculis subtus nebu- *Teucer.*  
lofis, posticis ocellis tribus, postremo maximo. *Syst.*  
*Ent.* 458. 66. *Linn. Syst. Nat.* 2. 753. 44. *Mus.*  
*Lud. Vlr.* 212.  
*Papilio Teucer. Cram. Inf.* 5. tab. 51. fig. A. B.  
*Papilio Eurylochus. Cram. Inf.* 3. tab. 33. fig. A.  
— — — — — tab. 34. fig. A.  
*Sloan. Iam.* 2. 219. 24.  
*Knorr. Delic.* tab. C. 1. fig. 1. 2.?  
*Sulz. Hist. Inf.* tab. 14. fig. 1.  
*Merian. Sur.* tab. 23.  
*Seb. Mus.* 4. tab. 31. fig. 3. 4.  
Habitat in MUSA Americæ.  
*Larva* brunnea dorso quadrispinosa, capite spinosa,  
cauda bifida.  
*Puppa* pallescens maculis vtrinque duabus argenteis.
103. P. E. A. alis angulato repandis fuscis subtus lividiu- *Automa-*  
sculis, ocello anguli ani. *Gen. Inf. Mant.* 253. \* *don.*



*Papilio Automedon*. Cram. Inf. 4. tab. 41. fig. A. B.  
Habitat Surinami.

*Hecuba*. 104. P. E. A. alis dentatis, anticis rubris posticis nigris,  
subtus vtrunque ocellatis. *Syst. Ent.* 459. 67. *Linn.*  
*Mant.* 1534.

*Aubent. miscell.* tab. 19.

Habitat in Caienna.

*Idomeneus*. 105. P. E. A. alis subcrenatis fuscis basi cyaneis subtus ne-  
bulosis, posticis oculo magno flavescente. *Syst. Ent.*  
459. 68.

*Papilio Idomeneus* alis subcrenatis coerulescenti fusce-  
scentibus, subtus nebulosis, ocellis duobus. *Linn.*  
*Syst. Nat.* 2. 753. 45. *Mus. Lud. Vlr.* 213.

*Papilio surinamensis* maximus subtus perbelle oculatus  
et marmoratus. *Petiv. Gazoph.* 43. tab. 28. fig. 1.

*Papilio Idomeneus*. Cram. Inf. 5. tab. 52. fig. B.

*Papilio Ilioneus*. Cram. Inf. 5. tab. 52. fig. A.

*Clerk. Icon.* tab. 20. fig. 1.

*Gron. Zooph.* 721.

*Merian. Sur.* 60. tab. 60.

*Aubent. miscell.* 1. tab. 96. fig. 1. 2.

*Seb. Mus.* 4. tab. 24. fig. 7. 8.

— — tab. 43. fig. 15. 16.

Habitat in America meridionali.

*Larva* rufa verrucis plumosis verticillata.

*Puppa* dentata, apice stylis duobus incurvis.

*Eurypy- 106. P. E. A. alis dentatis concoloribus nigris, fascia in-*  
*lus.* — — — — — interrupta maculisque viridibus, posticis subtus rubro  
maculatis. *Linn. Syst. Nat.* 2. 754. 49. *Mus. Lud.*  
*Vlr.* 216. *Clerk. Icon.* tab. 28. fig. 2.

*Cram. Inf.* 11. tab. 122. fig. C. D.

*Seb. Mus.* 4. tab. 26. fig. 17 — 20. tab. 45. fig.  
23. 24.

Habitat in Indiis.

Nimis affinis P. Sarpedoni.

\*\*\* Heliconii.

107. P. H. alis oblongis integerrimis luteis, anticis fribus *Calliope*  
tribus, posticis fasciis tribus nigris. *Syst. Ent.* 459.  
70. *Linn. Syst. Nat.* 2. 755. 56. *Mus. Lud. Vlr.*  
223.

*Clerk. Icon.* tab. 41. fig. 4.

*Seb. Musf.* 4. tab. 7. fig. 5. 6.

— — tab. 29. fig. 17. 18.

Habitat in Indiis.

Apex alae anticae niger albo punctatus.

108. P. H. alis oblongis integerrimis flauis, anticis supra *Melite*  
nigris, lineis duabus fasciaeque flauis. *Syst. Ent.* 460.  
71. *Linn. Syst. Nat.* 2. 755. 57. *Amoen. acad.* 6.  
403. 56.

*Papilio Melite. Cram. Inf.* 13. tab. 153. fig. C. D.

*Clerk. Icon.* tab. 44. fig. 5.

*Seb. Musf.* 4. tab. 17. fig. 8. 9.

— — tab. 20. fig. 8. 9.

— — tab. 30. fig. 5. 6.

Habitat in Indiis.

Alae posticae flavae margine postico nigro.

109. P. H. alis oblongis concoloribus fulvis apice nigris *Mopsa*  
albo punctatis. *Syst. Ent.* 460. 72.

*Papilio Mopsa* alis oblongis concoloribus luteis nigris-  
que, subtus punctis marginalibus octo albis. *Linn.*  
*Syst. Nat.* 2. 756. 59. *β. Musf. Lud. Vlr.* 235.

*Papilio Harmonia. Cram. Inf.* 16. tab. 190. fig. D.

*Seligm. Av.* 6. tab. 70.

*Seb. Musf.* 4. tab. 29. fig. 17. 18.

Habitat in India orientali.

Variat rarius punctis albis tantum in pagina inferiori.

110. P. H. alis oblongis integerrimis fulvis nigro macula- *Mneme*  
tis, anticis apice atris flauo maculatis.

*Papilio Mneme* alis oblongis integerrimis nigris basi  
fulvis, primoribus fascia maculisque tribus flauis.  
*Linn. Syst. Nat.* 2. 756. 59. *Amoen. acad.* 6. 403. 5.

*Papilio Mneme. Cram. Inf.* 16. tab. 190. fig. 5.

*Sulz. Hist. Inf.* tab. 15. fig. 3. 4.

Habitat

28 CLASS. VI. GLOSSATA. Papilio.

Habitat in India orientali.

Praecedenti valde affinis, attamen distinctus videtur.

- Lybia.** 111. P. H. alis oblongis integerrimis, anticis atris, fasciis duabus fulvis, posticis fulvis, fascia marginali atra. *Papilio Lybia* alis oblongis integerrimis fulvis, anticis strigis tribus, posticis fascia marginali fuscis. *Syst. Ent.* 460. 73. \*

*Papilio Hypsipyle. Cram. Inf.* 15. tab. 177. fig. C. D.  
Habitat in India. *D. Solander.*

- Viola.** 112. P. H. alis oblongis integerrimis fulvis nigro punctatis, posticis margine nigro, punctis albis. *Syst. Ent.* 460. 74. \*

*Papilio Cepheus. Salz. Hist. Inf.* tab. 15. fig. 2.  
Habitat in Viola, Boragine Indiae. *D. Koenig.*

- Terpsichors.** 113. P. H. alis oblongis integerrimis fulvis, posticis nigro punctatis. *Syst. Ent.* 460. 75. *Linn. Syst. Nat.* 2. 755. 55. *Mus. Lud. Vlr.* 222.

*Papilio inadera spatani croceus, nigris maculis adspersus. Petiv. Gazoph.* tab. 40. fig. 4.

Habitat in Asia.

Apex alae anticae albicat.

- Serena.** 114. P. H. alis oblongis integerrimis fulvis, posticis subtus basi nigro punctatis. *Syst. Ent.* 461. 76. \*

Habitat in India. *Mus. Dom. Comit. Thot.*

- Polymnia.** 115. P. H. alis oblongis integerrimis, anticis fascia flava, posticis fasciis tribus nigris. *Syst. Ent.* 461. 77. *Linn. Syst. Nat.* 2. 755. 58. *Mus. Lud. Vlr.* 224.

*Papilio surinamensis ex aurantio et nigro mixtus. Petiv. Gazoph.* tab. 12. fig. 8.

*Papilio Polymnia. Cram. Inf.* 16. tab. 191. fig. E.

*Roef. Inf.* 4. tab. 5. fig. 2.

*Edw. Av.* 175. tab. 175.

*Seb. Mus.* 4. tab. 37. fig. 15. 16.

— — — — — tab. 30. fig. 23. 24.

Habitat in America meridionali.

*Larva* flaescens sub stigmatibus spinosa.

P. H. alis oblongis integerrimis, anticis nigris, fasciis tribus hyalinis, posticis hyalinis. *Syst. Ent. 461. 78. \* Hippodamia.*

Habitat — — — — Mauduit.

7. P. H. alis oblongis integerrimis fuscis disco albis, posticis ocellis vtrinque binis. *Syst. Ent. 461. 79. Linn. Vrania.*

*Syst. Nat. 2. 756. 60. Mus. Lud. Vlr. 225.*

Habitat in Indiis.

Ocelli duo magni flavi iride coerulescente pupilla alba.

P. H. alis oblongis integerrimis atris, anticis flavo maculatis, posticis supra basi coeruleis. *Syst. Ent. 461. 80. Linn. Mant. 1. 536. Doris.*

*Papilio Quirina. Cram. Inf. 6. tab. 65. fig. A. B.*

*Aubert. miscell. tab. 72. fig. 1. 2.*

Habitat Surinami.

Forma affinis P. Ricini.

P. H. alis oblongis integerrimis nigris, anticis fascia, posticis subtus punctis marginalibus albis. *Gen. Inf. Mant. 254. \* Hecale.*

*Papilio Pasithoe. Cram. Inf. 2. tab. 17. fig. A. B.*

Habitat Surinami, D. Schulz.

P. H. alis oblongis integerrimis fuscis, anticis vtrinque fasciis duabus flavis, posticis basi radiatis. *Syst. Ent. 461. 81. Linn. Syst. Nat. 2. 756. 63. Mus. Lud. Vlr. 227. Ricini.*

*Papilio Rhea. Cram. Inf. 5. tab. 54. fig. C. D.*

*Papilio Amathusia. Cram. Inf. 15. tab. 177. fig. F.*

*Gram. Zooph. 740. 741.*

*Ehret. pict. tab. 1.*

*Roef. Inf. 4. tab. 5. fig. 1.*


*Merian. Sur. tab. 30.*

*Seb. Mus. 4. tab. 29. fig. 3. 4.*

Habitat in Americae Ricino.

Forma virescens, pilosa pilis albis.

Forma alis posticis basi purpureo, femina coeruleo radiatis.

- Brassolis.** 121. P. H. alis oblongis integris fuscis, anticis flauo maculatis, posticis subtus sanguineo radiatis. *Gen. Inf. Mant.* 254. \*  
*Papilio Bellona. Cram. Inf.* 2. tab. 13. fig. E. F.   
 Habitat Surinami.
- Thales.** 122. P. H. alis oblongis subintegris subconcoloribus atris basi rubro radiatis, anticis flauo maculatis. *Gen. Inf. Mant.* 255. \*  
*Papilio Thales. Cram. Inf.* 4. tab. 38. fig. C. D.  
*Aubent. miscell.* tab. 72. fig. 3. 4.  
 Habitat Surinami.  
 P. Erato nimis affinis.  
 Conf. *Papilio Cybele. Cram. Inf.* 16. tab. 188. fig. A.
- Myrti.** 123. P. H. alis oblongis integerrimis atris, anticis fasciis duabus flauis, posticis disco rufo. *Syst. Ent.* 462. 82. \*  
*Papilio Erycinia. Cram. Inf.* 15. tab. 177. fig. E. ?  
 Habitat Surinami. *Dom. Drury.*
- Psidii.** 124. P. H. alis oblongis integerrimis fuscis, anticis fasciis tribus, posticis duabus hyalinis. *Syst. Ent.* 462. 83.  
*Linn. Syst. Nat.* 2. 756. 64. *Mus. Lud. Vlr.* 228.  
*Phalaena Benghalensis fusco et diaphano mixta. Pet. Gazoph.* tab. 40. fig. 3.  
*Roef. Inf.* 4. tab. 2. fig. 3.  
*Merian. Sur.* tab. 19.  
*Edw. Av.* tab. 80.  
 Habitat in Indiae Psidio.  
*Larua nuda viridis, capite nigro.*
- Aegle.** 125. P. H. alis oblongis integerrimis atris, anticis maculis, posticis disco hyalinis. *Gen. Inf. Mant.* 255. \*  
 Habitat in India. *D. Schulz.*  
 Conf. *Papilio Eurimedia. Cram. Inf.* 11. tab. 126. fig. C. D.
- Charitonia.** 126. P. H. alis oblongis integerrimis atris concoloribus, anticis fasciis flauis tribus, posticis duabus. *Syst. Ent.* 462. 84. *Linn. Syst. Nat.* 2. 757. 65.  
*Papilio Charitonia. Cram. Inf.* 16. tab. 191. fig. F.

*Seligm.*

CLASS. VI. GLOSSATA. Papilio. 31

*Seligm. Av.* 4. tab. 55.

*Aubent. miscell.* tab. 70. fig. 4.

*Sloan. Iam.* 2. 217. tab. 239. fig. 15, 16.

*Edw. Av.* tab. 80.

Habitat in America.

127. P. H. alis oblongis integerrimis nigris, anticis fascia *Melpomene* rubra, posticis basi subtus punctis sanguineis. *Syst. ne.*

*Ent.* 462. 85. *Linn. Syst. Nat.* 2. 752. 71. *Muf.*

*Lud. Vlr.* 232.

*Papilio Melpomene. Cram. Inf.* 16. tab. 191. fig. 6.

*Seligm. Av.* 2. tab. 75.

*Roej. Inf.* 4. tab. 3. fig. 6.

*Sloan. Iam.* 2. 219. tab. 239. fig. 25, 26.

*Edw. Av.* tab. 38.

*Seb. Muf.* 4. tab. 13. fig. 15, 16.

— — tab. 29. fig. 13, 14.

Habitat in America.

CONF. *Papilio Calycopis. Cram. Inf.* 16. tab. 190. fig.

E. F.

128. P. H. alis oblongis integerrimis atris, anticis fascia *Phyllis* rubra, posticis flava. *Syst. Ent.* 463. 86.

*Papilio mexicanus* nigrescens linea lutescente, area miniacca notabili. *Petiv. Gazoph.* 7. tab. 4. fig. 2.

*Papilio Roxana. Cram. Inf.* 4. tab. 45. fig. E. F.

Habitat in Brasilia. *Muf. Dom. Banks.*

129. P. H. alis oblongis integerrimis fuscis, anticis albo *Clio* maculatis, posticis fascia alba. *Syst. Ent.* 463. 87.

*Linn. Syst. Nat.* 2. 757. 66. *Muf. Lud. Vlr.* 229.

*Merian. Sur.* tab. 35. fig. inf.

Habitat in Americas Sophora.

Larva pilosa, brunnea.

130. P. H. alis oblongis integerrimis fuscis, anticis flavo *Thalia* maculatis, posticis striatis. *Syst. Ent.* 463. 88. *Linn.*

*Syst. Nat.* 2. 757. 67. *Muf. Lud. Vlr.* 230.

*Clerk. Icon.* tab. 43. fig. 2.

*Seb. Muf.* 4. tab. 7. fig. 1, 2.

Habitat in Indiis.

Striae alarum posticarum bifidae.

*Euryta*. 131. P. H. alis oblongis integerrimis concoloribus fuscis, fascia alba, posticis striatis basi nigro punctatis. *Syst. Ent.* 463. 89.

*Papilio Euryta* alis oblongis integerrimis fuscis concoloribus nigro striatis fascia alba, posticis punctis decem atris. *Linn. Syst. Nat.* 2. 757. 69. *Mus. Lud. Vlr.* 221.

*Seb. Musf.* 4. tab. 29. fig. 15. 16.

Habitat in Indiis.

*Aedeia*. 132. P. H. alis oblongis integerrimis albo maculatis, anticis virefcentibus, posticis fascia flava. *Syst. Ent.* 463. 90. *Linn. Syst. Nat.* 2. 757. 68. *Amoen. acad.* 6. 403. 55.

*Clerk. Icon.* tab. 4. fig. 2.

Habitat in Indiis.

Alae anticae fasciis tribus e maculis albis.

*Thallo*. 133. P. H. alis oblongis integerrimis nigris, anticis fasciis duabus posticisque unica flavis. *Syst. Ent.* 463. 91. *Linn. Syst. Nat.* 2. 756. 62.

*Edw. Av.* 36. tab. 226.

Habitat in China.

Corpus coeruleum capite rubro.

*Antiocha*. 134. P. H. alis oblongis integerrimis nigris, anticis fasciis duabus albis. *Syst. Ent.* 464. 92. \* *Linn. Syst. Nat.* 2. 1068. 12.

*Papilio Antiochus*. *Cram. Inf.* 4. tab. 38. fig. E. F.

*Papilio Clytia*. *Cram. Inf.* 6. tab. 66. fig. C. D.

*Ehret. pict.* tab. 1.

Habitat in Indiis.

*Camoena*. 135. P. H. alis subintegris fuscis nigro punctatis, posticis fascia rubra. *Syst. Ent.* 464. 93. \*

*Papilio Camoena*. *Drury Inf.* 2. tab. 7. fig. 2.

Habitat in Africa.

Affinis videtur P. Zetes *Linnaei*.

*Gesa*. 136. P. H. alis oblongis integerrimis nigris, anticis maculis duabus, posticis disco fulvo nigro striato, subius basi nigro punctatis.

Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*

Medius.

Medius. Caput et thorax atra albo punctata. Abdomen fuluum maculis dorsalibus nigris. Alae anticae nigrae macula magna ad marginem tenuiorem et minori interdum diuisa ad apicem. Posticae nigrae disco fuluo, nigro striato. Subtus anticae concolores, posticae fuluae, basi nigro punctatae, apice nigro striatae.

Variat interdum basi omni alarum nigro punctata.

137. P. H. alis subintegris nigris, disco communi virescente, subtus nigro punctatis, posticis macula bascos rubra. *Cyparissia*.

*Papilio Cyparissa*. Cram. Inf. 4. tab. 39. fig. D. E.

— — — — — Inf. 13. tab. 156. fig. B.

Habitat in Sierra Leon Africae.

Statura P. Camenae. Macula virescens in alis anticis, subtus omnes virescentes nigro punctatae posticis macula bascos rubra, interdum disco flauescente.

Conf. *Papilio Lucilla*. Cram. Inf. 13. tab. 156. fig. A. C.

138. P. H. alis integerrimis concoloribus fuscis albo maculatis, posticis disco albo punctis nigris. *Lycia*.  
464. 94. \*

Habitat in Sierra Leon Africae. *Mus. Dom. Banks*.

139. P. H. alis oblongis integerrimis concoloribus fuscis flauo maculatis, posticis disco ferrugineo, nigro striato. *Pyrrha*.  
464. 95. \*

*Seb. Mus.* 4. tab. 29. fig. 1. 2.

Habitat in Brasilia. *Mus. Dom. Banks*.

140. P. H. alis oblongis integerrimis fuscis albo variis maculis margineque atris. *Pasphe*.

*Papilio Medea*. Cram. Inf. 7. tab. 31. fig. C. D.

Habitat in Guinea.

Alae posticae subtus basi flauescientes maculis atris.

141. P. H. alis oblongis integerrimis, anticis fuscis immaculatis, posticis flauescensibus, nigro punctatis. *Murcia*.

Habitat in Africa aequinoctiali. *Mus. Dom. Banks*.

Medius. Corpus nigrum abdominis lateribus albo maculatis. Alae anticae fuscae omnino immaculatae.

*Fabr. Sp. Inf. T. II.*

C

Posti-



Posticae flavescentes punctis margineque nigris, subtus praeterea puncta aliquot alba ad basin et striga punctorum nigrorum in margine nigro.

**Caecilia.** 142. P. H. alis oblongis medio albis nigro punctatis, posticis margine nigro subtus albo punctato.

Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*

Statura et summa affinitas praecedentis. Alae omnes supra basi apiceque nigrae, in medio vero albicantes nigro punctatae. Subtus omnes albiae nigro punctatae, anticae apice nigrae, posticae margine postico nigro albo punctato.

**Bonasia.** 143. P. H. alis oblongis integerrimis fuscis fascia communi fulva, posticis basi nigro punctatis. *Syst. Ent.* 464. 96. \*

Habitat in Sierra Leon Africae. *Mus. Dom. Banks.*

**Euterpe.** 144. P. H. alis oblongis integerrimis nigris albo punctatis, fascia fulva. *Syst. Ent.* 465. 97. *Linn. Syst. Nat.* 2. 756. 61. *Mus. Lud. Vlr.* 226.

*Seb. Mus.* 4. tab. 7. fig. 17. 18.

Habitat in America.

Alae omnes pone fasciam immaculatae.

**Erythrea.** 145. P. H. alis oblongis atris, anticis rubro fasciatis, posticis striatis.

*Papilio Erythrea. Cram. Inf.* 16. tab. 189. fig. A.

Habitat Surinami.

Alae anticae basi fasciaque media rubris, posticae striis sex rubris.

**Erato.** 146. P. H. alis oblongis integerrimis atris, anticis flavo maculatis basi rubris, posticis rubro striatis. *Syst. Ent.* 465. 98. *Linn. Syst. Nat.* 2. 757. 70. *Mus. Lud. Vlr.* 231.

*Clerk. Icon.* tab. 40. fig. 1.

*Papilio Vesta. Cram. Inf.* 10. tab. 119. fig. A.

Habitat in Indiis.

Alae anticae maculis decem flavis, posticae striis quinque rubris basi coeuntibus.

Conf. *Papilio Aegeria. Cram. Inf.* 3. tab. 34. fig. B.

G. Vix distinctus videtur.

\*\*\* Par.

\*\*\*\* Parnassii.

147. P. P. alis rotundatis integerrimis albis nigro maculatis, posticis supra ocellis quatuor, subtus sex. *Syst. Ent.* 465. 99. *Linn. Syst. Nat.* 2. 754. 50. *Fn. Sv.* 1032. *Scop. carn.* 447. *Apollo.*

*Papilio alpina maior*, alis albicantibus, exterioribus maculis nigris notatis, interioribus maculis ophthalmoidibus, iride rubra. *Raj. Inf.* 139. 2.

*Papilio Rhea. Pod. Inf.* 66. 11.

*Mouff. Inf.* 94. fig. 2. 3.

*Petiv. Mus.* 49. n. 502.

— — *Gazoph.* tab. 23. fig. 8.

*Degeer Inf.* 1. tab. 18. fig. 12. 13.

*Roes. Inf.* 4. 29. tab. 4. fig. 1. 2. Larva.

— — 3. tab. 45. fig. 1. 2. Imago.

*Sut. Inf.* tab. 13. fig. 41.

*Schaeff. monogr.* 1754. tab. 2. fig. 2. 3.

— — *Elem.* tab. 9. fig. 6.

— — *Icon.* tab. 36. fig. 4. 5.

*Aubent. miscell.* tab. 68. fig. 1. 2.

*Esp. Papilion.* 1. tab. 2. fig. 1.

Habitat in Sedo Telephio, Saxifraga cotyledone.

*Larva* solitaria, tentaculata, holosericea, atra, segmentis vtrinque punctis rubris duobus.

*Puppa* coerulescens antice vtrinque rubro punctata.

148. P. P. alis rotundatis integerrimis albis nigro nervosis, anticis maculis duabus nigris marginalibus. *Syst. Ent.* 466. 100. *Linn. Syst. Nat.* 2. 754. 51. *Fn. Sv.* 1033. *Mnemosyne.*

*Esp. Papil.* 1. tab. 2. fig. 2.

*Vdm. Diff.* 55.

*Scop. carn.* 449.

*Pod. Inf.* 63. 3. 8.

*Schaeff. Icon.* tab. 34. fig. 6. 7.

Habitat in Europa boreali.

Præcedente minor et absque ocellis.

149. P. P. alis rotundatis integerrimis albis, venis nigris. *Crataegi. Syst. Ent.* 466. 101. *Linn. Syst. Nat.* 2. 758. 72. *Fn. Sv.* 1034.

*Papilio* alis rotundatis albis, venis nigris. *Geoff. Inf.*

2. 71. 43.

*Papilio* alba nervis alarum nigris. *Raj. Inf.* 115. 5.

*Papilio* alba venis nigris. *Petiv. Mus.* 33. 301.

*Eruca* hyemalis luteo alboque striata. *Frisch. Inf.* 5.

16. tab. 5.

*Scop. carn.* 448.

*Huffn. Inf.* tab. 10. fig. 14.

*Merian. Europ.* 3. 38. tab. 35.

*Albin. Inf.* tab. 2. fig. 2.

*Reaum. Inf.* 2. tab. 2. fig. 9. 10.

*Degeer Inf.* 1. tab. 14. fig. 19. 20.

*Roes. Inf.* 1. Pap. 2. tab. 3.

*Wilk. papil.* 49. tab. 2. a. 1.

*Esp. papil.* 1. tab. 2. fig. 3.

*Scop. Ann. Hist. nat.* 5. p. 112. n. 114.

*Schaeff. Icon.* tab. 140. fig. 2. 3.

*Seb. Mus.* 4. tab. 3. fig. 1 -- 4.

*Harris Aurel.* tab. 9. fig. k. i.

Habitat in Pomonae arboribus, hortorum pestis.

*Larva* gregaria pilosa flaua, capite lincisque tribus nigris.

*Puppa* virefcens maculis punctisque nigris.

*Andromacha.* 150. P. P. alis rotundatis integerrimis concoloribus nigro punctatis, anticis denudatis, posticis flavescentibus.

*Syst. Ent.* 466. 102. \*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

*Horta.* 151. P. P. alis rotundatis integerrimis rubris, anticis apice hyalinis, posticis subtus albidis nigro punctatis. *Syst.*

*Ent.* 459. 69. *Linn. Syst. Nat.* 2. 755. 54. *Mus. Lud.*

*Flr.* 234.

*Papilio Horta. Sulz. Hist. Inf.* tab. 15. fig. 1.

*Seb. Mus.* 4. tab. 39. fig. 4 -- 9.

Habitat in Africa.

*Quirina.* 152. P. P. alis rotundatis integerrimis hyalinis, posticis disco fuluo nigro punctato.

Habitat in India, Madras. *Mus. Dom. Banks.*

Statura praecedentis. Alae omnes albo hyalinae nigro venosae, at posticae disco late fuluo punctis numerosis atris.

153. P. P.

CLASS. VI. GLOSSATA. Papilio. 37

153. P. P. alis rotundatis integerrimis hyalino diaphanis, posticis ocellis binis marginalibus. *Syst. Ent.* 467. 103. *Linn. Syst. Nat.* 2. 754. 52. *Mus. Lud. Vlr.* 220. *Sulz. Hist. Inf.* tab. 3. fig. 5. *Clerk. Icon.* tab. 36. fig. 4. *Gron. Zooph.* 734. 735. *Merian. Sur.* tab. 16. fig. 1. *Kleeman. Inf.* 1. tab. 6. fig. 1. *Seb. Mus.* 4. tab. 8. fig. 6. 7. Habitat in Indiis. *Pieris.*
154. P. P. alis rotundatis nigris albo subradiatis puncto centrali albo, posticis subtus luteis basi cruentatis. *Syst. Ent.* 467. 104. *Linn. Syst. Nat.* 2. 755. 53. *Papilio Porfenna. Cram. Inf.* 4. tab. 43. fig. D. E. Habitat in India. *Pasithoe.*  
Alae posticae subtus sulphureae marginem versus nigris et basi sanguineae.  
Variat alis posticis basi superiori etiam sanguineae.
155. P. P. alis rotundatis integerrimis hyalino virescentibus, posticis apice fuscis, ocellis duobus. *Syst. Ent.* 467. 105. *Linn. Syst. Nat.* 2. 758. 74. *Mus. Lud. Vlr.* 236. *Anacardii.*  
*Clerk. Icon.* tab. 28. fig. 3.  
*Merian. Sur.* tab. 16.  
*Seb. Mus.* 4. tab. 35. fig. 17. 18.  
Habitat in Anacardio Americas.  
Forte P. Pierae mera varietas.  
*Larva* alba pilosa pilis longis deflexis.  
*Puppa* brunnea.
156. P. P. alis dentatis, anticis hyalinis maculis duabus nigris, posticis nigris macula alba. *Syst. Ent.* 448. 24.\* *Cressida.*  
Habitat in noua Hollandia. *Mus. Dom. Banks.*
157. P. P. alis rotundatis integerrimis fusco hyalinis, posticis punctis quinque albis. *Syst. Ent.* 467. 106.\* *Rhea.*  
Habitat in Indiis. *Mus. Hagense.*
158. P. P. alis rotundatis integerrimis hyalinis albis, posticis oculo vtrinque unico. *Syst. Ent.* 467. 107.\* *Andromeda.*  
Habitat in India. *Mus. Dom. Gronovii.*

*diaphanus*. 159. P. P. alis rotundatis integerrimis albo hyalinis, marginibus fuscis. *Syst. Ent.* 468. 108. \*

*Papilio hyalinus*. *Drury Inf.* 2. tab. 7. fig. 3.

*Papilio Astrea*. *Cram. Inf.* 2. tab. 22. fig. D.

Habitat in Iamaica.

*Idea*. 160. P. P. alis rotundatis integerrimis denudato albis, venis maculisque nigris. *Syst. Ent.* 468. 109.

*Papilio Idea* alis rotundatis integerrimis albis, venis maculisque nigris, anticis nigro margine albo punctato. *Linn. Syst. Nat.* 2. 758. 73. *Amoen. acad.* 6. 405. 63.

*Papilio Idea*. *Cram. Inf.* 17. tab. 193. fig. A. B.

*Papilio Lynceus*. *Drury Inf.* 2. tab. 7. fig. 1.

*Clerk. Icon.* tab. 38. fig. 1.

*Aubert. miscell.* tab. 20. fig. 1.

Habitat in Indiis.

Præcedentibus maior.

\*\*\*\*\* Danai candidi.

*Brassicae*. 161. P. D. C. alis integerrimis rotundatis albis, anticis maculis duabus apicibusque nigris maior. *Syst. Ent.* 468. 110. *Linn. Syst. Nat.* 2. 759. 75. *Fn. Sv.* 1035.

*Papilio alba* maior, brassicaria vulgarissima. *Raj. Inf.* 113. 1.

*Papilio albus* vulgaris maior. *Petiv. Gazoph.* 62. fig. 3.

*Papilio alba* maior apicibus nigris. *Pettiv. Mus.* 85. 825.

*Geoff. Inf.* 2. 68. 40.

*Roeß. Inf.* 1. pap. 2. tab. 4.

*Goed. Inf.* 1. 59. tab. 11.

*Mouff. Inf.* 189. fig. 1.

*Esp. Papil.* 1. tab. 3. fig. 1.

*List. Goed.* p. 16. fig. 7.

*Merian. Europ.* 1. p. 16. tab. 45.

*Albin. Inf.* 1. tab. 1.

*Swammerd. Bibl.* tab. 37. fig. 6.

*Reaum. Inf.* 1. tab. 29. fig. 1.

*Blank. Inf.* tab. 4. fig. A. D.

*Vallijn. Nat.* 1. tab. 1.

*Wilk. pap.* 49. tab. 2. a. 2.

*Schaeff. Icon.* tab. 40. fig. 3. 4.

*Schaeff.*

*Schaeff. Icon. tab. 140. fig. 4. 5.*

*Sepp. Inf. 2. p. 7. tab. 2.*

Habitat in Europae, Chinae Brassica.

*Larva* solitaria cinerea, nigro punctata, lineisque tribus sulphureis, cauda nigra.

*Puppa* pallide virescens, lincis tribus flauis segmentisque tribus gibbosis.

Variat sexu absque maculis atris alarum anticarum.

162. P. D. C. alis integerrimis, anticis maculis duabus apicibusque nigris minor. *Syst. Ent.* 469. 111. *Linn. Syst. Nat.* 2. 759. 76. *Fn. Sv.* 1036. *Geoff. Inf.* 2. 69. 41.

*Papilio alba media*, alis exterioribus albis, duabus maculis nigris notatis, inferioribus subtus flauicantibus. *Raj. Inf.* 114. 2.

*Papilio* corpore et antennis liuescentibus, capite alisque pallidis. *Merret. pin.* 198. 5.

*Papilio vulgaris albus minor.* *Petiv. Mus.* 15. 826.

*Papilio alba vulgaris striata maculis duplicibus.* *Petiv. Gazoph.* 10. tab. 64. fig. 4.

*Mouff. Inf.* 971. fig. 9. 10.

*Robert. Icon.* tab. 6.

*Albin. Inf.* tab. 51. fig. C. D. E. F.

*Reaum. Inf.* 1. tab. 29. fig. 7. 8.

*Blank. Inf.* tab. 5. fig. A — D.

*Goed. Inf.* 1. tab. 27.

*Lift. Goed.* 22. fig. 8.

*Merian. Europ.* 2. tab. 39.

*Ammir.* tab. 16.

*Sepp. Inf.* 2. p. 1. tab. 1.

*Esp. Pap.* 1. tab. 3. fig. 2.

*Holl. Inf.* tab. 2. fig. 10.

Habitat in Europae brassicis.

*Larva* viridis, linea dorsali aenea lateribusque aeneo punctatis.

*Puppa* gibba, virescens, lincis tribus sulphureis.

163. P. D. C. alis integerrimis albis, subtus venis dilatis virescentibus. *Syst. Ent.* 469. 112. *Linn. Syst. Nat.* 2. 760. 77. *Fn. Sv.* 1037. *Napi.*

*Papilio* alis rotundatis albis, inferioribus subtus fasciis virescentibus. *Geoff. Inf.* 2. 70. 42.

*Papilio* brassicaria alis albis secundum nervos lineis e viridi nigricantibus subtus striatis. *Raj. Inf.* 114. 4.

*Wilk. pap.* 50. tab. 2. a. 4.

*Merian. Europ.* 2. tab. 39.

*Alb. Inf.* tab. 52. fig. F. G.

*Petiv. Gazoph.* tab. 62. fig. 4.

— — *Mus.* 33. 302.

*Esp. papil.* 1. tab. 3. fig. 3.

— — — tab. 24. fig. 2.

Habitat in Europae, Asiae oleraceis.

*Sinapis.* 164. P. D. C. alis integerrimis rotundatis albis, apicibus fusciscentibus. *Syst. Ent.* 470. 114. *Linn. Syst. Nat.* 2. 760. 79. *Fn. Sv.* 1038.

*Papilio* alba minor. *Raj. Inf.* 116. 8.

*Schaeff. Icon.* tab. 97. fig. 8-11.

*Degeer Inf.* 2. 183. 4. tab. 1. fig. 1.

*Esp pap.* 1. tab. 3. fig. 4.

*Bergstr. Nomencl.* 2. tab. 19. fig. 5. 6. 7.

*Harr. Inf.* tab. 19. fig. 1.

*Scop. carn.* 452.

Habitat in Brassica, Sinapi, Rapa.

Color apicis alae anticae variat sexu.

*Poppea.* 165. P. D. C. alis rotundatis integerrimis concoloribus albis, margine nigro maculato, anticis macula bascos ferruginea.

*Papilio Poppea.* *Cram. Inf.* 10. tab. 110. fig. D.

Habitat in Sierra Leon Africae.

Nimis affinis P. Sylulae, a quo tamen differt magnitudine maculis marginalibus nigris et macula ferruginea bascos in vtraque pagina. Forte mera varietas.

*Sylula.* 166. P. D. C. alis rotundatis integerrimis albis, anticis apice fuscis, subtus macula bascos fulva, *Syst. Ent.* 470. 115.\*

Habitat in Sierra Leon Africae. *Mus. Dom. Banks.*

*Argante.* 167. P. D. C. alis integerrimis rotundatis fulvis, subtus ferrugineo irroratis. *Syst. Ent.* 470. 116.\*

Habitat in Brasilia. *Mus. Dom. Banks.*

CLASS. VI. GLOSSATA. Papilio

45

168. P. D. C. alis rotundatis integerrimis albis, margine fuscis. *Syst. Ent.* 470. 117. *Linn. Syst. Nat.* 2. 760. 80. *Mus. Lud. Vlr.* 237.  
*Papilio Monusta. Cram. Inf.* 12. tab. 141. fig. F.  
*Papilio Alcyone. Cram. Inf.* 5. tab. 58. fig. A.  
*Klemm. Inf.* tab. 3. fig. 3.  
*Seligm. Av.* 6. tab. 79.  
 Habitat in America.  
 Alae posticae subtus flavescentes.
169. P. D. C. alis integerrimis rotundatis albis, anticis apice nigris. *Syst. Ent.* 470. 118.\*  
*Papilio Cassiopia. Cram. Inf.* 17. tab. 201. fig. A.  
 Habitat in Sierra Leon Africae. *Mus. Dom. Banks.*
170. P. D. C. alis rotundatis albis apice nigris, posticis subtus margine brunneo. *Amathonte.*  
*Papilio Amathonte. Cram. Inf.* 10. tab. 116. fig. A. B.  
 Habitat in America meridionali.  
 Statura P. Monuste. Abdomen niveum maculis dorsalibus atris.
171. P. D. C. alis rotundatis integerrimis griseis, anticis supra albis apice fuscis. *Syst. Ent.* 471. 119.\*  
 Habitat in America.
172. P. D. C. alis integerrimis rotundatis albis, anticis costa bascos apiceque fuscis. *Syst. Ent.* 471. 120.\*  
 Habitat in India orientali. *Mus. Tottianum.*
173. P. D. C. alis rotundatis integerrimis albis, anticis puncta apiceque nigris, posticis subtus flavescentibus, margine nigro punctato. *Syst. Ent.* 471. 121.\*  
*Papilio Calypso. Cram. Inf.* 13. tab. 154. fig. C. D. ♂  
 — — — — — 13. tab. 154. fig. E. F. ♀.  
*Papilio Calypso. Drury Inf.* 2. tab. 17. fig. 3. 4.  
*Seb. Mus.* 4. tab. 26. Fig. 3. 4. ♂  
 — — — — — tab. 6. fig. 9. 10. ♀  
 Habitat in Sierra Leon Africae.
174. P. D. C. alis integris rotundatis albis, margine fuscis, subtus luteo griseis albo maculatis. *Syst. Ent.* 471. 122. *Linn. Syst. Nat.* 2. 760. 81. *Daplidicta.*



42 CLASS. VI. GLOSSATA. Papilio.

*Papilio Daphidice.* Pall. *Spicil. fasc.* 9. tab. 2.

*Papilio Edusa.* Gen. Inf. Mant. 255.\*

*Cram. Inf.* 15. tab. 171. fig. C. D.

*Seb. Mus.* 4. tab. 23. fig. 11. 12. 15. 16.

*Petiv. Gazoph.* tab. 1. fig. 7.

*Schaeff. Icon.* tab. 79. fig. 2. 3.

Habitat in Africa, Germania, Kiloniae.

*Creona.* 175. P. D. C. alis integris rotundatis albis, margine albo, striga punctorum.

*Papilio Creona.* *Cram. Inf.* 8. tab. 95. fig. C. D. E. F.

Habitat in India orientali.

Alae omnes albae margine exteriori nigro, in quo striga punctorum, feminae albidorum, maris fuluorum.

Punctum atrum in medio alae anticae. Posticae subtus subuirescentes.

*Nerissa.* 176. P. D. C. alis integerrimis rotundatis albis, margine nigro, subtus nigro venosis. *Syst. Ent.* 471. 123.\*

*Papilio Amasine.* *Cram. Inf.* tab. 44. fig. A.

Habitat in China.

*Demophi.* 177. P. D. C. alis rotundatis albis, fasciis duabus margineque fuscis, posticis subtus subincarnatis. *Syst. Ent.*

472. 124. *Linn. Syst. Nat.* 2. 761. 81. *Amoen. acad.* 6. 406. 66.

*Clerk. Icon.* tab. 28. fig. 4.

Habitat in India.

*Hecabe.* 178. P. D. C. alis integerrimis rotundatis flauis extimo nigris, subtus fusco punctatis. *Syst. Ent.* 472. 124.

*Linn. Syst. Nat.* 2. 763. 96. *Mus. Lud. Vlr.* 249.

*Papilio Hecabs.* *Cram. Inf.* 11. tab. 124. fig. B. C. et maior.

*Papilio Nise.* *Cram. Inf.* 2. tab. 20. fig. K. L.?

*Papilio Agare.* *Cram. Inf.* 2. tab. 20. fig. H. I.?

*Sulz. Hist. Inf.* tab. 15. fig. 7.

*Gronov. Zooph.* 757.

*Edw. Av.* tab. 253. fig. 2.

*Petiv. Gazoph.* tab. 28. fig. 9.

Habitat in Aeschynomene pumila Indiae orientalis. *D. Kosnig.*

179. P. D. C. alis rotundatis integerrimis albis, posticis *Cardami-*  
subtus viridi marmoratis. *Syst. Ent.* 472. 126. *Linn. nes.*  
*Syst. Nat.* 2. 761. 85. *Fn. Sv.* 1039. *Geoff. Inf.* 2.  
71. 44.

*Papilio* minor alis exterioribus albis macula insigni  
crocea splendentibus, interioribus superne albis,  
subtus viridi colore variegatis. *Raj. Inf.* 115.

*Edw. Av.* tab. 125.

*Roeß. Inf. pap.* 2. tab. 8.

*Robert. Icon.* tab. 21.

*Schaeff. Icon.* tab. 91. fig. 1 - 3.

— — *Elem.* tab. 94. fig. 8.

*Wilk. pap.* 2. p. 50. tab. 2. a. 5.

*Seligm. Av.* 5. tab. 20.

*Esp. papil.* 1. tab. 4. fig. 1.

— — — tab. 27. fig. 2.

*Harris* tab. 32. fig. g. h.

*Hufn. Icon.* tab. 9. fig. 1.

Habitat in Cardamine, Brassica, Thlaspi.

Mas macula fulva alae anticae.

*Larva* solitaria, supra virescens, infra albicans.

*Puppa* viridis, linea vtrunque laterali alba, thorace  
conico adscendente.

180. P. D. C. alis rotundatis integerrimis albis, anticis *Xiphia.*  
macula apiceque nigris, posticis subtus viridi ir-  
roratis.

Habitat in India orientali. *Mus. Dom. Banks.*

Corpus paruum. Alae omnes albae, nitidae, anticis  
supra macula apiceque nigris, posticis immaculatis.  
Subtus anticae macula versus apicem nigra lituraque  
baeos et apicis viridi. Posticae lineolis transversis  
numerosis irroratae.

181. P. D. C. alis rotundatis integerrimis albis, anticis ap- *Eucharis.*  
ce fulvis, margine nigro, posticis immaculatis. *Syst.*  
*Ent.* 472. 127.\*

Habitat in India orientali. *Mus. Tottianum.*

182. P. D. C. alis rotundatis integerrimis albis, posticis *Nysa.*  
subtus fuscis, puncto albo lunulisque sex flanis.  
*Syst. Ent.* 473. 128.\*

Habitat in nova Hollandia, *Mus. Dom. Banks.*

183. P.

44 CLASS. VI. GLOSSATA. Papilio.

- Chloris*. 183. P. D. C. alis rotundatis integerrimis albis apice nigris, posticis subtus fulvis, margine late nigro.  
*Syst. Ent.* 473. 129.\*  
*Papilio Thermopyle. Cram. Inf.* 18. tab. 207. fig. F. G.  
 Habitat in Sierra Leon Africae. *Mus. Dom. Banks*.
- Rhodope*. 184. P. D. C. alis rotundatis integerrimis subconcoloribus externis nigris, anticis flavis, posticis albis. *Syst. Ent.* 473. 130.\*  
 Habitat in Sierra Leon Africae. *Mus. Dom. Banks*.
- Elathea*. 185. P. D. C. alis rotundatis integerrimis extimo nigris, anticis flavis, posticis albis, subtus cinereis.  
*Papilio Elathea. Cram. Inf.* 9. tab. 99. fig. C. D.  
*Roef. Inf.* 4. tab. 3. fig. 4.  
 Habitat in America.  
 Affinis videtur praecedenti, at margo niger haud flavo striatus et subtus omnes cinereae.
- Alcmeone*. 186. P. D. C. alis rotundatis concoloribus, basi flavis, apice albis.  
*Papilio Alcmeone. Cram. Inf.* 12. tab. 141. fig. E.  
 Habitat in Malabar.ia.  
 Alae omnes basi flavae, apice albae. Margo apicis alae anticae ater.
- Phryne*. 187. P. D. C. alis integerrimis albis, margine nigro, posticis subtus flavis. *Syst. Ent.* 473. 131.\*  
*Papilio Licinia. Cram. Inf.* 13. tab. 153. fig. E. F. vix differre videtur.  
 Variat alis subtus haud nigro venosis. Habitat in America. *Dom. Drury*.
- Pyranthe*. 188. P. D. C. alis rotundatis albis puncto apiceque nigris, subtus cinereo vndatis puncto fulvo. *Syst. Ent.* 473. 132. *Linn. Syst. Nat.* 2. 763. 98. *Mus. Lud. Vlr.* 245.  
 Habitat in Indiae orientalis Cassis. *D. Koenig*.
- Leucippe*. 189. P. D. C. alis integerrimis fulvis, posticis supra flavis.  
*Gen. Inf. Mant.* 256.\*  
*Papilio Leucippe. Cram. Inf.* 3. tab. 36. fig. A. B. C.  
 Habitat in India orientali, Amboina.

CLASS. VI. GLOSSATA. Papilio: 45

190. P. D. C. alis integris albis, anticis apice, posticis margine punctisque sex nigris. *Syst. Ent.* 473. 133.\* *Callirhoe.*  
*Aubert. miscell.* tab. 91. fig. 1. 2.  
 Habitat in Asia.
191. P. D. C. alis integerrimis rotundatis albis, anticis apice nigris, posticis subtus cinereis fusco vndatis. *Glaucippe.*  
*Syst. Ent.* 474. 134. *Linn. Syst. Nat.* 2. 762. 89.  
*Mus. Lud. Vlr.* 240.  
*Papilio Glaucippe. Cram. Inf.* 14. tab. 164. fig. A. B. C.  
*Seligm. Av.* 5. tab. 23.  
*Clerk. Icon.* tab. 35. fig. 1.  
*Drury Inf.* 1. tab. 10. fig. 3. 4.  
*Edw. Av.* 128. tab. 128.  
*Seb. Mus.* 4. tab. 45. fig. 5. 6. 13. 14.  
 Habitat in Asia.
192. P. D. C. alis integerrimis rotundatis flavescentibus, anticis apice nigris, posticis subtus albis. *Eupippe.*  
*Syst. Ent.* 474. 135. *Linn. Syst. Nat.* 2. 762. 87. *Mus. Lud. Vlr.* 239.  
*Papilio Eupippe. Cram. Inf.* tab. 91. fig. D. E. F. G. ?  
*Aubert. miscell.* tab. 92. fig. 1. 2.  
*Seb. Mus.* 4. tab. 26. fig. 1. 2.  
*Gron. Zooph.* 758.  
*Petiv. Gazoph.* tab. 9. fig. 10.  
 Habitat in Asia.
193. P. D. C. alis integerrimis nigro venosis, supra albis, subtus virefcentibus. *Coronnis.*  
*Gen. Inf. Mant.* 256.\*  
*Papilio Coronnis. Cram. Inf.* 4. tab. 44. fig. B. C.  
 Habitat in China.
194. P. D. C. alis integerrimis albis nigro venosis, posticis subtus flavis, margine rubro maculato. *Hyparete.*  
*Syst. Ent.* 474. 36.\*  
*Papilio Hyparete alis integerrimis rotundatis albis, posticis subtus venis nigris maculisque sanguineis.*  
*Linn. Syst. Nat.* 2. 763. 92. *Mus. Lud. Vlr.* 247.  
*Papilio Antonoe. Cram. Inf.* 16. tab. 187. fig. C. D.  
*Papilio Eucharis. Cram. Inf.* 17. tab. 201. fig. B. C.  
 — — — tab. 202. fig. C.  
*Salz.*

46 CLASS. VI. GLOSSATA. Papilio.

*Sulz. Hist. Inf.* tab. 15. fig. 8.

*Schroet. Abhandl.* 1. tab. 1. fig. 4.

*Papilio Eucharis. Drury Inf.* 2. tab. 10. fig. 5. 6.

*Clerk. Icon.* tab. 38. fig. 2. 3.

*Aub. Miscell.* tab. 93. fig. 3. 4.

Habitat in Indiae orientalis Lorantho.

Conf. *Papilio Iffe. Cram. Inf.* 5. tab. 55. fig. E. F.

*Iphigenia.* 195. P. D. C. alis integerrimis albis apice atris, posticis subtus flavescentibus, fascia media atra, striga abbreviata fulva. *Gen. Inf. Mant.* 256.\*

*Papilio Iphigenia. Naturf.* 9. tab. 2. fig. C.

Habitat Surinami. *D. Schulz.*

*Teutonia.* 196. P. D. C. alis integerrimis rotundatis albis, posticis subtus nigro venosis flavo maculatis. *Syst. Ent.* 474. 137.\*

*Papilio Teutonia. Sulz. Hist. Inf.* tab. 15. fig. 9.?

Habitat in nova Hollandia. *Muf. Dom. Banks.*

*Myfis.* 197. P. D. C. alis integerrimis rotundatis albis, posticis subtus margine atro, fascia sanguinea. *Syst. Ent.* 475. 138.\*

Habitat in nova Hollandia. *Muf. Dom. Banks.*

*Nigrina.* 198. P. D. C. alis integerrimis rotundatis albis, apice nigris, subtus nigris, posticis striga flexuosa sanguinea. *Syst. Ent.* 475. 139.\*

Habitat in nova Hollandia. *Muf. Dom. Banks.*

*Saba.* 199. P. D. C. alis rotundatis integerrimis nigris, fascia communi alba.

Habitat in Africa aequinoctiali. *Muf. Dom. Banks.*

Medius. Alae omnes supra nigrae, fascia lata alba in medio. Puncta aliquot alba versus apicem alae anticae. Subtus concolores litura tantum flava ad basin et alae posticae magis albae.

*Pyrrha.* 200. P. D. C. alis integerrimis albis, posticis subtus atris, striga flava rubraque fasciaeque alba.

*Papilio Pyrrha. Cram. Inf.* 6. tab. 63. fig. A. B.

Habitat Surinami.

Alae

Alae supra albae apice nigrae. Alae posticae subtus atrae striga bascos flava, quae fere cum fascia postica alba coit. Striga maior abbreviata rubra et in medio fascia lata postice dentata alba.

201. P. D. C. alis integerrimis albis, margine nigris, sub- *Coronea*  
tus atris basi fuluo radiatis.

*Papilio Coronea. Cram. Inf. 6. tab. 68. fig. B. C.*

Habitat Surinami.

Alae omnes subtus atrae basi fuluo radiatae et versus marginem striga punctorum fuluorum.

202. P. D. C. alis rotundatis integerrimis albidis, apice ni- *Melania*  
gris, posticis subtus obscure glaucis. *Syst. Ent.*  
475. 140.\*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

203. P. D. C. alis integerrimis rotundatis flauis, anticis *Scylla*  
supra albis, limbo nigro, subtus omnibus nebulosis.

*Syst. Ent. 475. 142. Linn. Syst. Nat. 2. 763. 95.*

*Mus. Lud. Vir. 242. Amoen. acad. 6. 404. 57.*

*Papilio Scylla. Cram. Inf. 1. tab. 12. fig. C. D.*

*Sulz. Hist. Inf. tab. 15. fig. 6.*

*Seb. Mus. 4. tab. 28. fig. 19. 20.*

Habitat in India.

204. P. D. C. alis rotundatis fuluis, limbo nigro, subtus *Amata*  
virescentibus. *Syst. Ent. 476. 143.\**

Habitat in India orientali. *Mus. Tottianum.*

205. P. D. C. alis rotundatis albis, anticis apice croceis *Danae*  
margine fasciaque nigris, subtus striga monilliformi  
abbreviata. *Syst. Ent. 476. 144.\**

Habitat in India orientali. *Mus. Tottianum.*

Striga in pagina inferiori alae anticae e punctis quinque subocellaribus fuscis pupilla albicante, alae posticae e punctis quinque subocellaribus ferrugineis.

206. P. D. C. alis rotundatis flauis apice (medio fuluo) *Sesia*  
nigris, posticis subtus maculis septem ocellaribus al-  
bis. *Gen. Inf. Mant. 257.\**

*Papilio Pyrene. Cram. Inf. 11. tab. 125. fig. A. B.*

*Drury Inf. 1. tab. 5. fig. 2.*

Habitat in Americae meridionalis insulis. *D. Spengler.*  
207. P,

*Aenippe.* 207. P. D. C. alis rotundatis flavis, omnibus sub punctis ocellaribus, anticis sex, posticis septem.

*Papilio Aenippe.* *Cram. Inf.* 9. tab. 105. fig. C. D.

— — — — — 14. tab. 157. fig. C. D.

Habitat in China.

Affinis videtur praecedenti. Alae anticae flavae, apice nigrae macula magna fulva. Subtus flavae puncto medio atro sexque ocellaribus albis, inde nigro cinctis. Posticae flavae margine nigro maculato, subtus flavae puncto medio atro septemque ocellaribus albis, iride atra cinctis.

*Rhexia.* 208. P. D. C. alis rotundatis integerrimis albis apice (medio fulvo) nigris, singulis subtus puncto medio atro. *Syst. Ent.* 476. 745. \*

Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*

*Trite.* 209. P. D. C. alis integerrimis flavis, anticis supra margine nigris, singulis subtus striga ferruginea. *Syst. Ent.* 476. 146. *Linn. Syst. Nat.* 2. 763. 97. *Mus. Lnd. Vir.* 248.

*Papilio Trite.* *Cram. Inf.* 12. tab. 141. fig. C. D.

*Seb. Mus.* 4. tab. 23. fig. 17. 18.

Habitat in Africa, Europa australi.

*Palatino.* 210. P. D. C. alis integerrimis flavis, apice nigris margineque fulvis, posticis subtus puncto argenteo. *Syst. Ent.* 476. 147. *Linn. Syst. Nat.* 2. 764. 99. *Fn. Sv.* 1041.

*Papilio hexapus* alis rotundatis albis oculo parvo fusco oblongo apicibus fuscis. *Vdm. Diff.* 56.

*Bergstr. Nomencl.* 2. tab. 16. fig. 5. 6.

*Esp. pap.* 1. tab. 4. fig. 2.

*Schaeff. Icon.* tab. 149. fig. 4. 5.

Habitat in Europae Pteride aquilina.

Antennae rubrae.

*Hyale.* 211. P. D. C. alis rotundatis flavis, posticis macula fulva, subtus puncto selqui altero argenteo. *Syst. Ent.* 477. 148. *Linn. Syst. Nat.* 2. 764. 100. *Fn. Sv.* 1040.

*Papilio* alis luteis limbo nigro, primariis macula nigra, secundariis fulva. *Geoff. Inf.* 2. 75. 48.

*Papilio*

**CLASS. VI. GLOSSATA. Papilio.** 49

*Papilio* mediae magnitudinis, alis exterioribus summa parte flavis, ad margines nigricantibus. *Raj. Inf.* 112. 6.

*Papilio* croceus apicibus nigricantibus. *Petiv. Gazoph.* tab. 14. fig. 11.

*Schaeff. Icon.* tab. 149. fig. 1. 2. 3.

*Cram. Inf.* 2. tab. 14. fig. F. G.

*Esp. papil.* 1. tab. 4. fig. 3.

*Bergstr. Nomencl.* 2. tab. 16. fig. 3. 4.

*Seb. Mus.* 4. tab. 61. fig. 1-5.

*Roef. Inf.* 3. tab. 46. fig. 5. 6.

Habitat in Europa, Africa, America.

112. P. D. C. alis integerrimis fulvis, anticis puncto fusco, *Lesbia*.  
omnibus subtus pallidis, puncto niueo. *Syst. Ent.* 477. 149.\*

Habitat in Patagoniae litoribus. *Mus. Dom. Banks.*

113. P. D. C. alis integerrimis rotundatis flavis, singulis  
subtus puncto gemino ferrugineo. *Syst. Ent.* 477.  
150. *Linn. Syst. Nat.* 2. 764. 103.

*Sloan. Iam.* 2. tab. 236. fig. 11. 12.

*Merian. Sur.* 58. tab. 58.

*Raj. Inf.* 112. 5.

*Seb. Mus.* 4. tab. 10. fig. 13. 14. 17. 18.

— — — tab. 23. fig. 13. 14.

— — — tab. 26. fig. 5. 6.

— — — tab. 38. fig. 15. 16.

Habitat in Americae Cassia.

Variat alis supra flauissimis apicibus nigris et immaculatis.

Conf. *Papilio* *Cipris*. *Cram. Inf.* 2. tab. 99. fig. E. F.

114. P. D. C. alis integris rotundatis flavis, singulis sub-*Marcellina*.  
tus puncto gemino argenteo. *na.*

*Papilio* *Marcellina*. *Cram. Inf.* 14. tab. 163. fig. A. B. C.

Habitat Surinami.

Nimis affinis *P. Sennae*, vix distinctus. Alae omnes supra flauissimae margine nigro punctato, subtus punctis duobus approximatis argenteis.

*Fabr. Spec. Inf. T. II.* D 215. P.



*Eubule.* 215. P. D. C. alis integerrimis rotundatis flavis, margine punctis nigris, posticis subtus puncto gemino ferrugineo argenteo. *Syst. Ent.* 477. 151. *Linn. Syst. Nat.* 2. 764. 102.

*Papilio Eubule.* *Cram. Inf.* 10. tab. 120. fig. F. G.

*Papilio Arioge.* *Cram. Inf.* 8. tab. 94. fig. A. B.?

*Seb. Mus.* 4. tab. 38. fig. 17. 18. 19. 20.

— — — tab. 23. fig. 19. 20.

*Edw. Av.* tab. 304.

Habitat in Carolina.

Variat sexu. Alter alis anticis ocello medio ferrugineo et margine fusco maculato. Alter alis supra immaculatis.

Conf. *Papilio Herfilia.* *Cram. Inf.* 15. tab. 173. fig. C. D.

*Proterpia.* 216. P. D. C. alis integerrimis angulatis fulvis, anticis margine exteriori nigro. *Syst. Ent.* 478. 152.\*

Habitat in Jamaica. *Mus. D. Hunter.*

*Gnoma.* 217. P. D. C. alis integris rotundatis albis, anticis puncto atro, subtus omnibus puncto argenteo. *Syst. Ent.* 828.\*

Habitat in India. *Mus. D. Hunter.*

*Drya.* 218. P. D. C. alis integerrimis rotundatis flavis, subtus anticis puncto ferrugineo, posticis argenteo. *Syst. Ent.* 478. 153.\*

*Papilio Statira.* *Cram. Inf.* 10. tab. 120. fig. C. D.

Habitat in America.

*Ecclipsis.* 219. P. D. C. alis integerrimis angulatis flavis, anticis punctis duobus maculaque nigris, posticis ocellis coeruleo. *Syst. Ent.* 478. 154. *Linn. Syst. Nat.* 2. 765. 107. *Almoen. acad.* 6. 406. 67.

*Pitt. Gacopa.* 6. tab. 10. fig. 6.

Habitat in America.

*Rhamni.* 220. P. D. C. alis integerrimis angulatis flavis, singulis puncto ferrugineo. *Syst. Ent.* 478. 155. *Linn. Syst. Nat.* 2. 755. 106. *Fa. Sv.* 1042. *Groff. Inf.* 2. 74. 47

*Papilio praecox* sulphurea siue flavo viridis, singulis alis macula ferruginea notatis. *Raj. Inf.* 112. 4.

*Robert*

CLASS. VI. GLOSSATA. Papilio.

51

*Robert. Icon.* tab. 13.

*Roef. Inf.* 3. tab. 46. fig. 1. 2. 3.

— — 4. tab. 26. fig. 1 - 5.

*Alb. Inf.* tab. 2. fig. 3.

*Sulz. Inf.* tab. 13. fig. 84.

*Wilk. pap.* 48. tab. 1. a. 2.

*Schaeff. Elem.* tab. 94. fig. 7.

— — *Icon.* tab. 35. fig. 1 - 3.

*Degeer Inf.* 1. tab. 15. fig. 1 - 10.

*Esp. pap.* 1. tab. 4. fig. 4.

*Bergstr. Inf.* 2. tab. 15. fig. 5 - 8.

Habitat in Europae, Africae Rhamno.

Mas sulphureus, femina alba.

*Larva* laevis viridis, linea dorsali obscuriore.

*Puppa* gibbosa, antice acuminata.

221. P. D. C. alis integerrimis angulatis flavis, anticis ma- *Philea.*  
cula, posticis limbo luteis. *Syst. Ent.* 478. 156.

*Linn. Syst. Nat.* 2. 764. 104. *Amoen. acad.* 6.

404. 59.

*Papilio Philea. Cram. Inf.* 15. tab. 173. fig. E. F.

*Roef. Inf.* 4. tab. 3. fig. 5.

Habitat in Indiis.

Alae omnes subtus puncto gemino albo annulo purpureo cincto.

222. P. D. C. alis integris angulatis flavis, anticis supra *Maerula.*  
macula atra, reliquis ferruginea. *Syst. Ent.* 479.

157\*.

*Papilio Ecclipsis. Cram. Inf.* 11. tab. 129. fig. A. B.

Habitat in America. *Mus. D. Hunter.*

223. P. D. C. alis subangulatis albis, anticis puncto ni- *Pomona.*  
gro, subtus flavis, posticis punctis duobus argenteis.

*Syst. Ent.* 479. 158.\*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

224. P. D. C. alis subangulatis albis, anticis puncto fusco, *Florilla.*  
posticis subtus punctis tribus argenteis. *Syst. Ent.*

479. 159.\*

Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*

- Cleopatra*. 225. P. D. C. alis integerrimis angulatis flauis, anticis supra disco fuluo, reliquis puncto ferrugineo. *Syst. Ent.* 479. 160. *Linn. Syst. Nat.* 2. 765. 105.  
*Papilio Cleopatra*. *Cram. Inf.* 11. tab. 131. fig. E.  
*Esp. pap.* 1. tab. 48. suppl. 24. fig. 1.  
*Seb. Mus.* 4. tab. 26. fig. 7. 8.  
 Habitat in Africa, in Lusitania. *D. Gray.*

## † Danai festiui.

- Cyparissias*. 226. P. D. F. alis integerrimis nigris, fasciis duabus albis anticarum obliquis, posticarum punctatis. *Gen. Inf. Mant.* 257.\*  
*Papilio Daedalus*. *Cram. Inf.* 1. tab. 1. fig. A. B.  
 Habitat in America meridionali.

- Midas*. 227. P. D. F. alis integerrimis nigris albo punctatis, anticis supra coerulescentibus, posticis supra punctorum alborum striga. *Syst. Ent.* 479. 161. *Linn. Syst. Nat.* 2. 765. 108. *Mus. Lud. Vlr.* 251.  
*Sulz. Hist. Inf.* tab. 16. fig. 4. a. b.  
*Alt. Holm.* 1748. tab. 6. fig. 1. 2.  
*Ehret. pict.* tab. 3. et 11.  
*Kleman. Inf.* 1. tab. 9.  
*Seb. Mus.* 4. tab. 19. fig. 13. 14.  
 Habitat in Indiae orientalis Paretta. *D. Koenig.*  
 Conf. *Papilio Mulciber*. *Cram. Inf.* 11. tab. 127. fig. 2.  
 C. D. alis posticis supra immaculatis.

- Niarius*. 228. P. D. F. alis integerrimis nigris, anticis fascia alba, posticis macula communi baseos alba. *Syst. Ent.* 480. 162. *Linn. Syst. Nat.* 2. 766. 109. *Mus. Lud. Vlr.* 253.  
*Cram. Inf.* 1. tab. 2. fig. F. G.  
*Clerk. Icon.* tab. 32. fig. 2.  
 Habitat in Africa aequinoctiali.

- Liberius*. 229. P. D. F. alis integerrimis coeruleis, subtus anticis viridibus, posticis fuluis.  
*Papilio Liberia*. *Cram. Inf.* 18. tab. 210. fig. G. H.  
 Habitat in Amboina.  
 Margo omnis vtrunque tenuissime ater.

30. P. D. F. alis integerrimis atris, anticis maculis tribus, posticis disco albo. *Syst. Ent.* 480. 163.\* *Zoilus.*  
Habitat in noua Hollandia. *Mus. Dom. Banks.*

31. P. D. F. alis integerrimis atris coeruleo maculatis, posticis subtus punctis sanguineis. *Arethusa.*  
*Papilio Arethusa. Cram. Inf.* 7. tab. 17. fig. E. F.  
Habitat in Surinami.

Corpus atrum thorace punctis coeruleis. Alae omnes supra atrae maculis plurimis coeruleis. Subtus anticae immaculatae, posticae punctis bascos apicisque coccineis.

*Conf. Papilio Laodamia. Cram. Inf.* 11. tab. 100. fig. A. Differt tantum fascia alae anticae alba.

32. P. D. F. alis integris, fascia cyanea, subtus anticis sanguineis, posticis albis annulis punctisque atris. *Clymenus.*  
*Papilio Clymena. Cram. Inf.* 2. tab. 24. fig. E. F.  
Habitat in Brasilia. *Mus. Dom. Banks.*

Ad Pap. Nymphal. phaler. refert *Cramer*, at alae integrae, supra atrae fascia viridi, subtus anticae rufae apice nigrae lineolis duabus albis, posticae albae striis duabus marginis tenuioris, annulis duobus contiguïs disci strigisque duabus marginis atris. Puncta duo atra in singulo annulo.

33. P. D. F. alis integerrimis rotundatis nigris, supra fascia communi fulua, *Syst. Ent.* 480. 164.\* *Linn. Antiochus.*  
*Mant.* I. 537.  
*Papilio Eupalemon. Cram. Inf.* 12. tab. 143. fig. B. C.  
*Aubent. miscell.* tab. 68. fig. 3. 4.  
Habitat in China.

34. P. D. F. alis integerrimis atris, anticis fascia alba, posticis rubra marginali. *Syst. Ent.* 48. 165.\* *Syphax.*  
Habitat in Indiis. *Mus. Lugdunense.*

35. P. D. F. alis integerrimis fuscis, anticis fasciis duabus, posticis vnica albis maculisque rubris. *Eualthe.*  
*Papilio Eualthe* alis integerrimis fuscis, anticis fasciis duabus flauis, posticis fascia flaua rubraque, *Syst. Ent.* 480. 166.\*

54 CLASS. VI. GLOSSATA. Papilio.

*Papilio Dardanus*. Cram. Inf. 2. tab. 17. fig. E. F.

Habitat in Indiis.

Posticae subtus rufae fascia media maculari alba.

*Pelaeus*. 236. P. D. F. alis integerrimis concoloribus, anticis virescentibus, fascia fulva, posticis nigris immaculatis.

*Papilio Pelaeus*. Cram. Inf. 17. 202. fig. D.

Habitat Surinami.

Antennae nigrae, vinctatae.

*Phaëton*. 237. P. D. F. alis integerrimis nigris albo rufoque maculatis. *Syst. Ent.* 481. 167.

*Papilio Phaëton*. Drury Inf. 1. tab. 21. fig. 3. 4.

Cram. Inf. 17. tab. 193. fig. C. D.

Habitat in America.

*Galanthis*. 238. P. D. F. alis integerrimis supra atris sanguineo fasciatis, anticis punctis duobus apicis albis. *Gen. Inf.*

*Mant.* 257.\*

*Papilio Galanthis*. Cram. Inf. 3. tab. 25. fig. D. E.

Habitat Surinami.

*Caeneus*. 239. P. D. F. alis integerrimis subfuscis, posticis subtus orbe macularum novem rubro. *Syst. Ent.* 481. 168.

*Linn. Syst. Nat.* 2. 766. 111. *Mus. Lud. Vlr.* 271.

Habitat in Indiis.

Alae supra fuscae flavo maculatae basi pallidae.

*Tytius*. 240. P. D. F. alis integerrimis fuscis albo maculatis anticis punctis quatuor ocellaribus atris.

*Papilio Tytia*. Cram. Inf. 11. tab. 121. fig. C. D.

Habitat Surinami.

Alae omnes supra fuscae albo maculatae, ad basin anticae maculae quatuor approximatae atrae, annulo albo cinctae, posticarum puncto obscuriori. Subtus pallidiores albo maculatae punctis plurimis atris sparsis subocellaribus.

*Enpalus*. 241. P. D. F. alis integerrimis fuscis, posticis fascia coerulea, subtus viridibus, puncto atro.

*Papilio Harpalyce*. Cram. Inf. 13. tab. 145. fig. D. E.

Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*

Medius

**Medius.** Alae omnes supra fuscae auro parum nitentes. Anticae apice terminantur puncto albo, posticae fascia cyanea, quae ad angulum ani latior. Alae omnes subtus virides puncto baseos atro. In quibusdam striga punctorum albidorum. Antennae nigrae apice ferrugineae. Corpus nigrum.

242. P. D. F. alis integerrimis fuscis concoloribus, anticis maculis tribus flavis, posticis dimidiato albis. *Pinthaeus*.  
*Syst. Ent.* 481. 169. *Linn. Syst. Nat.* 2. 766. 114.  
*Mus. Lud. Vlr.* 253.

Habitat in Indiis.

243. P. D. F. alis integerrimis fulvis, venis nigris dilatis, margine nigro, punctis albis. *Plexippus*.  
*Syst. Ent.* 481. 170. *Linn. Syst. Nat.* 2. 767. 117. *Mus. Lud. Vlr.* 262.  
*Papilio Iamaicensis* maior, alis amplissimis, media parte fulvis cum nervis nigris maculis et punctis albis crebris sparsis. *Raj. Inf.* 138. 3.

*Papilio carolinarius* rufescens limbis nigris, albis guttulis adspersis. *Petiv. Mus.* 527.

*Papilio Plexippus.* *Cram. Inf.* 18. tab. 206. fig. E. F.

*Papilio Eriippus.* *Cram. Inf.* 1. tab. 3. fig. A. B.

*Papilio Genutia.* *Cram. Inf.* 18. tab. 206. fig. C. D.

*Sloan. Iamaic.* 2. 214. tab. 239. fig. 5; 6.

*Catesb. Carol.* 2. tab. 88.

*Seb. Mus.* 4. tab. 6. fig. 15. 16.

— — — tab. 41. fig. 9. 10. 13. 14.

Habitat in America boreali.

*Larva* albo nigroque annulata, tentaculis duobus colli caudaeque.

*Puppa* incrassata viridis striga punctisque aureis.

Conf. *Papilio Berenice.* *Cram. Inf.* 18. tab. 205. fig. E. F.

244. P. D. F. alis subintegriss fulvis, margine nigro albo punctato, posticis arcu nigro. *Missippus*.  
*Syst. Ent.* 481. 171.

*Papilio Missippus* alis subintegerrimis fulvis margine nigro alboque lineatis, posticis macula nigra. *Linn.*

*Syst. Nat.* 2. 767. 118. *Mus. Lud. Vlr.* 263.

*Papilio Archippus.* *Cram. Inf.* 2. tab. 16. fig. A. B.

*Seb. Mus.* 4. tab. 6. fig. 17. 18.

Habitat in America.

Simillimus P. *Plexippo*, at minor.

Conf. *Papilio Melanippus*. *Cram. Inf.* 11. tab. 127. fig. A. B.

*Chrysippus*. 245. P. D. F. alis integerrimis fulvis, margine nigro punctis albis, posticis disco punctis nigris. *Syst. Ent.* 481. 172. *Linn. Syst. Nat.* 2. 767. 119. *Mus. Lud. Vlr.* 263.

*Papilio lamaicensis* minor fulvescens, imis alis limbo nigro guttulis albis adperso cinctis. *Raj. Inf.* 139. 1.

*Papilio Aegyptius*. *Schreb. Inf.* 9. fig. 11. 12.

*Papilio Chrysippus*. *Cram. Inf.* 10. tab. 118. fig. B. C.

*Papilio Diocippus*. *Cram. Inf.* 3. tab. 28. fig. B. C.

*Seligm. Av.* 6. tab. 84.

*Edw. Av.* 189. tab. 189.

*Sulz. Hist. Inf.* tab. 16. fig. 3.

*Kleman. Inf.* 1. tab. 1. fig. 1.

*Seb. Mus.* 4. tab. 6. fig. 9 — 12.

— — — tab. 13. fig. 13. 14.

Habitat in Asclepiade fruticosa Indiae orientalis.

Femina alis posticis magis maculatis.

Conf. *Papilio Erisimus*. *Cram. Inf.* 15. tab. 175. fig. G.

H. vii et *Papilio Iparia*. *Cram. Inf.* 18. tab. 214. fig. A. B.

*Alcippus*. 246. P. D. F. alis integerrimis fulvis, margine nigro punctis albis, posticis disco albo, punctis nigris.

*Papilio Alcippus*. *Cram. Inf.* 11. tab. 127. fig. E. F.

Habitat in America.

Vix a praecedente satis distinctus.

*Gilippus*. 247. P. D. F. alis integerrimis concoloribus fulvis albo maculatis, margine atro, punctis albis. *Gen. Inf. Mant.* 258. \*

*Papilio Gilippus*. *Cram. Inf.* 3. tab. 26. fig. C. D.

Habitat in America meridionali.

*Hegesippus*. 248. P. D. F. alis integerrimis concoloribus nigris albo maculatis, anticis basi fulva posticis albo lineatis.

*Papilio Hegesippus*. *Cram. Inf.* 15. tab. 180. fig. A.

Habitat in Sumatra.

Statura praecedentium. Alae anticae atrae, basi maculis tribus oblongis fulvis lineatae, apice albo maculatae. Posticae fuscae, basi albo lineatae, apice strigis duabus punctorum alborum.

**CLASS. VI. GLOSSATA. Papilio. 57.**

1. P. D. F. alis integerrimis fuscis, anticis albo maculatis, posticis subtus arcu nigro. *Syst. Ent. 482. 173.\** *Darius.*  
*Habitat in Brasilia. Mus. Dom. Banks.*
2. P. D. F. alis integerrimis fuscis albo maculatis, subtus brunneis. *Syst. Ent. 482. 174.\** *Daedalus.*  
*Habitat in Guinea. Mus. Dom. Banks.*
3. P. D. F. alis integerrimis fuscis, anticis ocello, posticis supra fascia rubra marginali punctisque nigris ocellaribus. *Syst. Ent. 482. 175.\** *Nicaeus.*  
*Papilio Salome. Cram. Inf. 1. tab. 12. fig. G. H.*  
*Habitat in India.*  
*Variat sexu absque ocello alae anticae.*  
*Conf. Papilio Lamia. Cram. Inf. 13. tab. 150. fig. C.*
4. P. D. F. alis integerrimis atris, anticis maculis duabus, posticis unica fulvis. *Numilius.*  
*Papilio Numilia. Cram. Inf. 7. tab. 81. fig. E. F. at alae in nostro integerrimae.*  
*Habitat in Brasilia. Mus. Dom. Banks.*  
*Medius. Corpus fuscum. Alae anticae supra atrae maculis duabus dilute fulvis, anteriore maiore, subtus fulvescentes fascia apicis fusca. Posticae supra atrae macula transversa baseos fulvae lituraeque marginis postici coerulescente. Subtus griseo fuscac.*
53. P. D. F. alis integerrimis atris sanguineo fasciatis, posticis subtus lineis annularibus flavis punctisque coeruleis. *Codomanus.*  
*Habitat in Brasilia. Mus. Dom. Banks.*  
*Minor praecedente. Alae anticae supra atrae basi fasciaeque, quae margines haud attingit, sanguineis. Punctum fulvum transversum versus apicem et margo apicis albo punctatus. Subtus fere concolores fascia tantum flava et striga coerulea apicis. Posticae supra atrae vitra abbreviata fulvae, subtus atrae lineis annularibus flavis punctisque coerulescentibus. Pectus albo punctatum.*
54. P. D. F. alis integerrimis fuscis coeruleo micantibus, posticis subtus flavis, lineis annularibus nigris punctisque tribus coeruleis. *Hyaspes.*



58 CLASS. VI. GLOSSATA. Papilio.

Habitat in Brasilia. *Mus. Dom. Banks.*

Præcedente minor. Alae anticae concolores fascia sanguinea lituraque apicis flava. Posticae supra immaculatae, subtus flavae lineis annularibus atris punctisque tribus subocellaribus in medio strigaque postica cyaneis.

*Gripus.* 255. P. D. F. alis integris subfuscis, subtus striga punctorum alborum. *Syst. Ent. App.* 829. \*

*Papilio Eumea.* *Cram. Inf.* 16. tab. 183. fig. C. D.

*Drury Inf.* 1. tab. 2. fig. 3.

Habitat in India. *Mus. D. Hunter.*

*Cramer* alas pingit dentatas. Forte potius *Papilio Nymphalis*.

*Sophorae.* 256. P. D. F. alis integerrimis fuscis, fascia ferruginea, posticis subtus brunneis ocellis tribus. *Syst. Ent.* 483. 176. *Linn. Syst. Nat.* 2. 767. 121. *Mus. Lud. Vlr.* 266.

*Clerk. Icon.* tab. 35. fig. 3.

*Merian. Sur.* tab. 35.

*Roef. Inf.* 4. tab. 4. fig. 1. 2.

*Seb. Mus.* 4. tab. 11. fig. 3—6. 7—9.

— — — tab. 15. fig. 10. 11.

Habitat in Sophora Americæ.

*Larva* gregaria, nuda, lineata.

*Xanthus.* 257. P. D. P. alis integerrimis fuscis, fascia ferruginea, posticis subtus griseo marmoratis ocellis duobus. *Syst. Ent.* 483. 177. *Linn. Syst. Nat.* 2. 767. 122. *Mus. Lud. Vlr.* 267.

*Papilio Xanthus.* *Cram. Inf.* 16. tab. 183. fig. A. B.

*Clerk. Icon.* tab. 34. fig. 1. 2.

*Seb. Mus.* 4. tab. 11. fig. 1. 2.

— — — tab. 15. fig. 5. 6.

— — — tab. 20. fig. 15. 16.

Habitat in Asia.

Ocellus alter lunula alba notatus.

*Cassiae.* 258. P. D. F. alis subintegris fuscis, fascia ferruginea apice bifida, posticis subtus cinereis ocellis duobus. *Syst. Ent.* 483. 178.

*Papilio*

*Papilio Cassiae* alis integerrimis fuscis, primoribus posticarumque margine fascia ferruginea, omnibus subtus ocellatis. *Linn. Syst. Nat.* 2. 767. 120. *Mus. Lud.* Vlr. 265.

*Papilio Cassiae.* *Cram. Inf.* 9. tab. 105. fig. A. B. at alae magis dentatae.

*Papilio Cassiae.* *Cram. Inf.* 9. tab. 106. fig. A.

*Clerk. Icon.* tab. 34. fig. 3. 4.

*Merian. Sur.* tab. 32.

Habitat in Americis Cassia.

*Larva* viridis rubro striata, capite bispinoso, cauda bifida.

*Puppa* rufescens maculis argenteis.

259. P. D. F. alis integerrimis supra atris, fascia marginali flava, subtus posticis maculis sex ocellaribus. *Berecynthus.*

*Papilio Berecynthia.* *Cram. Inf.* 16. tab. 184. fig. B. C.

*Seb. Mus.* 4. tab. 13. fig. 2. 3.

Habitat Surinami.

Magnus. Alae omnes supra atrae fascia marginali fulva, anticarum angulata. Puncta tria alba ad apicem alae anticae. Subtus variegatae anticis macula ocellari unica, posticis sex.

260. P. D. F. alis integerrimis fuscis, striga marginali fulva, posticis subtus subundatis ocellis septem. *Oethon.*

Habitat Surinami. *Mus. Dom. Teats.*

Statura omnino P. Cassiae. Alae anticae fuscae striga apicis fulva punctisque aliquot albis, subtus oculo paruo atro. Posticae supra fuscae striga marginali fulva, subtus strigis albis fuscisque obsoletis undatae ocellis septem.

261. P. D. F. alis integerrimis supra fuscis, anticis fascia, posticis disco omnibusque subtus albis. *Pirithous.* *Syst. Ent.* 483. 179. \*

Habitat in America boreali. *Mus. Dom. Banks.*

262. P. D. F. alis integris fuscis, posticis subtus griseo fuscoque nebulosis. *Polixes.* *Syst. Ent.* 484. 180. \*

Habitat in America boreali. *Mus. Dom. Banks.*

*Cupanius*. 263. P. D. F. alis integris atris albo maculatis, posticis subtus griseis, fascia vtrinque alba.

*Papilio Cupania*. *Cram. Inf.* 17. tab. 193. fig. E. F.

Habitat in Guinea, Sierra Leon.

Alae posticae supra atrae, subtus griseae, vtrinque fascia albida, abbreviata.

*Marthesius*. 264. P. D. F. alis integris atris, anticis disco fuluo, posticis striga punctorum alborum.

*Papilio Marthesia*. *Cram. Inf.* 16. tab. 191. fig. A. B.

Habitat Surinami.

Affinis videtur P. Orion. Alae anticae supra fulvae, apice atrae punctis aliquot albis, subtus pallidiores. Posticae atrae striga e punctis septem albis, subtus coerulescentes fasciis rufis atrisque.

*Polycar-mes*. 265. P. D. F. alis integerrimis supra nigris, subtus brunneis immaculatis. *Syst. Ent.* 484. 181. \*

Habitat in Indiis. *Mus. Lugdunense*.

*Chorinaeus*. 266. P. D. F. alis integerrimis fuscis, anticis falcatis fascia fulva, posticis caudatis. *Syst. Ent.* 484. 182. \*

*Papilio Arcefilaus*. *Sulz. Inf.* tab. 14. fig. 4.

Habitat Surinami. *Dom. Drury*.

*Eribotes*. 267. P. D. F. alis integerrimis subcaudatis fulvis, basi coerulescentibus, subtus griseis, *Syst. Ent.* 484. 183. \*

Habitat in India. *Mus. D. Hunter*,

*Itys*. 268. P. D. F. alis subcaudatis fulvis, apice flavo nigris, subtus variegatis, posticis basi coeruleis.

*Papilio Itys*. *Cram. Inf.* 10. tab. 119. fig. F. G.

Habitat Surinami.

Margo exterior alae anticae subtus coerulco punctatus.

*Laertes*. 269. P. D. F. alis caudatis fuscis basi coeruleis, posticis supra punctis quinque albis, subtus quatuor ocellaribus,

*Papilio Laertes*. *Cram. Inf.* 7. tab. 73. fig. C. D.

Habitat Surinami.

Affinis sequenti.

270. P. D. F. alis integris caudatis, supra fuscis, basi coeruleis, subtus griseo virescentibus. *Morus.*

*Papilio Morus* alis integris subcaudatis, supra fuscis coeruleo nitentibus, subtus griseis. *Syst. Ent.* 484. 184.

*Papilio Arachne. Cram. Inf.* 4. tab. 48. fig. A. B.

Habitat in India. *Mus. D. Hunter.*

Medius. Alae anticae fuscae, basi coerulecentes margineque interiori valde emarginato. Posticae caudatae fuscae, basi coerulecentes. Subtus omnes griseo virescentes.

271. P. D. F. alis subcaudatis fuscis, anticis fascia lutescente exteriore maculaque coerulea postica subtusque ocello. *Syst. Ent.* 533. 386. *Linn. Syst. Nat.* 2. 794. 262. *Amoen. acad.* 6. 409. 78.

*Clerk. Icon.* tab. 36. fig. 2.

*Cram. Inf.* 5. tab. 49. fig. C. D.

*Seb. Mus.* 4. tab. 41. fig. 17. 18.

— — — tab. 4. fig. 3. 4.

Habitat Surinami.

Macula coerulea nitida ad marginem interiorem alae anticae.

272. P. D. F. alis subcaudatis fuscis, basi coeruleo nitidis, posticis ocello vtrinque unico. *Cymodoc.*

*Papilio Cymodocé. Cram. Inf.* 9. tab. 99. fig. G. H.

Habitat Surinami.

Alae anticae fuscae, basi coeruleo nitidae, versus apicem litura albida strigaeque atra. Subtus subconcolores. Posticae subcaudatae fuscae, basi cyaneae ocello ferrugineo pupilla alba. Subtus dilutiores striga ferruginea coeruleo adnata. Versus apicem ocellus punctaque aliquot atra.

273. P. D. F. alis subcaudatis fulvis, apice nigris, subtus punctis ocellaribus, anticis duobus, posticis tribus. *Bifaltide.*

*Papilio Bifaltide. Cram. Inf.* 9. tab. 102. fig. C. D.

Habitat Surinami.

Ante apicem maculae duae lineolaeque flavae, et in apice puncta duo alba. Subtus omnes basi flavescentes striga atra.

- Orion.** 274. P. D. F. alis integerrimis, anticis falcatis rufis, apice nigris, macula marginali alba. *Syst. Ent.* 485. 185. \*  
Habitat Surinami. *Mus. D. Hunter.*
- Obrinus.** 275. P. D. F. alis integerrimis atris, fascia anticarum coerulea, posticarum ferruginea, subtus viridibus. *Syst. Ent.* 485. 186.  
*Papilio Obrinus. Naturf.* 6. 128. tab. 6. fig. 1. 2.  
*Papilio Ancaea. Linn. Syst. Nat.* 2. 781. 184.  
*Papilio Obrinus. Linn. Mus. Lud. Vlr.* 255. β.  
*Sulz. Hist. Inf.* tab. 16. fig. 1. 2.  
Habitat in India.
- Ancaeus.** 276. P. D. F. alis integerrimis, supra fuscis fascia coerulea maculaque ferruginea, subtus viridibus fascia albicante. *Syst. Ent.* 485. 187. \*  
*Papilio Obrinus. Linn. Syst. Nat.* 2. 766. 113.  
*Clerk. Icon.* tab. 31. fig. 2. 3.  
*Cram. Inf.* 5. tab. 49. fig. E. F.  
Habitat in India.
- Lucindus.** 277. P. D. F. alis integerrimis nigro fasciatis, supra coeruleis, subtus fulvis. *Gen. Inf. Mant.* 258. \*  
*Papilio Lucinda. Cram. Inf.* 1. tab. 1. fig. E. F.  
*Seb. Mus.* 4. tab. 13. fig. 25. 26.  
Habitat Surinami.
- Morphus.** 278. P. D. F. alis integerrimis fulvis nigro maculatis, posticis vtrunque striga punctorum nigrorum. *Syst. Ent.* 530. 370. \*  
*Papilio Cocyta. Cram. Inf.* 9. tab. 101. fig. A. B. C.  
Habitat in America boreali.  
*Conf. Papilio Tharos. Cram. Inf.* 15. tab. 169. fig. E. F.
- Lyfidice.** 279. P. D. F. alis integerrimis coeruleis, supra limbo, subtus fasciis atris.  
*Papilio Lyfidice. Cram. Inf.* 15. tab. 169. fig. C. D.  
Habitat Surinami.  
Alae anticae coeruleae nervis marginibusque atris. Posticae coeruleae margine atro. Subtus omnes coeruleae

ruleae fasciis atris. In fascia tertia anticarum punctum argentum et in posticarum tria.

280. P. D. F. alis integerrimis fuscis, fascia alba, posticarum vtrinque terminata macula fulva. *Eumens.*

Habitat in Caiennia. *Mus. Dom. Teats.*

Corpus fuscum. Alae omnes concolores fuscae fascia alba, quae in alis posticis ad marginem tenuiorem et crassiorem macula fulva terminatur. Apex alae albicat.

281. P. D. F. alis integerrimis fuscis, posticis disco albo, supra ocello, subtus duabus. *Gen. Inf. Mant.* 258. \* *Iairus.*

*Papilio Iairus. Cram. Inf.* 1. tab. 6. fig. A. B.

— — — 16. tab. 185. fig. A. B. C.

*Clerk. Icon.* tab. 29. fig. 3.

Habitat in India Orientali.

282. P. D. F. alis integerrimis coeruleis, anticis albo maculatis, posticis macula atra albaque marginali. *Arcas.*

*Papilio Arcas. Cram. Inf.* 15. tab. 179. fig. E. F.

Habitat Surinami.

Macula magna atra, albo terminata, ad marginem exteriorrem alae posticae.

283. P. D. F. alis integerrimis fuscis, anticis fascia flava ocelloque subtripupillato, posticis ocellis quinque. *Clytus.*

*Syst. Ent.* 485. 188. *Linn. Syst. Nat.* 2. 768. 124. *Mus.*

*Lud. Vlr.* 268.

*Papilio Clytus. Cram. Inf.* 8. tab. 86. fig. C. D.

Habitat in Africa.

284. P. D. F. alis integerrimis fuscis area rufa, anticis ocello bipupillato, posticis subtus coecis. *Hyperbius.*

*Syst. Ent.* 485. 189. *Linn. Syst. Nat.* 2. 769. 130. *Mus. Lud.*

*Vlr.* 257.

*Papilio Hyperbius. Cram. Inf.* 14. tab. 168. fig. C. D. E. F.

Habitat in Africa.

Ocellus ad apicem alae anticae niger punctis duobus albis.

*Philocles*. 285. P. D. F. alis integerrimis fuscis, anticis ocello tripupillato, posticis lineis transversis vndatis. *Gen. Inf. Mant.* 259. \* *Linn. Syst. Nat.* 2. 791. 240. *Mus. Lud. Vlr.* 321.

*Papilio Philocles*. *Cram. Inf.* 16. tab. 124. fig. D. E. F.

*Papilio Ofinia*. *Cram. Inf.* 10. tab. 115. fig. F.

*Papilio Thymetus*. *Cram. Inf.* 16. tab. 124. fig. G.

*Clerk. Icon.* tab. 45. fig. 3.

Habitat in Indiis.

Ocellus alae anticae vtrinque tripupillatus.

Variat alis supra atris fascia coerulescente maculaque magna alba anguli ani et alis fuscis lineis transversis vndatis fasciaque alba.

*Croesus*. 286. P. D. F. alis integerrimis cyaneis strigis atris numerosis, anticis supra macula ocellari tripupillata. *Gen. Inf. Mant.* 259. \*

Habitat Surinami. *D. Schulz.*

*Aeropus*. 287. P. D. F. alis integerrimis fuscis fascia lutea, antica subtus basi ocello unico. *Syst. Ent.* 486. 190. *Linn. Syst. Nat.* 2. 769. 122. *Mus. Lud. Vlr.* 256.

*Papilio Aeropus*. *Cram. Inf.* 10. tab. 111. fig. F. G.

*Clerk. Icon.* tab. 39. fig. 1. 2.

*Sed. Mus.* 4. tab. 9. fig. 3. 4.

Habitat in Indiis.

*Canthus*. 288. P. D. F. alis integra, supra fuscis immaculatis, posticis subtus ocellis sex. *Syst. Ent.* 486. 191.

*Papilio Canthus* alis integerrimis fuscis, subtus primis ocellis quatuor, posticis senis. *Linn. Syst. Nat.* 2. 788. 120.

*Papilio Eurydice*. *Amoen. acad.* 6. 456. 65.

*Papilio Argantha*. *Cram. Inf.* 17. tab. 204. fig. C. D.

Habitat in America boreali.

*Hyperan*. 289. P. D. F. alis integerrimis fuscis, subtus anticis ocellis tribus, posticis duobus tribusque. *Syst. Ent.* 486. 192. *Linn. Syst. Nat.* 2. 752. 127. *Fa. Sv.* 204.

*Cress. In.* 2. 47. 14.

*Papilio Fugurea*. *Sey. carn.* 424.

*Papilio* media tota pulla prona alarum parte ocellis aliquot e puncto albo, duplici circulo nigro et for-  
dide luteo cincto compositis. *Raf. Inf.* 129. 7.

*Schaeff. Icon.* tab. 127. fig. 1. 2.

*Degeer Inf.* 2. tab. 2. fig. 9. 10.

*Sepp. Inf.* 1. tab. 4.

*Naturf.* 8. tab. 3. fig. D.

*Esp. pap.* 1. tab. 5. fig. 1.

*Bergstraeff. Inf.* 2. tab. 16. fig. 7. 8.

— — — tab. 17. fig. 5. 6. 7. 8.

*Harris Inf.* tab. 35. fig. h. k.

Habitat in Europae Poa annua ad radicem.

Variat alis supra ocellatis et immaculatis.

*Larua* solitaria, pilosa, nigra.

190. P. D. F. alis integerrimis fuscis, omnibus subtus *Zangis*.  
ocello vnico, anticarum bipupillato. *Syst. Ent.* 486.

193. \*

Habitat in Carolina. *Mus. D. Hunter.*

191. P. D. F. alis integerrimis fuscis, anticis ocellis vtrin. *Eurytus*.  
que duobus, posticis supra vnico subtus quatuor,  
omnibus pupilla gemina. *Syst. Ent.* 487. 194. \*

*Papilio Cymela. Cram. Inf.* 11. tab. 132. fig. C. D.

Habitat in Iamaica. *Mus. D. Hunter.*

192. P. D. F. alis integerrimis supra fuscis immaculatis, *Hermes*.  
posticis subtus ocellis sex, tribus coecis. *Syst. Ent.*  
487. 195. \*

*Papilio Antonoe. Cram. Inf.* 5. tab. 60. fig. E. F. at  
ocelli tantum quinque postico bipupillata.

Habitat in Brasilia. *Mus. Dom. Banks.*

193. P. D. F. alis integerrimis disco flauo, posticis ocello *Irius*.  
supra vnico subtus duobus. *Syst. Ent.* 487. 196. \*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

Variat alis totis flauescens marginibus solo fusco.

194. P. D. F. alis integerrimis fuscis, anticis supra ocello *Minus*.  
vnico, posticis subtus septem fasciaque alba. *Syst.*  
*Ent.* 488. 197. \*

*Fabr. Sp. Inf. T. II.*

8

*Papilio*



66 CLASS. VI. GLOSSATA. Papilio.

*Papilio Mineus* alis integerrimis subfuscis, subtus posticis ocellis septem, anticis duobus supra vnico. *Linn. Syst. Nat.* 2. 768. 126.

*Papilio Drusia*. *Cram. Inf.* 7. tab. 84. fig. C. D. at alae dentatae.

Habitat in China.

*Medus.* 295. P. D. F. alis integerrimis fuscis, subtus striga niuea ocellisque anticarum duobus, posticarum tribus. *Syst. Ent.* 488. 198. \*

Habitat ad Cap. bon. sp.

*Perseus.* 296. P. D. F. alis integerrimis fuscis, subtus punctis ocellaribus, anticis tribus posticis septem. *Syst. Ent.* 488. 199. \*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

*Terminus.* 297. P. D. F. alis integris fuscis area rufa, supra anticis ocello vnico, posticis quatuor. *Syst. Ent.* 488. 200. \*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

*Sirius.* 298. P. D. F. alis integerrimis obscure rufis, supra anticis ocellis duobus, posticis quatuor. *Syst. Ent.* 488. 201. \*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

*Daus.* 299. P. D. F. alis integerrimis fuluis, anticis ocellis duobus, posticis sex coecis, subtus pupillatis. *Gen. Inf. Mant.* 259. \*

Habitat Hamburgi. *D. Schulz. Kiloniae Dom. de Scheffelt.*

Affinis P. Sabaco.

*Pamphilus.* 300. P. D. F. alis integerrimis flauis, subtus anticis ocello vnico, posticis cinereis, fascia ocellisque quatuor oblitteratis. *Syst. Ent.* 529. 368. *Linn. Syst. Nat.* 2. 791. 239. *Fn. Sv.* 1044.

*Papilio Menaleas.* *Scop. carn.* 458.

*Papilio* alis rotundatis fuluis, oris fuscis, primariis subtus ocello vnico, secundariis subtus albo cinereoque variegatis. *Geoff. Inf.* 2. 53. 21.

*Papilio* paruus e fuluo et fusco bicolor fuluo mediam alam fusco oras extremas occupante cum ocello ad extre-

**CLASS. VI. GLOSSATA. Papilio. 67**

**extremum angulum alarum exteriorum. Raj. Inf.**  
125. 19.

*Merian. Europ. tab. 134.*

**Habitat in Europae pratis syluaticis in Cynosuro cristato.**

**Larua** virescens linea dorsali albida caudaque bidentata. *Naturf. 6. 10.*

301. P. D. F. alis integerrimis ferrugineis, subtus anticis *Arcan-*  
ocello vnico, posticis quinque, primo fascia remoto. *us.*  
*Syst. Ent. 530. 369. Linn. Syst. Nat. 2. 791. 242. Fn.*  
*Sv. 1045.*

**Papilio** alis rotundatis, primoribus fuluis oris fuscis,  
subtus ocello vnico, posticis supra fuscis, infra cin-  
reis, fascia alba ocellisque quinque. *Geoff. Inf. 2.*  
53. 22.

**Papilio** *Amyntas. Scop. carn. 457.*

*Schaeff. Elem. tab. 94. fig. 3.*

**Habitat in Europae pratis syluaticis.**

302. P. D. F. alis integerrimis fuscis, posticis supra ocel- *Sabaeus.*  
lis quatuor coecis, subtus sex pupillatis. *Syst. Ent.*  
530. 371. \*

**Habitat in Lipsiae nemoribus.**

303. P. D. F. alis integerrimis fuluis, subtus anticis ocel- *Hero.*  
lo, posticis senis. *Syst. Ent. 530. 372. Linn. Syst.*  
*Nat. 2. 793. 253. Fn. Sv. 1047.*

**Habitat in Europae pratis syluaticis.**

304. P. D. F. alis integerrimis fuscis, anticis vtrinque *Arctous.*  
ocello bipupillato, posticis supra subcoecis. *Syst. Ent.*  
489. 202. \*

**Habitat in noua Hollandia. Mus. Dom. Banks.**

305. P. D. F. alis integerrimis concoloribus albis fusco *Hypheus.*  
fasciatis, anticis ocello atro, posticis coecis.

**Papilio** *Hyphea. Cram. Inf. 8. tab. 92. fig. 6.*

**Habitat in India occidentali.**

**Alae omnes concolores albae fasciis quatuor fuscis vlti-  
ma terminali. Ocellus anticarum in fascia prima ater  
iride flaua.**

*Baldus.* 306. P. D. F. alis integerrimis fuscis, anticis vtrinque ocello, pupilla gemina, posticis supra ocellis quatuor, subtus sex. *Syst. Ent. App.* 329.  
Habitat in India. *Mus. D. Hunter.*

*Eumenes.* 307. P. D. F. alis integerrimis dimidiato fuscis, anticis vtrinque ocello atro tripupillato.

*Papilio Eumene.* *Cram. Inf.* 8. tab. 92. fig. F. G.

Habitat in India occidentali.

Praecedentibus paullo maior. Alae anticae fuscae margine interiori dimidiato coeruleo apiceque fasciis tribus obscurioribus. Ocellus magnus ater pupillis tribus albis in medio. Posticae coeruleae margine interiori fusco fasciis tribus versus apicem atris. Subtus omnes basi fuscae apice albae fasciis nigris ocelloque atro anticarum tripupillato, posticarum coeco.

*Ephynes.* 308. P. D. F. alis integerrimis coeruleis purpureo fasciatis, anticis vtrinque ocello, inferiori tripupillato.

*Papilio Ephyne.* *Cram. Inf.* 8. tab. 93. fig. E. F.

Habitat Surinami.

Affinis praecedenti. Alae supra coeruleae fasciis quatuor apicis purpureis. Ocellus ater pupilla alba in medio alae anticae. Subtus fuscae flavo fasciatis, anticis ocello atro pupillis tribus albis.

### \* Nymphales gemmati.

*Io.* 309. P. N. G. alis angulato dentatis fulvis nigro maculatis, singulis ocello coeruleo. *Syst. Ent.* 489. 203. *Lin.* *Syst. Nat.* 2. 769. 131. *Fn. Sv.* 1048. *Geoff. Inf.* 2. 36. 2.

*Papilio elegantissima*, singulis alis singulis maculis oculos imitantibus, perbelle depictis. *Raj. Inf.* 422. 14.

*Oculus Pavonis.* *Goed.* tab. 1. fig. 23.

*Huffnag. pict.* tab. 12. fig. 9.

*Merian. Europ.* 1. tab. 26.

*Albin. Inf.* tab. 3. fig. 4.

*Ejp. pap.* 1. tab. 5. fig. 2.

*Reaum. Inf.* 1. tab. 25. fig. 1. 2.

*Roef. Inf.* 1. Pap. 1. tab. 3.

*Bergstræff. Inf.* 2. tab. 21. fig. 6. 7.

*Harris Aurel.* tab. 8. fig. k. i.

*Wilks pap.* tab. 3. a. 2.

*Schæff. Icon.* tab. 94. fig. 1.

Habitat in Vrtica, Humulo.

*Larva* gregaria, spinosa, atra, albo punctata, pedibus posticis ferrugineis.

*Puppa* decemdentata, viridis, aureo punctata, postice bifida.

310. P. N. G. alis angulato dentatis fulvis apice nigris, *Polynice*.  
subtus fuscis, strigis coeruleis ocellisque anticarum  
sex, posticarum quinque.

*Papilio Polynice. Cram. Inf.* 17. tab. 195. fig. D. E.

Habitat in Sumatra Insula, Indiae orientalis.

Alae omnes supra fulvae apice nigrae sinubus albis. In  
posticis puncta tria nigra ante marginem et in mar-  
gine maculae quatuor fulvae.

311. P. N. G. alis angulatis fulvis nigro maculatis, posti- *Almana*.  
cis oculo maiori pupilla gemina, omnibus subtus  
brunneis. *Syst. Ent.* 490. 204.

*Papilio Almanae* alis angulatis luteis, singulis supra  
ocellis sesquialteris, anteriore pupilla gemina. *Linn.*

*Syst. Nat.* 2. 769. 132. *Mus. Lud. Vlr.* 272.

*Papilio Almanae. Cram. Inf.* 5. tab. 58. fig. F. G.

*Seligm. Av.* 3. tab. 36.

*D'Aubent. miscell.* tab. 94. fig. 3. 4.

*Edw. Av.* tab. 84.

Habitat in Asia.

312. P. N. G. alis dentatis luteo variis, posticis oculo ma- *Asterie*.  
iori, pupilla gemina, subtus pallidis ocellis subtri-  
bus. *Syst. Ent.* 490. 205.

*Papilio Asterie* alis dentatis luteo variis, singulis vtrin-  
que ocellis sesquialteris, anteriore pupilla gemina.

*Linn. Syst. Nat.* 2. 769. 133.

*Papilio Asterie. Cram. Inf.* 5. tab. 58. fig. D. E.

*Klem. Inf.* 1. tab. 5. fig. 3. 4.

*Aubent. miscell.* tab. 94. fig. 1. 2.

Habitat in Acantho spinoso Indiae orientalis.

70 CLASS. VI. GLOSSATA. Papilio.

*Oenone*. 313. P. N. G. alis denticulatis luteis margine nigris, posticis basi cyaneis. *Syst. Ent.* 490. 206.

*Papilio Oenone* alis denticulatis, primoribus albido maculatis subbiocellatis, posticis basi cyaneis, ocellis duobus. *Linn. Syst. Nat.* 2. 770. 135. *Mus. Lud. Vlr.* 274. 275.

*Papilio Oenone*. *Cram. Inf.* 3. tab. 35. fig. A. B. C.

*Kleman. Inf.* 1. tab. 3. fig. 1. 2.

*Petiv. Gazoph.* tab. 4. fig. 3.

*Edw. Av.* tab. 37.

*Seb. Musf.* 4. tab. 14. fig. 15. 16.

— — — tab. 30. fig. 17. 18.

Habitat in Asia.

Ocelli valde variant.

*Lemoni*- 314. P. N. G. alis dentatis fuscis flauo punctatis, omnibus  
as. ocello vnico. *Syst. Ent.* 490. 207. *Linn. Syst. Nat.* 2. 770. 136. *Musf. Lud. Vlr.* 277.

*Sulz. Hist. Inf.* tab. 16. fig. 7. 11.

β. *Papilio Ononis* alis angulatis griseis, primoribus fluorescenti nebulosis supra ocellis sesquialteris. *Linn. Syst. Nat.* 2. 769. 134.

*Cram. Inf.* 3. tab. 35. fig. D. E. F.

Habitat in Indiae orientalis Ruelliis. *D. Roenig.*

*Orithya*. 315. P. N. G. alis dentatis fuscis, omnibus supra ocellis duobus, anticis subtus vnico. *Syst. Ent.* 490. 208.

*Linn. Syst. Nat.* 2. 770. 137. *Musf. Lud. Vlr.* 278.

*Papilio Orithya*. *Cram. Inf.* 2. tab. 19. fig. C. D.

*Papilio Lauinia*. *Cram. Inf.* 2. tab. 21. fig. C. R. E. F.?

*Cram. Inf.* 3. tab. 32. fig. E. F.

*Seligm. Av.* 2. tab. 51.

*Sulz. Hist. Inf.* tab. 17. fig. 1. 2.

*Roef. Inf.* 4. tab. 6. fig. 2.

*Edw. Av.* 26. tab. 26.

*Seb. Musf.* 4. tab. 5. fig. 13. 14.

— — — tab. 14. fig. 15. 16.

— — — tab. 22. fig. 5. 6.

Habitat in China, in noua Hollandia. *Musf. Dom. Banks.*

Variat alis posticis supra disco coeruleo et fusco.

Conf.

CLASS. VI. GLOSSATA. Papilio. 71

Conf. *Papilio Euaete*. *Cram. Inf.* 17. tab. 203. fig. C. D.

316. P. N. G. alis dentatis supra fuscis, subtus omnibus *Penelea*. ocellis duobus.

*Papilio Penelea*. *Cram. Inf.* 9. tab. 101. fig. G.

Habitat Surinami.

Alae supra fuscae, subtus variegatae ocellis duobus antillarum minoribus.

317. P. N. G. alis dentatis supra fuscis coeruleo micantibus, anticis ocellis duobus punctisque duobus albis. *Fidia*.

*Syst. Ent.* 491. 209. *Linn. Syst. Nat.* 2. 770. 138.

*Naturf.* 6. pag. 13. 14.

*Esp. pap.* 1. tab. 49. *Suppl.* 25. fig. 3.

Habitat in Africa.

Ocelli ad marginem anteriorem, inter quos puncta duo alba.

318. P. N. G. alis dentatis supra coeruleo fusco alboque marmoratis, omnibus ocellis sex. *Feronia*. *Syst. Ent.* 491. 210.

*Papilio Feronia* alis dentatis supra coeruleo fuscoque vndatis, omnibus ocellis sex. *Linn. Syst. Nat.* 2.

770. 140. *Mus. Lud. Vlr.* 283.

*Cram. Inf.* 16. tab. 192. fig. E. F.

*Clerk. Icon.* tab. 31. fig. 1.

*Drury Inf.* tab. 10. fig. 1. 2.

*Aubent. miscell.* tab. 92. fig. 7. 8.

*Seb. Mus.* 4. tab. 38. fig. 10. 11. 15. 16.

— — tab. 40. fig. 19.

Habitat in India orientali.

Ocelli supra pupilla alba subtus coeci.

319. P. N. G. alis dentatis fuscis, vtrinque anticis sesquicoello, posticis ocellis supra tribus, subtus sex. *Macra*. *Syst. Ent.* 491. 211. *Linn. Syst. Nat.* 2. 771. 141. *Fn. Sv.* 1049.

*Papilio* alis fulvo fuscoque nebulosis, primariis sesquicoello, secundariis supra tribus, infra septem ocellis. *Geoff. Inf.* 2. 50.

70 CLASS. VI. GLOSSATA. *Papilio*.

*Papilio* media alis fuluo s. rufo et nigricante colore variis, cum ocello prope extimum angulum alarum exteriorum. *Raj. Inf.* 123. 15.

*Robert. Icon.* tab. 15. fig. 2.

*Esp. pap.* tab. 6. fig. 2.

*Degeer Inf.* 2. tab. 2. fig. 1. 2.

*Schaeff. Icon.* tab. 58. fig. 2. 3.

*Naturf.* 8. tab. 3. fig. A. B. at paullo maior.

*Seb. Mus.* 4. tab. 3. fig. 2.

*Merian. Europ.* 1. tab. 27. fig. 16. 17.

Habitat in Europae gramine.

*Larna* subuullosa, virescens, cauda bidentata.

*Puppa* virescens, obtuse bifida, lateribus echinatis.

*Papilio Hiera. Gen. Inf. Mant.* 259. \* Huius mera varietas.

*Roxelana.* 320. P. N. G. alis dentatis fuscis supra coecis, posticis subtus vndatis ocellis septem, interiori pupilla gemina.

*Papilio Roxelana. Cram. Inf.* tab. 141. fig. C. D.

Habitat in Asia. *Mus. Dom. Teats.*

Statura omnino praecedentis. Alae anticae supra fuscae disco fuluo maculisque flauis, subtus ocello atro pupilla alba. Posticae supra fuscae punctis ocellaribus tribus iride fulua, subtus griseae fusco vndatis ocellis septem atris iride flaua pupilla alba, 3. 4. minutissimis, ultimo pupilla gemina.

*Helima.* 321. P. N. G. alis angulatis flauis fusco subfasciatis, posticis vtrinque ocellis sex, subtus striga alba. *Syst. Ent.* 492. 212. \*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

*Tirra.* 322. P. N. G. alis angulato dentatis supra fuscis, fascis fulua anticisque striga punctorum alborum, postice ocellorum.

*Papilio Tirra. Drury Inf.* 2. tab. 18. fig. 3. 4.

*Papilio Tirra. Cram. Inf.* 12. tab. 158. fig. E. F.

Habitat in Sierra Leon Africae.

Alae omnes angulato dentatae. Anticae basi pallide fuscae lineolis quatuor transuersis nigris, in medio fascia communis fulua striga sanguinea, apice obscurae strigae punctis septem albis margineque strigoso. Subtus basi flavescentes fasciis obscurioribus, apice fuscae punctis ocellaribus albis. Posticae basi pallide fuscae, in medio fascia communis fulua striga nigra, tunc ocelli septem fere antiqui ferruginei iride pupillaeque nigris, subtus cinerei flauo varii punctis ocellaribus atris.

333. P. N. G. alis angulatis supra nigris, fascia vtrinque *Leodice*.  
virescente, striga punctorum ocellatorum.

*Papilio Leodice. Cram. Inf. 12. tab. 138. fig. G. H.*  
Habitat in Guinea.

Alae anticae nigrae basi maculis fuluis albo marginatis, in medio fascia sinuata, communis, virescens, quae tamen marginem crassioretm haud attingit, et in hac puncta aliquot ocellaria atra. Apice puncta aliquot alba. Posticae nigrae fascia media virescente, in qua striga punctorum nigrorum. Ad angulum anilineolae duae albae. Subtus omnes variegatae fascia virescente et in hac striga punctorum alborum iride atra.

334. P. N. G. alis dentatis luteis fusco fasciatis, anticis *Meguera*.  
ocello, posticis supra quinque subtus sex. *Syst. Ent.*

492. 213. *Linn. Syst. Nat. 2. 771. 142.*

*Wilks pap. 53. tab. 2. 6. 3.*

*Amiral. tab. 5.*

*Harris Inf. tab. 27. fig. c. f. g.*

*Esp. pap. tab. 6. fig. 3.*

*Hollar. Inf. tab. 1. fig. 3.*

*Schaeff. Icon. tab. 148. fig. 3. 4.*

Habitat in Europae graminosis.

*Larva* villosa, bicaudata.

335. P. N. G. alis dentatis fuscis luteo maculatis, vtrinque *Aegeria*.  
anticis ocello, posticis supra ocellis subtus punctis  
quatuor. *Syst. Ent.* 492. 214. *Linn. Syst. Nat. 2.*  
771. 143.



*Papilio* alis rotundatis dentatis fuscis fulvo macul  
anticis ocello unico, secundariis superne qua  
plici. *Geoff. Inf.* 2. 48. 16.

*Reaum. Inf.* 1. tab. 27. fig. 16. 17.

*Wilk. pap.* 53. tab. 2. 6. 4.

*Roef. Inf.* 4. tab. 33. fig. 3. 4.

*Edw. Av.* tab. 178.

*Sepp. Inf.* 1. 33. tab. 6.

*Schaeff. Leon.* tab. 75. fig. 1. 2.

*Seligm. Av. B.* tab. 73.

*Esp. Papil.* 1. tab. 7. fig. 1.

*Bergstr. Inf.* 2. tab. 22. fig. 6. 7.

*Harr. Inf.* tab. 41. fig. h. i. k.

*Petiv. Gazoph.* tab. 24. fig. 3.

*Raj. Inf.* 128. 5.

Habitat in Europae Graminibus.

*Larva* viridis albo lineata, cauda bispinosa.

*Puppa* brevis, obesa viridis.

*Xiphia.* 326. P. N. G. alis dentatis fuscis flavo maculatis, vtrin  
anticis ocello posticisque supra tribus subtus quat  
*Syst. Ent.* 492. 215.\*

Habitat in Madera. *Mus. Dom. Banks.*

*Minerva.* 327. P. N. G. alis dentatis fulvis apice nigris albo ma  
latis, posticis ocellis supra quinque coecis, sub  
septem pupillatis. *Syst. Ent.* 493. 216.\*

*Papilio Arcadia.* *Cram. Inf.* 10. tab. 116. fig. E. I

Habitat in America. *Dom. v. Rohr.*

*Ocypte.* 328. P. N. G. alis dentatis supra fuscis immaculatis ful  
glaucis, strigis tribus obscuris, posticis ocellis qu  
que. *Gen. Inf. Mant.* 260.\*

*Papilio Helle.* *Cram. Inf.* 17. tab. 194. fig. F. G.

Habitat Surinami. *D. Schulz.*

*Ocirrhoe.* 329. P. N. G. alis dentatis supra albis apice fuscis, ful  
fuscis, fasciis duabus albis, posticis ocellis quin  
*Gen. Inf. Mant.* 260.\*

*Papilio Cessia.* *Cram. Inf.* 17. tab. 194. fig. D. E.

*Sulz. Hist. Inf.* tab. 17. fig. 3. 4.

*Seb. Mus.* 4. tab. 38. fig. 3. 4.

Habitat Surinami. *D. Schulz.*

CLASS. VI. GLOSSATA. Papilio. 75

30. P. N. G. alis subdentatis fuscis ocello vnico, posticis *Penelope*  
subtus duobus. *Syst. Ent.* 493. 217.\*  
Habitat in America. *Dom. Drury.*
31. P. N. G. alis subdentatis fuscis, posticis coerulescentibus, subtus ocellis quinque. *Gen. Inf. Mant.* 260.\*  
*Papilio* *Lea. Cram. Inf.* 13. tab. 151. fig. C. D.?  
Habitat Surinami. *D. Schulz.*
32. P. N. G. alis angulatis pallidis strigis vndatis fuscis, *Iatro-*  
anticis ocello vnico, posticis duobus. *Syst. Ent. phae;*  
493. 218.  
*Papilio Iatrophae* alis angulatis pallidis, lineis vndu-  
latis, primoribus puncto vnico, posticis duobus.  
*Linn. Syst. Nat.* 2. 779. 172. *Mus. Lud. Vr.* 289.  
*Amoen. acad.* 6. 408. 73.  
*Gronov. Zooph.* 2. 197. 790.  
*Cram. Inf.* 17. tab. 202. fig. E.  
*Merian. Surin.* tab. 4.  
Habitat in Americae meridionalis Iatrophis, in noua  
Hollandia. *Mus. Dom. Banks.*
33. P. N. G. alis dentatis cinerascens, supra anticis *Laome-*  
ocellis sex, posticis quinque, quibusdam coecis. *dia.*  
*Syst. Ent.* 494. 219. *Linn. Syst. Nat.* 2. 772. 144.  
*Papilio Laomedia. Cram. Inf.* 1. tab. 8. fig. F. G.  
*Papilio Atlites. Linn. Amoen. acad.* 6. 407. 72.  
*Drury Inf.* 1. tab. 5. fig. 3.  
*Sulz. Hist. Inf.* tab. 16. fig. 10.  
Habitat in India orientali.
34. P. N. G. alis dentatis fuscis subtus lineatis, posticis *Libye.*  
ocellis sex, pupilla argentea. *Syst. Ent.* 494. 220.  
*Linn. Syst. Nat.* 2. 772. 146.  
*Sulz. Hist. Inf.* tab. 17. fig. 7.  
Habitat in India.  
Alae anticae subtus ocello vnico, posticae sex fuscis.
35. P. N. G. alis dentatis concoloribus griseis, singulis *Hedonia.*  
ocellis sex ferrugineis. *Syst. Ent.* 494. 221. *Linn.*  
*Syst. Nat.* 2. 774. 153. *Mus. Lud. Vr.* 279.  
*Papilio Luzonicus* fuliginosus oculatus et vndulatus.  
*Petiv. Gazoph.* tab. 39. fig. 4.

*Papilio*

76 CLASS. VI. GLOSSATA. *Papilio*.

*Papilio Hedonia*. *Cram. Inf.* 6. tab. 69. fig. C. D.

*Seb. Musf.* 4. tab. 14. fig. 5. 6.

Habitat in Asia.

*Dejanira*. 336. P. N. G. alis dentatis fuscis, anticis ocellis vtrinque quinque, posticis sex fasciaque alba repanda. *Syst. Ent.* 494. 222. *Linna. Syst. Nat.* 2. 774. 154. *Musf. Lud. Vlr.* 282.

*Papilio Achine*. *Scop. carn.* 433.

*Bergstr. Inf.* 2. tab. 37. fig. 6. 7.

*Ejp. Pap.* 1. tab. 9. fig. 2.

Habitat in Lolio temulento Germaniae.

*Tullia*. 337. P. N. G. alis dentatis fuscis maculis ocellaribus, fuscus fascia violacea, posticarum ocellis duobus.

*Papilio Tullia*. *Cram. Inf.* 7. tab. 81. fig. A. B.

Habitat in China.

Alae anticae fuscae maculis ocellaribus albis iride cor-  
rulea, posticae pallidiores maculis albis iride rufa  
flava. Subtus omnes testaceae fascia violacea, fuscus  
fusca, posticis ocellis duobus in fascia violacea.

*Pegala*. 338. P. N. G. alis dentatis fuscis, anticis fascia rufa ocellis  
loque vnico, posticis supra ocello subtus sex.

Habitat in America. *Musf. D. Hunter.*

Variat interdum ocello primo et quarto obsoletis.

*Briſeis*. 339. P. N. G. alis dentatis fuscis viridi micantibus fascia  
alba, anticis ocellis duobus, posticis coecis.

*Papilio Briſeis* alis subdentatis supra fuscis viridi micantibus, primoribus ocellis duobus, subtus nigro  
bimaculatis. *Linna. Syst. Nat.* 2. 770. 139. *Musf. Lud. Vlr.* 276.

*Naturf.* 10. tab. 2. fig. 3. 4.

Habitat in Germania.

Ocelli in pagina superiori interdum defunt.

*Semele*. 341. P. N. G. alis dentatis fuscis fuluo subfasciatis, posticis albo nigroque variis. *Syst. Ent.* 494. 224.

*Papilio Semele* alis dentatis fuluo nigroque nebulosis, anticis vtrinque ocellis duobus, posticis supra vnico  
*Linna. Syst. Nat.* 2. 772. 148. *Fn. Sv.* 1051.

*Papilio*

*Papilio maufeuila*, alis pullis cum duplici in exteri-  
bus macula lutea et duplici ocello nigro. *Raj. Inf.*  
123. 6.

*Huffn. Inf.* 2. tab. 8.

*Petiv. Gazoph.* tab. 14. fig. 9.

*Esp. pap.* 1. tab. 8. fig. 1.

*Sulz. Hist. Inf.* tab. 17. fig. 5. 6.

*Schaeff. Icon.* tab. 207. fig. 3. 4.

Habitat in Europae sylvis.

Alae anticae subtus flavescentes.

1. P. N. G. alis dentatis fuscis, fascia pallida, anticis *Hermio-*  
ocellis supra duobus, subtus vnico. *Syst. Ent.* 495. *ne.*  
225.

*Papilio Hermione* alis dentatis fuscis fascia pallida pri-  
moribus ocello, posticis supra puncto. *Linn. Syst.*  
*Nat.* 2. 773. 149. *Mus. Lud. Vlr.* 281.

*Papilio* alis rotundatis nigrofuscis, omnibus fascia al-  
bida, primariis ocello duplici, secundariis vnico.  
*Geoff. Inf.* 2. 46. 13.

*Papilio lusitanicus* oculatus et marmoratus. *Petiv. Ga-*  
*zoph.* 12. tab. 7. fig. 5.

*Papilio Fagi. Scop. carn.* 428.

*Esp. pap.* 1. tab. 8. fig. 2.

*Schaeff. Icon.* tab. 274. fig. 5. 6.

— — — tab. 277. fig. 1. 2.

*Roef. Inf.* 3. tab. 34. fig. 5. 6.

Habitat in Germania, Gallia.

Similis *P. Semele*, at paullo maior.

In nostro specimine ocellus parvus vtrinque in alis po-  
sticis.

2. P. N. G. alis dentatis fuscis, fascia vtrinque alba, an- *Circe.*  
ticis ocello vnico. *Syst. Ent.* 495. 226.

*Papilio Proserpina* alis dentatis nigris, fascia candida,  
omnibus subtus ocello, primoribus supra ocello coe-  
co, *Wien. Verz.* 299. tab. 1. a. 9. 1. 6. 9.

*Papilio Viliada. Naturf.* 6. 17. 3.

*Schaeff. Icon.* tab. 82. fig. 1. 2.

Habitat in Europae Gramine.

*Larva* nuda lineis longitudinalibus fuscis albis luteis,  
cauda furcata.

78 CLASS. VI. GLOSSATA. Papilio.

*Phaedra*. 343. P. N. G. alis dentatis fuscis concoloribus, ant  
ocellis duobus violaceis. *Syst. Ent.* 495. 227. *L.*  
*Syst. Nat.* 2. 773. 150. *Mus. Lud. Vlr.* 280.

*Papilio Dryas*. *Scop. carn.* 429.

*Esp. pap.* tab. 40. fig. 1. 2.

*Bergstr. Inf.* tab. 10. fig. 3. 4.

*Schaeff. Icon.* tab. 210. fig. 1. 2.

Habitat in Europa.

*Periboea*. 344. P. N. G. alis dentatis fuscis, anticis oculo obsolet  
subtus striga flavescente, posticis punctis ocellarib  
atris. *Gen. Inf. Mant.* 261.\*

Habitat Surinam. *D. Schulz.*

*Arfinoe*. 345. P. N. G. alis caudato dentatis fulvis nigro maculat  
posticis ocellis vtrinque duobus.

*Papilio Arfinoe*. *Cram. Inf.* 13. tab. 160. fig. B. C.

*Seb. Mus.* 4. tab. 19. fig. 7. 8.

— — — tab. 37. fig. 11. 12.

Habitat in Amboina.

Alae omnes vtrinque fulvae maculis strigisque vari  
nigris, ocelli duo nigri in alis posticis, superiorib  
pupilla coerulea, inferioribus alba. Punctum albu  
subtus ad angulum ani.

*Merops*. 346. P. N. G. alis dentatis fulvis, apice nigris, oculo vn  
co, posticis subtus cinereis ocellis tribus minutis. *Syst. Ent.* 495. 228.\*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

*Ligea*. 347. P. N. G. alis dentatis fuscis, fascia rufa, vtrinq  
anticis ocellis quatuor, posticis tribus. *Syst. Ent.*  
495. 229. *Linn. Syst. Nat.* 2. 772. 144. *Fa. S*  
1050.

*Alb. Inf.* tab. 5. fig. 1.

*Esp. Papil.* 1. tab. 7. fig. 2.

*Schaeff. Icon!* tab. 183. fig. 2. 3.

*Naturf.* 8. tab. 3. fig. 6.?

*Degeer Inf.* 2. tab. 2. fig. 7. 8.

Habitat in Europae syluis.

*Iole*. 348. P. N. G. alis dentatis fulvis nigro punctatis, postic  
subtus ocellis quinque, ultimo bipupillato.

*Papil*

*Papilio Laodice*. Cram. Inf. 14. tab. 157. fig. E. F.  
Habitat in Guinea.

Corpus fulvum. Alae anticae fulvae punctis quinque strigisque marginalibus fuscis, subtus magis strigisae. Posticae fulvae macula magna bascos fusca, punctis duobus anguli ani strigisque marginalibus fuscis. Subtus flauescentes fascia coerulea ocellisque tribus duobusque approximatis.

349. P. N. G. alis dentatis albo nigroque variis, subtus *Galathea* anticis ocello unico, posticis quinque obsoletis. *Syst. Ent.* 496. 230. Linn. *Syst. Nat.* 2. 772. 147.

*Papilio* alis rotundatis albis lineis maculisque nigris pulcre tessellatis. *Geoff. Inf.* 2. 74. 46. tab. 11. fig. 3. 4.

*Papilio* media, alis albo et nigro coloribus pulcre variegatis. *Raj. Inf.* 116. 9.

*Papilio* leucamelanos. *Petiv. Mus.* 3. 3. tab. 1. fig. 3.

*Wilks pap.* 52. tab. 2. 6. 1.

*Roef. Inf.* 3. tab. 37. fig. 1. 2.

*Schaeff. Icon.* tab. 98. fig. 7. 8. 9.

*Bergstr. Inf.* 2. tab. 16. fig. 1. 2.

*E/p. Pap.* 1. tab. 7. fig. 3.

— — — tab. 25. suppl. 1.

*Harris Aurel.* tab. 11. fig. k. 1.

Habitat in Europae Gramine.

*Larva* viridis, capite spinisque duabus caudalibus carneis. *Roef. Inf.* 3. tab. 70. fig. 6.

*Puppa* nuda, gibba, maculata *Fuesl. Magaz.* 1. tab. 2. fig. 3.

350. P. N. G. alis dentatis fuscis, subtus striga obliqua alba, anticis ocellis duobus, posticis sesequitertiis. *Hesione*. *Gen. Inf. Mant.* 260.\*

*Papilio Hesione*. Cram. Inf. 1. tab. 11. fig. C. D.

*Sulz. Hist. Inf.* tab. 17. fig. 3. 4.

Habitat Surinami.

351. P. N. G. alis dentatis, supra nigris, fascia communi alba, posticis subtus ocellis duobus. *Syst. Ent.* 496. 231.\* *Egata*.

Habitat in America. *Mus. D. Hunter*.

- Serina.* 352. P. N. G. alis subdentatis, supra viridibus, posticis fuscis disco albo, ocellis duobus. *Syst. Ent.* 497. 23.  
Habitat in Iamaica. *Mus. D. Hunter.*
- Doris.* 353. P. N. G. alis dentatis fuscis, subtus coeruleis nigro fasciatis, ocellis trium parium. *Gen. Inf. Mant.* 162.  
*Papilio Doris.* *Cram. Inf.* 1. tab. 8. fig. B. C.  
Habitat Surinami.
- Ianthus.* 354. P. N. G. alis dentatis, supra fuscis albo subfasciatis, posticis subtus ocellis quinque coecis.  
Habitat Cayennae. *Mus. Dom. Teats.*  
Statura praecedentium. Alae omnes supra fuscae, antice punctis, posticis fascia maculari albis. Subtus anticae basi flavae, apice fuscae albo maculatae, posticae fuscescentes maculis, fascia lunulisque albis. Pone medium ocelli quinque atri iride flava absque pupilla.
- Pilosellae.* 355. P. N. G. alis dentatis fuscis, disco fulvo, anticis utrinque ocello nigro, pupilla gemina. *Syst. Ent.* 497. 233.  
*Papilio* alis rotundatis fulvis, oris fuscis, primariis ocello duplici continuo, secundariis duobus parvis, infra fusco cinereoque nebulosis. *Geoff. Inf.* 1. 52. 60.  
*Papilio Tithonius* alis subdentatis concoloribus disco luteis, primoribus ocello bipupillato, posticis punctis duobus albis. *Linn. Mant.* 1. 537.  
*Sepp. Inf.* 1. 21. tab. 3.  
*E/p. pap.* 1. tab. 9. fig. 1.  
*Bergstr. Inf.* 2. tab. 18. fig. 5. 6. 7. 8. 9.  
*Naturf.* 3. tab. 2. fig. a. b. c.  
Habitat in Plantagine, Pilosella Germaniae.  
*Larva* virescens, albolineata, capite fusco.  
*Variat* alis posticis supra uniocellatis et coecis.  
*Conf. Papilio Lycaon.* *Naturf.* 3. tab. 2. fig. d.  
*Schaeff. Icon.* tab. 213. fig. 1. 2.  
Vix satis a P. Pilosellae distinctus.
- Cadma.* 356. P. N. G. alis dentatis fulvis, posticis subtus disco albo, ocellis duobus, pupilla didyma coerulea. *Syst. Ent.* 497. 234.\*

## CLASS. VI. GLOSSATA. Papilio.

*Papilio Cadma. Drury Inf.* 2. tab. 18. fig. 1. 2.  
Habitat in America meridionali.

157. P. N. G. alis angulato dentatis, posticis supra fuscis *Laethes*.  
puncto albo, subtus ocellis sex coeruleis.

*Papilio Laethes. Cram. Inf.* 11. tab. 132. fig. A. B.  
Habitat Surinami.

Alae anticae fulvae, apice fuscae, subtus punctis duobus  
albis. Posticae supra fuscae puncto parvo albo, sub-  
tus variegatae puncto baseos albo, fasciaque media  
brunnea et in hac ocelli sex coeruli pupilla alba.

158. P. N. G. alis dentatis fuscis, anticis subtus luteis ocel- *Iantira*.  
lo vtrinque vnico, posticis subtus punctis tribus.  
*Syst. Ent.* 497. 235. *Linn. Syst. Nat.* 2. 774. 156.  
*Fn. Sv.* 1053.

*Papilio* alis rotundatis fuscis primariis subtus fulvis  
ocello vnico. *Geoff. Inf.* 2. 49. 17.

*Papilio* media alis pullis. *Raj. Inf.* 124. 16.

*Esp. Pap.* 1. tab. 10. fig. 1.

*Bergstr. Inf.* 2. tab. 23. fig. 5.

*Holl. Inf.* tab. 10. fig. 1.

*Alb. Inf.* tab. 53.

*Schaeff. Icon.* tab. 273. fig. 1. 2. 5. 6.

Habitat in Europae sylvis.

159. P. N. G. alis dentatis fuscis, anticis supra litura fla- *Iurtina*.  
ua, ocello vtrinque vnico, posticis coecis. *Syst. Ent.*  
498. 236. *Linn. Syst. Nat.* 2. 774. 155. *Fn. Sv.* 152.

*Papilio* alis rotundatis dentatis supra fuscis, primariis  
macula fulva, infra e fuluo et fusco undulatis, pri-  
moribus vtrinque ocello. *Geoff. Inf.* 2. 50. 18.

*Papilio* media, alis exterioribus superne media parte  
rufis, oris pullis. *Raj. Inf.* 114. 17.

*Esp. pap.* 1. tab. 10. fig. 2.

*Hufn. Inf.* 1. tab. 1. fig. 1.

*Harr. Inf.* tab. 32. fig. c. d. p.

*Alb. Inf.* tab. 53.

*Reaum. Inf.* 1. tab. 11. fig. 1.

*Wilk. pap.* 33. tab. 2. 6. 2.

*Roes. Inf.* 3. tab. 34. fig. 7. 8.

*Sepp. Inf.* 1. 29. tab. 5.

*Fabr. Sp. Inf.* T. II.

F

Habl.



Habitat frequens in Europae graminosis.

*Larva* villosa viridis, linea laterali alba, cauda bifida

*Puppa* flavescentes.

Varietas sexus praecedentis videtur.

*Tiphone*. 360. P. N. G. alis dentatis fuscis, anticis ocellis utrinque duobus, posticis sex, pupilla violacea. *Gen. In Mant.* 263.\*

*Papilio Tiphone*. *Naturf.* 6. 18. tab. 1. fig. 1. 2.

Habitat in Germaniae graminosis.

*Gonerilla*. 361. P. N. G. alis dentatis nigris albo maculatis, fasci communi rufa, posticis ocellis quatuor. *Syst. Ent.* 498. 237.\*

Habitat in noua Zelandia. *Mus. Dom. Banks.*

*Itea*. 362. P. N. G. alis dentatis nigris, anticis fascia punctisque flauis, posticis disco rufo, ocellis quatuor. *Syst. Ent.* 498. 238.\*

Habitat in noua Zelandia. *Mus. Dom. Banks.*

*Portlandia*. 363. P. N. G. alis dentatis fuscis, posticis supra ocellis quinque coecis, subtus septem pupillatis.

Habitat in America meridionali. *Mus. Dom. Teats.*

*Paruus*. Alae anticae fuscae versus apicem fascia flavescente maculis tribus ocellaribus atris. Posticae fuscae maculis quinque atris iride flavescente aboque pupilla. Subtus fuscae strigis obscurioribus. Anticae versus apicem fascia lata apice bifida alba et in hac ocelli quatuor atris iride flaua, exterioribus pupilla alba. Posticae fascia versus apicem alba et ocello unico ante fasciam, sex pone fasciam atris ultimis duobus connatis iride flaua pupillaque oblonga argentea.

*Cardui*. 364. P. N. G. alis dentatis fuluis albo nigroque variegatis, posticis subtus ocellis quatuor. *Syst. Ent.* 499. 239. *Linn. Syst. Nat.* 2. 774. 157. *Fn. Sv.* 1054. *Geoff. Inf.* 2. 41. 7.

*Papilio* maior pulkra, nigro, albo, rubro coloribus varia. *Raj. Inf.* 122. 13.

*Papilio Cardui*. *Cram. Inf.* 3. tab. 26. fig. E. F.

*Esp. pap.* 1. tab. 10. fig. 3.

*Huf-*

*Hufn. Inf.* tab. 7. fig. 3.

*Alb. Inf.* tab. 56.

*Reaum. Inf.* 1. tab. 26. fig. 11. 12.

*Holl. Inf.* tab. 2. fig. 2. 3.

*Seb. Mus.* 4. tab. 1. fig. 2. 3. 4.

*Harris Aurel.* tab. 11. fig. c. f.

*Wilks Inf.* 56. tab. 3. a. 3.

*Roes. Inf.* 1. pap. 1. tab. 10.

*Schaeff. Icon.* tab. 97. fig. 5. 6.

Habitat in Europae, Africae carduis.

*Larva* solitaria spinosa fusca lineis flauis, lateralibus interruptis.

*Pappa* fusca, lineis cinereis punctisque aureis.

365. P. N. G. alis subangulatis fuluis albo nigroque variis *Huntera.*  
posticis subtus albo reticulatis ocellisque duobus.

*Syst. Ent.* 499. 240.

*Papilio Cardui Virginienfis.* *Drury Inf.* 1. tab. 5.  
fig. 3.

*Papilio Belladonna* Virginiana oculis subtus minoribus.

*Petiv. Gazoph.* Dec. 4. tab. 33. fig. 5.

*Papilio Iole. Cram. Inf.* 1. tab. 12. fig. E. F.

Habitat in Americae Balsamita.

Præcedenti affinis, at distinctus.

*Larva* viridis, nigro annulata.

366. P. N. G. alis dentatis fuscis, anticis subtus ocellis *Allionia.*  
duobus, posteriori coeco.

Habitat in Lusitania. *D. Gray.*

Magnitudo et statura P. Iurtinae. Corpus fuscum immaculatum. Alae dentatae, fuscae ocellis paginae inferioris obscure perspicuis absque pupilla, subtus basi fuscae strigis duabus obscurioribus, apice pallidiores, anticis ocellis duobus magnis, anteriori pupilla alba, posteriori coeco. Inter ocellos macula alba. Posticae loco ocellorum punctis tribus albis maioriq. atro.

367. P. N. G. alis dentatis subconcoloribus fuscescentibus, *Tulbag-*  
fascia lutea, posticis ocellis quinque. *Syst. Ent.* 499. *hia.*

241. *Lin. Syst. Nat.* 2. 775. 158. *Mus. Lud.*  
*Plr.* 284.

84 CLASS. VI. GLOSSATA. Papilio.

*Seb. Mus.* 4. tab. 19. fig. 11. 12. ?

*Cram. Inf.* 1. tab. 3. fig. E. F.

Habitat ad Cap. Bon. Spei.

*Pipileis.* 368. P. N. G. alis crenatis nigris concoloribus, post disco fulvo; ocellis septem. *Syst. Ent.* 499. 1

*Linn. Syst. Nat.* 2. 775. 159. *Mus. Lud. Vlr.* 28

*Papilio Pipileis. Cram. Inf.* 5. tab. 60. fig. A. B.

*Clerk. Icon.* tab. 26. fig. 2.

*Seb. Mus.* 4. tab. 44. fig. 16. 17.

— — — tab. 46. fig. 13. 14.

Habitat in Indiis.

*Claudia.* 369. P. N. G. alis crenatis nigris, arca sanguinea, post cis subtus marmoratis, ocellis septem. *Gest. Mant.* 263.\*

*Papilio Claudia. Naturf.* 9. tab. 2.

Habitat in India. *D. Schultz.*

*Blomfieldia.* 370. P. N. G. alis dentatis flavis apice nigris, posticis fuscis albo vndatis, ocellis quatuor intermedium coecis.

Habitat in Brasilia. *Mus. Dom. Blomfield.*

Magnus. Alae omnes fulvae anticis apice nigris, et ruleo micantibus maculis tribus albis, posticis stri marginali. Subtus anticae subconcolores, posticae fuscae lineolis annulisque albis vndatae. Vnde marginem posticum ocelli quatuor exteriori atro et pillula alba, iride ferruginea; intermediis duobus rescentibus coecis iride grisea, interiori testaceo et pillula alba iride atra.

*Bankia.* 371. P. N. G. alis angulatis supra fuscis, anticis disco fuscescente, ocello atro; pupilla gemina. *Syst. Ent.* 499. 24.\*

*Papilio Ismene. Cram. Inf.* 3. tab. 26. fig. A. B.

Habitat in nova Hollandia. *Mus. Dom. Banks.*

*Nolandra.* 372. P. N. G. alis angulatis fuscis, anticis ocello sesquialtero, posticis supra duobus, subtus quatuor. *Syst. Ent.* 500. 244.\*

Habitat in Insula Otaheity maris pacifici. *Mus. Dom. Banks.*

373. P. N. G. alis dentatis fuscis, anticis ocello, posticis supra duobus, subtus septem. *Polydecta*.

*Papilio Polydecta*. Cram. Inf. 12. tab. 144. fig. E. F.  
Habitat in Malabaria.

Alae anticae fuscae ocello magno atro iride flava pupillaeque alba, subtus basi fuscae striga alba, apice palidiores ocellis quatuor margineque strigoso.

374. P. N. G. alis dentatis atris, posticis subtus strigis ferrugineis coeruleisque, ocellis quinque, postico maiori. *Tolumnia*.

*Papilio Tolumnia*. Cram. Inf. 11. tab. 130. fig. F. G.  
Habitat in India orientali.

Paruus. Alae, ut videtur, dentatae supra atrae, anticis immaculatis, posticis macula coerulefcente puncto atro. Subtus anticae fuscae strigis ferrugineis mediaeque coerulea ocelloque paruo apicis. Posticae coerulefcentes strigis duabus baseos ferrugineis, tunc ocelli quinque atri pupilla alba, postico maiori pupilla didyma, denique strigae duae coeruleae et flavae ad marginem.

375. P. N. G. alis dentatis fuscis, posticis ocellis utrinque duobus punctisque tribus albis. *Morna*.  
*Syst. Ent.* 500. 245.\*

Habitat in India. *Mus. D. Hunter*.

376. P. N. G. alis angulatis fuscis, anticis supra ocello geminato, subtus griseo reticulatis, posticis ocellis sex. *Leda*.  
*Syst. Ent.* 500. 246. *Linn. Syst. Nat.* 2. 773. 151.

*Papilio Leda*. Cram. Inf. 17. tab. 196. fig. C. D.

*Edw. Av.* 297.

*Drury Inf.* 1. tab. 15. fig. 5. 6.

*Seb. Mus.* 4. tab. 42. fig. 5. 6.

Habitat in Africae Sierra Leon.

Variat alis subtus obscure ferrugineis immaculatis.

377. P. N. G. alis subdentatis fuscis, posticis postice atris albo coeruleoque punctatis, subtus biocellatis. *Lena*.  
*Syst. Ent.* 514. 301. *Linn. Syst. Nat.* 2. 784. 206.

*Papilio Lena*. Cram. Inf. 17. tab. 198. fig. D. E.

*Klemann. Inf.* 1. tab. 10. fig. 3. 4.

Habitat in America.

*Dyndime*. 378. P. N. G. alis dentatis fuscis, posticis supra violaceis, punctis quatuor albis.

*Papilio Dyndimene*. Cram. Inf. 17. tab. 198. fig. F. G.

*Papilio Lamia*. Sultz. Hist. Inf. tab. 18.

Habitat Surinami.

Affinis praecedenti. Alae subtus griseae atomis fuscis strigisque tribus obscuris. Anticae punctis duobus apicis albis, posticis puncto ocellari baseos punctis que quinque apicis albis.

*Iphita*. 379. P. N. G. alis angulato dentatis fuscis, obscurius fasciatis, omnibus subtus ocellis quinque.

*Papilio Iphita*. Cram. Inf. 18. tab. 209. fig. C. D.

Habitat in China.

Statura et magnitudo P. Ledaec. Alae anticae vtrunque puncto albo versus apicem et ocelli duo parui fere obsoleti in pagina superiori alae posticae.

*Europa*. 380. P. N. G. alis angulatis fuscis, posticis subtus fascia ocellari albida. Syst. Ent. 500. 247.\*

*Papilio Beroë*. Cram. Inf. 7. tab. 79. fig. C. D.

Habitat in America. Mus. Dom. Monson.

*Calisto*. 381. P. N. G. alis dentatis atris, posticis fascia sanguinea, ocellis supra quinque subtus septem. Gen. Inf. Mant. 263.\*

*Papilio Calisto*. Cram. Inf. 2. tab. 24. fig. A. B.

Aubert. miscell. tab. 20. fig. 2. 3.

Seb. Mus. 4. tab. 44. fig. 12. 13.

Habitat in Africa.

*Iris*. 382. P. N. G. alis dentatis vtrunque fascia albida interrupta ocelloque unico. Syst. Ent. 501. 248.

*Papilio Iris* alis subdentatis griseis, fascia vtrunque alba interrupta, posticis supra vniocellatis. Linn. Syst. Nat. 2. 775. 161.

*Papilio* alis angulatis supra nigro violaceis, albo fasciatis, subtus fulvo fusco albidoque variis, singulis ocello nigro coeruleo. Geoff. Inf. 2. 61. 29.

*Papilio maior nigra* s. pulla. Raj. Inf. 126. 2.

Harr. Aurel. tab. 3.

Roef. Inf. 3. tab. 42.

*Wilks Inf.* 63. tab. 1. a. 2.

*Ledermull. microsc.* 89. tab. 49.

*Sulz. Inf.* tab. 14. fig. 86.

*Esp. pap.* 1. tab. 11. fig. 1.

— — — tab. 25. Suppl. 1. fig. 4.

— — — tab. 43. Suppl. 19. fig. 1.

*Schaeff. Icon.* tab. 52. fig. 1. 2. 3.

Habitat in Salice, Fraxino.

Variat colore nigro et flavo, at semper colore coerulescens.

*Larva* scabra, viridis, lineis obliquis albis, capite spinis duabus. *Roef. Inf.* 4. tab. 31. fig. 6.

*Puppa* compressa, bicornis, virescens. *Kleeman. Inf.* 1. tab. 9. fig. 3.

\* Nymphales phalerati.

33. P. N. P. alis dentato caudatis, anticis basi rubris apice nigris albo maculatis. *Syst. Ent.* 501. 249.\* *Acheronta.*

*Papilio Cadmus. Cram. Inf.* 2. tab. 22. fig. A. B.

Habitat in Brasilia. *Dom. Drury.*

34. P. N. P. alis dentato caudatis rufis, striga margineque postico fuscis. *Syst. Ent.* 502. 250.\* *Troglo-dita.*

Habitat in America. *Mus. D. Hunter.*

35. P. N. P. alis bicaudato dentatis fulvis, strigis nigris, posticis subtus glaucis nigro punctatis. *Gen. Inf.* 264.\* *Thetys.*

*Papilio Petreus. Cram. Inf.* 8. tab. 87. fig. D. E.

Habitat in America meridionali. *Dom. Spengler.*

36. P. N. P. alis dentatis fuscis albo fasciatis maculatisque, subtus luteis albo fasciatis, maculis coerulescentibus. *Syst. Ent.* 502. 251. *Linn. Syst. Nat.* 2. 776. 162. *Fn. Sv.* 1055. *Populi.*

*Papilio maior nigra* f. pulla, alis supina parte maculis albis notatis. *Raj. Inf.* 126. 2.

*Vdm. Diff.* 54. fig. 15.

*Clerk. At. Holm.* 1753. 278. tab. 7.

*Schaeff. Icon.* tab. 40. fig. 8. 9.

*Seb. Mus.* 4. tab. 12. fig. 25. 26. 27 - 30.

*Roef. Inf.* 3. tab. 33. fig. 1. 2.

— — — 4. tab. 31. fig. 1. 2.

*Esp. papil.* 1. tab. 12. fig. 1.

88 **CLASS. VI, GLOSSATA. Papilio.**

*Bergstr.* 2, tab. 15, fig. 1. 2.

— — — tab. 19, fig. 1. 2.

Habitat in populo tremula.

*Larva* spinosa, variegata, capite caudaque fulvis.

*Puppa* flavescent, nigro punctata postice gibbere notata.

*Chrysis* 387. P. N. P. alis dentatis nigris, area communi rufa, posticis subtus fuscis atro maculatis. *Syst. Ent.* 503. 252.

Habitat in noua Hollandia. *Mus. Dom. Banks.*

*Cydippe* 388. P. N. P. alis dentatis nigris albo maculatis, area communi rufa, subtus basi testaceis nigro coerulesque variis. *Syst. Ent.* 503. 253. *Linn. Syst. Nat.* 2, 776. 163. *Amoen. acad.* 6. 409. 76.

*Papilio Ina. Cram. Inf.* 6, tab. 62, fig. A. B.

*Clerk. Icon.* tab. 36, fig. 1.

Habitat in India.

*Athaea* 389. P. N. P. alis dentatis fuscis, fascia strigaque angulato dentatis albis.

*Papilio Athaea. Cram. Inf.* 8, tab. 89, fig. E. F.

Habitat in Guinea.

Statura et magnitudo praecedentis. Alae basi fuscas anticis maculis duabus, posticis vnica obscure ferrugineis. In medio fascia angulato dentata alba, post hanc striga angulato dentata alba nigrae innata. Subtus flavescentes lituris atris fascia strigisque iisdem.

*Peuthesilea* 390. P. N. P. alis dentatis atris albo maculatis, posticis subtus fasciis flauis albaque nigro maculatis.

*Papilio Peuthesilea. Cram. Inf.* 13, tab. 145, fig. B. C.

*Papilio Biblis. Cram. Inf.* 15, tab. 175, fig. A. B.

*Papilio Biblis. Drury Inf.* 1, tab. 4, fig. 2.

Habitat in India.

Affinis P. Cydippe, Alae omnes atrae lunulis albis. Area communis rufa punctis marginalibus atris. Subtus anticae basi maculis tribus transuersis nigro foveatis, apice variegatae. Margo omnis niger striga flexuosa flaua.

*Smire* 391. P. N. P. alis dentatis subconcoloribus fuscis viridi fuluoque maculatis basi fulvis nigro punctatis.

*Papi-*

*Papilio Semire. Cram. Inf.* 17. tab. 194. fig. B. C.

Habitat in Sierra Leon Africae.

Affinis *Heliconiis*, *Violae*, *Terpsicore*, at alae dentatae.

Alae omnes basi fuluae punctis nigris, in medio fascia lata macularis viridis et postice fascia vnica alae anticae et duae alae posticae e maculis fuluis. Subtus concolores, at posticae fascia tantum vnica maculari alba.

392. P. N. P. alis dentatis nigris, anticis fascia, posticis Cyane. disco nigro punctato albis. *Syst. Ent.* 503. 254.

*Papilio Cyane. Drury Inf.* 1. tab. 4. fig. 1.

Habitat in India.

393. P. N. P. alis dentatis nigris, fascia communi alba, posticis subtus basi flauis nigro striatis. *Syst. Ent.* 503. 255. \*

Habitat in Sierra Leon. *Mus. Dom. Banks.*

394. P. N. P. alis angulatis nigris, limbo albida. *Syst. Ent.* 503. 256. *Linn. Syst. Nat.* 2. 776. 165. *Fn. Sv.* 1056. *Geoff. Inf.* 2. 35. 1. *Antiope.*

*Papilio maxima nigra*, alis vtrisque limbo lato albo cinctis. *Raj. Inf.* 135. 136.

*Schaeff. Elem.* tab. 94. fig. 1.

— — *Icon.* tab. 70. fig. 1. 2.

*Degeer Inf.* 1. tab. 21. fig. 8. 9.

*Wilk. pap.* 58. tab. 2. a. 10.

*Sulz. Inf.* 1. tab. 14. fig. 85.

*Roef. Inf.* 1. pap. 1. tab. 1.

*Esp. pap.* 1. tab. 12. fig. 2.

*Bergstraeff.* 2. tab. 39. fig. 1. 2. 3. 4.

*Seb. Mus.* 4. tab. 32. fig. 1. 2.

Habitat in Europae, Americae *Betula alba*, *Salice*,

*Larua* gregaria spinosa nigra, maculis dorsalibus quadratis ferrugineis.

*Puppa* dentata nigra fuluo punctata.

*Imago* variat aetate, limbo albo et flauescente.

395. P. N. P. alis dentatis fuluis, limbo nigro albo maculato. *Protogeneia.*

*Papilio Protegenia. Cram. Inf.* 16. tab. 139. fig.

F. G.

Habitat in Insula Iaua.



90 CLASS. VI. GLOSSATA. *Papilio*.

Alae valde dentatae sinibus inter dentes albis, fulvis  
limbo late fusco maculis aliquot albis. Subtus  
fasciae ferrugineo irroratae, anticis macula apice  
posticis puncto albo fasciaque cinerea.

*Ceres*. 396. P. N. P. alis dentatis nigris, anticis fascia alba pos-  
ticis disco coerulescente, puncto atro. *Syst. Ent.* 504.  
257. \*

Habitat in Sierra Leon Africae. *Mus. Dom. Banks.*

*Atalanta*. 397. P. N. P. alis dentatis nigris albo maculatis, fascia  
communi purpurea, anticis vtrinque, posticis mar-  
ginali. *Syst. Ent.* 504. 258. *Linn. Syst. Nat.* 2. 778.  
175. *Fn. Sv.* 1060.

*Papilio* alis nigris albo maculatis, omnibus fasciis  
cincta coccinea. *Geoff. Inf.* 2. 40. 6.

*Papilio* maior nigricans, alis maculis rubris et  
pulcre illustratis. *Raj. Inf.* 126. 1.

*Albin. Inf.* tab. 3. fig. 4.

*Reaum. Inf.* 1. tab. 10. fig. 8. 9.

*Degeer Inf.* 1. tab. 22. fig. 5.

*Harris Aur.* tab. 6. fig. a — h.

*Roef. Inf.* 1. pap. 1. tab. 6.

*Wilk. pap.* 55. tab. 3. a. 1.

*Ammir. Inf.* tab. 24.

*Sepp. Inf.* 1. tab. 1.

*Esp. pap.* 1. tab. 14. fig. 1.

*Bergsraeff. Inf.* 2. tab. 20. fig. 1 — 11.

*Schaeff. Icon.* tab. 148. fig. 1. 2.

Habitat in Europae, Asiae, Americae Vrticis.

*Larva* solitaria, spinosa, virescens linea laterali  
fasciata.

*Puppa* dentata nigricans subtus cinerea, punctis  
reus.

Conf. *Papilio Atalanta*. *Cram. Inf.* 7. tab. 84. fig.  
E. F.

*Charonia*. 398. P. N. P. alis dentatis nigris, fascia coerulescente. *Syst.*  
*Ent.* 504. 259. \*

*Papilio Charonia*. *Cram. Inf.* 4. tab. 47. fig. A. B. C.

*Drury Inf.* 1. tab. 15. fig. 1. 2.

Habitat in India.

90. P. N. P. alis dentatis fuscis, omnibus subtus striga punctorum alborum. *Echo.*

*Papilio Echo. Cram. Inf. 5. tab. 57. fig. C. D.*

Habitat Surinami.

Alae supra fuscae macula anticarum versus apicem ferruginea.

91. P. N. P. alis dentatis fuscis, fascia communi fulva, anticarum punctis duobus nigris. *Syst. Ent. 504. 260. \**

Habitat in noua Hollandia. *Mus. Dom. Banks.*

92. P. N. P. alis dentatis concoloribus atris, posticis fascia maculari sanguinea. *Syst. Ent. 505. 261. \**

Habitat in America.

93. P. N. P. alis dentatis fuscis, disco angulato fuluo nigro punctato, subtus maculis baseos flauis. *Ostania.*

*Papilio Ostania. Cram. Inf. 12. tab. 135. fig. B. C.*

Habitat in Sierra Leon Africae.

Magnitudo P. Biblis. Corpus fuluum annulis abdominalibus nigris. Limbus alarum supra fuscus lunulis marginalibus albis, anticarum punctis aliquot fuluis coeruleisque, posticarum strigis atris anguloque anticoeruleo. Discus varie angulatus, fuluus punctis aliquot nigris. Subtus omnes basi nigrae flauo maculatae, in medio albae fuluo marginatae strigaeque punctorum nigrorum. Margo posticus ater strigis tribus lunularum albidarum.

94. P. N. P. alis dentatis obscure virescentibus albo nigro rufoque maculatis. *Lubentina.*

*Papilio Lubentina. Cram. Inf. 13. tab. 155. fig. C. D.*

Habitat in China.

Alae anticae subconcolores obscure virescentes punctis duobus baseos nigris, in medio maculae duae transversae rufae, utrinque nigro marginatae, albam includentes, tunc fascia macularis alba et denique apex fusco striatus punctis quatuor albis. Posticae itidem virescentes maculis tribus baseos nigris fasciisque duabus apicis e maculis nigris, quibusdam nucleo rufo foetis, subtus pallidiores, basi nigro maculatae, apice

apice fasciis duabus e maculis rufis nigro marginatis.

*polychloros*. 404. P. N. P. alis angulatis fulvis nigro maculatis, antice supra punctis quatuor nigris. *Syst. Ent.* 505. 26 *Linn. Syst. Nat.* 2. 777. 166. *Fn. Sv.* 1057. *Geoff. Inf.* 2. 37. 3.

*Papilio vtricarum* referens maior, quam *ulmarum* vocant soliti sumus. *Raj. Inf.* 118. 2.

*Merian. Europ.* 52. tab. 52.

*Alb. Inf.* tab. 55.

*Frisch. Inf.* 6. tab. 3.

*Reaum. Inf.* 1. tab. 23. fig. 2.

*Ammiral. Inf.* tab. 15.

*Roes. Inf.* 1. pap. 1. tab. 2.

*Wilk. pap.* 56. tab. 3. a. 5.

*Esp. pap.* 1. tab. 13. fig. 1.

*Bergstræff. Inf.* 2. tab. 30. fig. 1 — 3.

*Schæff. Icon.* tab. 146. fig. 1. 2.

*Seb. Mus.* 4. tab. 1. fig. F.

Habitat in Pomonae arboribus.

*Larva* gregaria, spinosa, nigricans, linea laterali fissa

*Puppa* incarnata.

*Vrticariae*. 405. P. N. P. alis angulatis fulvis nigro maculatis, antice supra punctis nigris tribus. *Syst. Ent.* 505. 263. *Linn. Syst. Nat.* 2. 777. 167. *Fn. Sv.* 1058. *Geoff. Inf.* 2. 37. 4.

*Papilio vrticaria* vulgarissima, rufo nigro coerulesco albo coloribus varia. *Raj. Inf.* 117. 1.

*Robert. Icon.* tab. 5.

*Huffn. pict.* 2. fig. 16.

*Bradt. nat.* tab. 27. fig. 2.

*Merian. Europ.* 44. tab. 44.

*Albin. Inf.* tab. 4. fig. 51.

*Swamm. Bibl. nat.* tab. 35. fig. 12.

*Esp. pap.* 1. tab. 13. fig. 2.

*Bergstræff. Inf.* 2. tab. 29. fig. 1 — 12.

*Schæff. Icon.* tab. 142. fig. 1. 2.

*Harr. Aur.* tab. 11. fig. 2 — 12.

*Goed. Inf.* 3. tab. 3.

*Reaum. Inf.* 1. tab. 26. fig. 6. 7.

*Roes.*

*Roef. Inf.* 1. pap. 1. tab. 4.

*Wilk. pap.* 56. tab. 3. a. 3.

*Sep. Inf.* 1. 17. tab. 2.

Habitat in Vrtica dioica vulgarissimus.

*Larus* gregaria, spinosa fusco viridique variegata, capite nigra.

*Puppa* dentata, fusca, maculis collaribus aureis variat interdum tota aurea.

406. P. N. P. alis angulatis nigris subtus albo maculatis, *Clytemnestra* anticis vtrinque fascia flaua.

*Papilio Clytemnestra*. *Cram. Inf.* 12. tab. 137. fig. A. B. Habitat Surinami.

Statura P. Dirce. Alae anticae supra nigrae fascia lata flaua, posticae nigrae, immaculae. Subtus fuscae basi albo maculatae, posticis apice subserrugineis.

407. P. N. P. alis angulatis fuscis, fascia anticarum flauescente, subtus nigro undulatis. *Syst. Ent.* 506. 264. *Dirce*.

*Linn. Syst. Nat.* 2. 778. 271. *Mus. Lud. Vlr.* 287.

*Seb. Musf.* 4. tab. 10. fig. 1. 2. 7. 8.

— — — tab. 13. fig. 9. 10.

*Cram. Inf.* 18. tab. 212. fig. C. D.

*Sulz. Hist. Inf.* tab. 17. fig. 11. 12.

Habitat in Indiis.

408. P. N. P. alis dentatis fulvis, punctis margineque albo punctato nigris, subtus griseis. *Progne*. *Gen. Inf. Mant.*

264. \*

*Papilio Progne*. *Cram. Inf.* 1. tab. 5. fig. E. F.

Habitat in America meridionali.

409. P. N. P. alis angulatis fulvis nigro maculatis, posticis *C. album*, subtus C. albo notatis. *Syst. Ent.* 506. 265. *Linn.*

*Syst. Nat.* 2. 778. 168. *Fn. Sv.* 1059. *Geoff. Inf.* 2.

38. 5.

*Papilio vlmariae* similis. *Raj. Inf.* 118. 3.

*Robert. Icon.* tab. 23.

*Merian. Eur.* tab. 14. fig. 1.

*Frisch. Inf.* 4. tab. 4.

*Reaum. Inf.* 1. tab. 27. fig. 9. 10.

*Degeer Inf.* 1. tab. 20. fig. 9. 10.

Harris

94 CLASS. VI. GLOSSATA. *Papilio*.

*Harris Aur.* tab. 1. fig. a — d.

*Roef. Inf.* 1. pap. 1. tab. 5.

*Wilk. pap.* 57. tab. 3. a. c.

*Holl. Inf.* tab. 6. fig. 8.

*Esp. pap.* 1. tab. 13. fig. 3.

*Bergstræss. Inf.* 2. tab. 38. fig. 1 — 5.

*Schæff. Icon.* tab. 147. fig. 3. 4.

*Seb. Mus.* 4. tab. 1. fig. G.

Habitat in Vrtica, Salice, Ribe.

*Larva* solitaria spinosa fulua, dorso antice flavitice albo.

*Puppa* carnea medio coarctata punctis aureis.

Conf. *Papilio Egia. Cram. Inf.* 7. tab. 78. fig.

*C. aureum.* 410. P. N. P. alis dentato caudatis fulvis nigro mac posticis subtus C. aureo notatis. *Syst. Ent.* 506

*Lin. Syst. Nat.* 2. 778. 169.

*Papilio C. aureum. Cram. Inf.* 2. tab. 19. fig. E.

Habitat in Asia.

*Merione.* 411. P. N. P. alis angulatis fulvis, margine fusco, p subtus fuscis maculis strigisque albis coeruleisq

*Papilio Ariadne. Cram. Inf.* 15. tab. 180. fig. B.

Habitat Surinami.

Statura omnino praecedentis. Alae angulatae, fulvae margine postico fusco. Subtus anticae scentes lineolis tribus fulvis transuersis apice macula ocellari strigaeque vndata coeruleis. Posticae medio subferrugineae puncto baseos albo medio lunulae duae albae maculaeque quatuor ruleae subocellares et pone has striga albida ciliaeque vndatae.

*Ariadne.* 412. P. N. P. alis angulatis fulvis, strigis nigris vntis, anticis puncto marginali albo. *Syst. Ent.* 267. *Lin. Syst. Nat.* 2. 778. 170. *Amoen. act.* 407. 71.

*Papilio Merione. Cram. Inf.* 12. tab. 144. fig. G.

Habitat in Iaua.

Conf. *Papilio Coryta. Cram. Inf.* 8. tab. 86. fig.

Vix differt.

413. P. N. P. alis dentato caudatis fulvis nigro maculatis, *Portia*.  
 subtus griseis nigro irroratis. *Syst. Ent.* 507. 268. \*  
*Papilio Peleus* alis caudatis falcatis dentatisque rufis  
 fulvis lineis tribus nigris parallelis, posticarum mar-  
 gine caudaque nigris. *Sulz. Hist. Inf.* tab. 13. fig. 4. ?  
 Habitat in America. v. *Rohr*.  
 Varietas paullo minor, alis anticis minus falcatis fulvis,  
 puncto unico nigro. An mas?
414. P. N. P. alis dentatis nigris, anticis maculis duabus, *Bolina*,  
 posticis solitaria albo coeruleis. *Syst. Ent.* 507. 269.  
*Linn. Syst. Nat.* 2. 781. 188. *Mus. Lud. Vlr.* 295.  
*Papilio Bolina*. *Cram. Inf.* 6. tab. 65. fig. E. F.  
*Papilio Auge*. *Cram. Inf.* 16. tab. 190. fig. A. B.  
*Clerk. Icon.* tab. 21. fig. 2.  
*Drury Inf.* 1. tab. 14. fig. 1. 2.  
*Seb. Mus.* 4. tab. 25. fig. 15. 16.  
 Habitat in Portulaca Indiae orientalis. *D. Koenig*.  
 Conf. *Papilio Lifianassa*. *Cram. Inf.* 18. tab. 205. fig.  
 A. B. Vix distinctus videtur.
415. P. N. P. alis dentatis nigris, anticis margine exteriori albo maculatis, posticis albo luteoque triplici ordine. *Clytia*.  
*Syst. Ent.* 507. 270. *Linn. Syst. Nat.* 2. 781.  
 189. *Mus. Lud. Vlr.* 296.  
 Habitat in India.  
 Corpus nigrum albo maculatum.
416. P. N. P. alis angulato dentatis supra fuscis albo punctatis, fascia rubra, linea vndata nigra. *Amatheia*.  
*Syst. Ent.* 507. 271. *Linn. Syst. Nat.* 2. 779. 174.  
*Papilio Amatheia*. *Cram. Inf.* 18. tab. 209. fig. A. B.  
*Clerk. Icon.* tab. 40. fig. 3.  
 Habitat in Indiis.
417. P. N. P. alis dentatis flavis nigro variis apice radiatis, posticis punctis septem rubris. *Hypsipyle*.  
 265. \*  
*Papilio Hypermnestra*. *Scop. Ent.* 425.  
*Papilio Polyxena*. *Wien. Verz.* tab. tit. praef. p. 162.  
*Hollar. Inf.* tab. 8.  
 Habitat in Europae australioris Aristolochia Clematidis.

*Ramina.* 418. P. N. P. alis dentatis variegatis, supra anticis punctatis sex, posticis quatuor rubris. *Syst. Ent.* 313. 299.  
*Linn. Syst. Nat.* 2. 783. 200.

*Papilio tetrapus*, alis ex coccineo, luteo, argenteo, nigroque variegatis, primoribus rotundioribus. *Osh. Iter.* 51.

*Catesb. Carol.* 2. 95. tab. 95.

Habitat in Lusitania, Europa australi.

*Meleagris.* 419. P. N. P. alis subdentatis albo punctatis, supra fuscis, subtus flauescentibus.

*Papilio Meleagris. Cram. Inf.* 6. tab. 66. fig. A. B.

Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*

Magnitudo et statura P. Leucothoe. Supra alae fuscatae, albo punctatae, subtus flauescentes albo punctatae striga penultima punctis nigris foeta.

*Melantha.* 420. P. N. P. alis dentatis supra fuscis albo punctatis, subtus brunneis. *Syst. Ent.* 313. 297.\*

Habitat in Guinea. *Mus. Dom. Banks.*

*Leucothea.* 421. P. N. P. alis dentatis supra fuscis fasciis tribus albis, subtus luteis, fasciis tribus albis nigro notatis. *Syst. Ent.* 508. 272. *Linn. Syst. Nat.* 2. 780. 179. *Mus. Lud. Vlr.* 292.

*Papilio Erofina. Cram. Inf.* 17. tab. 203. fig. E. F. Vir differt.

*Seb. Mus.* 4. tab. 45. fig. 1. 2.

*Aubent. miscell.* tab. 91. fig. 3. 4.

*Sulz. Hist. Inf.* tab. 18. fig. 2. 3.

Habitat in Asia.

Conf. *Papilio Sulpitia. Cram. Inf.* 18. tab. 214. fig. E. F.

*Medea.* 422. P. N. P. alis dentatis falcatis nigris, fasciis tribus macularibus flavis, angulo ani rubro. *Syst. Ent.* 508. 273.\*

*Papilio Chione. Cram. Inf.* 8. tab. 90. fig. E. F.

Habitat Surinami. *Dom. Drury.*

*Melicerita.* 423. P. N. P. alis dentatis concoloribus atris albo punctatis fasciisque communi alba anticarum interrupta. *Syst. Ent.* 508. 274.\*

*Papilio*

*Papilio Melicerta. Drury Inf. 2. tab. 19. fig. 3. 4.*

*Papilio Heliodore. Cram. Inf. 18. tab. 212. fig. E. F.*

Habitat in Sierra Leon Africae.

424. P. N. P. alis dentatis albis margine flavo, posticis *Dorcas.*  
subtus flavis, fasciis duabus albis. *Syst. Ent. 508.*

275. \*

Habitat in Carolina. *Mus. D. Hunter.*

425. P. N. P. alis dentatis fuscis albo maculatis, subtus ma- *Ophione.*  
culis atris ocellaribus numerosis.

*Papilio Ophione. Cram. Inf. 10. tab. 114. fig. E. F.*

Habitat in Guinea.

Statura P. Melicertae.

426. P. N. P. alis dentatis fulvis nigro fasciatis, posticis *Ilithuia.*  
subtus fasciis albis nigro maculatis.

*Papilio Ilithuia. Drury Inf. 2. tab. 17. fig. 1. 2.*

*Papilio Ilithuia. Cram. Inf. 18. tab. 213. fig. A. B.*

— — — tab. 214. fig. C. D.

Habitat in India orientali. *Dom. Lund.*

Alae subtus fulvae anticis nigro fasciatis lituris albis,  
posticis fasciis duabus baseos albis, quarum anterior  
lineis duabus punctorum nigrorum, posterior vni-  
ca, tunc fascia nigra albo punctata. Margo niger  
striga alba.

427. P. N. P. alis dentatis nigris coeruleo nebulosis, an- *Amphino-*  
ticis fascia vtrinque alba, posticis subtus rubro ra- *me.*  
diatis. *Syst. Ent. 508. 276. Linn. Syst. Nat. 2. 779.*

176.

*Kloman. Inf. 1. tab. 10. fig. 1. 2.*

*Merian. Sur. tab. 8.*

*Aubert. miscell. tab. 92. fig. 7. 8.*

*Cram. Inf. 5. tab. 54. fig. E. F.*

Habitat in America meridionali.

428. P. N. P. alis dentatis fuscis, anticis flavo reticulatis, *Opis.*  
posticis fasciatis.

*Papilio Opis. Drury Inf. 1. tab. 18. fig. 5. 6.*

*Papilio Opis. Cram. Inf. 12. tab. 138. fig. A. B.*

Habitat in Sierra Leon Africae.

*Fabr. Sp. Inf. T. II.*

G

Alae



Alae omnes fuscae. In medio fascia communis flava, anticarum abbreviata. Anticae flavo maculatae apice reticulatae strigaque punctorum alborum, posticae pone fasciam reticulatae. Subtus anticae fusco, coeruleo, nigro flavoque variegatae, posticae apice late flavae arcibus duobus fuscis.

- Crithea.* 439. P. N. P. alis dentatis fuscis maculis ocellaribus, posticis subtus cinereis punctis duobus strigaque fuscis. *Papilio Crithea.* *Drury Inf.* 2. tab. 16. fig. 5. 6. *Papilio Crithea.* *Cram. Inf.* 12. tab. 138. fig. C. D. Habitat in Sierra Leon Africae.

*Praecedente minor.* Alae anticae fuscae maculis plurimis variis ocellaribus atris, iride flava cinctis, et praeterea lineolae aliquot flavae, subtus basi pallidae annulis atris, apice fuscae lunula coerulea. Posticae fuscae strigis flavescentibus margineque interiori luteo, subtus griseae punctis duobus strigaque media fuscis.

- Nerina.* 430. P. N. P. alis dentatis atris, fascia alba, anticarum terminata macula rufa. *Syst. Ent.* 509. 277. Habitat in nova Hollandia. *Mus. Dom. Banks.*

- Venilia.* 431. P. N. P. alis dentatis fuscis, fascia communi arcuata alba marginibus coerulecente. *Syst. Ent.* 509. 278. *Linn. Syst. Nat.* 2. 780. 177. *Mus. Lud. Vir.* 290. *Clerk. Icon.* tab. 32. fig. 4. *Seb. Mus.* 4. tab. 16. fig. 3. 4. 8. 9. Habitat in Indiis.

- Allmena.* 432. P. N. P. alis dentatis atris, fascia coerulea interrupta, singulis punctis septem albis marginalibus. *Syst. Ent.* 509. 279. *Linn. Syst. Nat.* 2. 780. 178. *Mus. Lud. Vir.* 291. *Seb. Mus.* 4. tab. 11. fig. 13. 14. Habitat in Indiis.

- Phaerusa.* 434. P. N. P. alis dentatis fulvis, fasciis tribus fuscis, anticarum longitudinalibus, posticarum transuersis. *Syst. Ent.* 509. 280. *Linn. Syst. Nat.* 2. 780. 180. *Mus. Lud. Vir.* 293.

CLASS. VI. GLOSSATA. Papilio. 99

*Papilio Phaerusa*. Cram. Inf. 11. tab. 130. fig. B. C.

Roef. Inf. 4. tab. 2. fig. 1.

Habitat in Indiis.

434. P. N. P. alis dentato caudatis, flavo nigroque variis, *Hippona*.  
posticis punctis quatuor albis. Gen. Inf. Mant. 265.\*

*Papilio Fabius*. Cram. Inf. 8. tab. 90. fig. C. D.

Habitat in India. D. Schulz.

435. P. N. P. alis dentatis fulvis, anticis strigis duabus, *Julia*.  
posticis margine postico nigris. Syst. Ent. 509. 281.\*

*Papilio Aliconea*. Cram. Inf. 18. tab. 215. fig. A. F. G.

Habitat in America. Dom. v. Rohr.

436. P. N. P. alis angulato dentatis fuscis, fascia commu- *Laura*.  
ni dimidiato fulva, posticis subtus argenteis, fascia  
niuea. Syst. Ent. 510. 282.

*Papilio Laure*. Drury Inf. 2. tab. 17. fig. 5. 6.

Habitat in Jamaica.

437. P. N. P. alis dentatis flavis supra margine nigro, sub- *Lydia*.  
tus striga punctorum nigrorum.

*Papilio Lydia*. Cram. Inf. 13. tab. 148. fig. C. D.

Habitat Surinami.

Alae anticae supra flavae basi obscuriores apice nigrae.

Subtus basi flavae striga fulva apice obscuriores striga

e punctis sex nigris. Margo striga nigra. Posticae

obscurae fascia flava maculisque quinque ferrugineis

lunulaeque nigra, apice nigrae strigis duabus pallidio-

ribus. Subtus basi flavae striga communi fulva, in

medio luteae fascia fulva et in hac striga e lunulis

cinereis punctisque nigris, apice striga fulva, cine-

rea flavaeque.

438. P. N. P. alis dentatis supra fuscis, fascia communi *Iphicla*.  
alba maculaque ferruginea anguli ani. Syst. Ent. 510.

283. Linn. Syst. Nat. 2. 780. 181. Mus. Lud. Vlr. 311.

*Papilio capucinus* alis fuscis, in primoribus flava in  
posticis alba linea. Naturf. 6. tab. 6. fig. 2. a. b.

*Papilio Iphicla*. Cram. Inf. 16. tab. 188. fig. D. E. F.

Drury Inf. 1. tab. 14. fig. 3. 4.

Clerk. Inf. tab. 41. fig. 3.

*Seb. Musf.* 4. tab. 13. fig. 5. 6.

— — — tab. 30. fig. 15. 16.

Habitat in America meridionali.

*Delila.* 439. P. N. P. alis dentatis oblongis fulvis, anticis lunula nigra. *Syst. Ent.* 510. 284. \*

*Papilio Cellene. Cram. Inf.* 18. tab. 215. fig. D. E.

Habitat in America. D. *Eaton.*

*Dido.* 440. P. N. P. alis dentatis oblongis nigris, viridi maculatis, posticis fascia maculisque septem transuersis coerulescentibus. *Syst. Ent.* 510. 285. *Linn. Syst. Nat.* 2. 782. 192. *Amoen. acad.* 6. 408. 74.

*Papilio Dido. Cram. Inf.* 17. tab. 96. fig. E. F.

*Seb. Musf.* 4. tab. 4. fig. 13. 14.

*Merian. Sur.* tab. 2.

Habitat in Americae Ananas.

Heliconiis affinis ob alas oblongas, at dentatas.

*Larua* spinosa viridis linea rubra albaque, spinis duobus caudalibus elongatis.

*Neaerea.* 441. P. N. P. alis dentatis, supra fuscis, albo maculatis, subtus maculis albis, linea rubra cinctis. *Syst. Ent.* 510. 286. *Linn. Syst. Nat.* 2. 782. 190. *Musf. Lud. Vlr.* 297.

*Papilio Neaerea. Cram. Inf.* 7. tab. 75. fig. C. D.

*Edw. Av.* 33. tab. 33.

*Seb. Musf.* 4. tab. 19. fig. 1. 2.

β. *Papilio Tipha* alis subcaudatis nigris, primoribus maculis duabus, posticis disco rubro marginato. *Linn. Syst. Nat.* 2. 776. 164. *Musf. Lud. Vlr.* 308.

*Papilio Tipha. Cram. Inf.* 1. tab. 8. fig. D. E.

*Clerk. Icon.* tab. 32. fig. 3.

Habitat in India.

Angulus ani ruber.

*Aceste.* 442. P. N. P. alis subdentatis, anticis nigris basi fascisque flauis, posticis flauis, subtus fasciis fuscis. *Syst. Ent.* 511. 287. *Linn. Syst. Nat.* 2. 782. 191. *Musf. Lud. Vlr.* 298.

*Papilio Aceste. Cram. Inf.* 11. tab. 121. fig. E. F.

*Seb.*

CLASS. VI. GLOSSATA. Papilio. 101

*Seb. Mus.* 4. tab. 27. fig. 1. 2.

— — — tab. 30. fig. 7. 8.

Habitat in India.

Alae posticae fulvae margine punctato subtus nigro fasciatae.

443. P. N. P. alis dentatis nigris, albo maculatis, posticis *Eurino-*  
basi albis. *me.*

*Papilio Eurinome. Cram. Inf.* 6. tab. 70. fig. A.

Habitat in India orientali.

Statura omnino P. dissimilis.

444. P. N. P. alis dentatis dilatato venosis nigris, maculis albis sagittatis, posticis subtus maculis marginalibus luteis. *Syst. Ent.* 511. 288. *Linn. Syst. Nat.* 2. 782.  
195. *Mus. Lud. Vlr.* 301.

*Papilio dissimilis. Cram. Inf.* 7. tab. 82. fig. C. D.

*Clerk. Icon.* tab. 16. fig. 3.

*Ehret. pict.* tab. 17.

*Sulz. Hist. Inf.* tab. 18. fig. 6.

Habitat in Asia.

445. P. N. P. alis subrepandis concoloribus nigris, punctis coerulefcenti albidis, posticis punctis coccineis. *affimilis.*  
*Syst. Ent.* 511. 289. *Linn. Syst. Nat.* 2. 782. 194.  
*Mus. Lud. Vlr.* 300.

*Papilio affimilis. Cram. Inf.* 13. tab. 154. fig. A.

*Papilio finilis. Cram. Inf.* 1. tab. 9. fig. B. C. ?

*Clerk. Icon.* tab. 16. fig. 1.

*Drury Inf.* 1. tab. 17. fig. 3. 4.

Habitat in China.

446. P. N. P. alis subrepandis concoloribus, punctis coerulefcenti albis versus basin lineatis. *ffimilis.*  
290. *Linn. Syst. Nat.* 2. 782. 193. *Mus. Lud. Vlr.* 299.

*Papilio Melaneus. Cram. Inf.* 3. tab. 30. fig. D.

*Papilio Limniace. Cram. Inf.* 5. tab. 59. fig. D. E.

*Papilio Aventina. Cram. Inf.* 5. tab. 59. fig. F.

*Clerk. Icon.* tab. 16. fig. 2.

*Petiv. Gazoph.* tab. 9. fig. 13.

Habitat in Asia.

- affinis.* 447. P. N. P. alis dentatis nigris, albo maculatis, posticis subtus limbo nigro flavo alboque maculato. *Syst. Ent.* 511. 291. \*

Habitat in nova Hollandia. *Mus. Dom. Banks.*

- Lais.* 448. P. N. P. alis dentatis nigris, coerulefcenti striatis, subtus fuscis.

*Papilio Lais.* *Cram. Inf.* 10. tab. 110. fig. A. B.

*Seb. Mus.* 4. tab. 7. fig. 23. 24.

Habitat in India orientali.

Statura omnino praecedendum, at paullo minor. Distinctus alis subtus fuscis.

- Egialea.* 449. P. N. P. alis dentatis, anticis atris maculis, posticis fuscis disco albis.

*Papilio Egialea.* *Cram. Inf.* 16. tab. 192. fig. D.

Habitat in Asia, Africa.

Statura P. Semele. Corpus fuscum thorace albo punctato.

- Panope.* 450. P. N. P. alis dentatis fuscis concoloribus, margine exteriori albo maculato, angulo ani fulvo. *Syst. Ent.* 512. 292.

*Papilio Panope* alis dentatis fuscis concoloribus, limbo exteriori albo maculatis, posticis margine luteo maculatis. *Linn. Syst. Nat.* 2. 782. 196.

*Papilio Perimela.* *Cram. Inf.* 6. tab. 65. fig. C. D.

*Seb. Mus.* 4. tab. 41. fig. 21. 22.

Habitat in Asia.

Conf. *Cram. Inf.* 6. tab. 67. fig. A. B. C. D. E. huius videtur varietas, at puncta alarum minora.

- Sibilla.* 451. P. N. P. alis dentatis supra nigris subtus ferrugineis, fascia vtrinque maculari alba. *Syst. Ent.* 512. 294.

*Linn. Syst. Nat.* 2. 781. 186. *Mus. Lud. Vlr.* 303.

*Papilio* alis dentatis nigris subtus fusco rubris, vtrinque maculis albis fasciatim positis. *Geoff. Inf.* 2. 73. 45.

*Papilio* mediae magnitudinis elegantissima, alis supinis nigris. *Raj. Inf.* 126. 3.

*Papilio Sibilla.* *Drury Inf.* 2. tab. 16. fig. 1. 2.

*Papilio Sibilla.* *Cram. Inf.* 10. tab. 114. fig. C. D.

*Papilio rinularis.* *Scop. carn.* 443.

*Roef. Inf.* 3. tab. 70. fig. 1. 2. 3.

*Fuesl. Magaz.* 256. tab. 2. fig. 4 - 7.

**CLASS. VI. GLOSSATA. Papilio.** 103

*Esp. pap.* 1. tab. 38. suppl. 14. fig. 2.

*Schaeff. Icon.* tab. 152. fig. 1. 2.

β. *Papilio Camilla* alis dentatis fuscis subconcoloribus albo fasciatis maculatisque, angulo ani rubro. *Linn. Syst. Nat.* 2. 781. 187.

*Roef. Inf.* 3. tab. 33. fig. 3.

*Petiv. Gazoph.* 72. tab. 12. fig. 10.

Habitat in Xylosteo, Caprifolio.

*Larva* viridis, capite spinisque rubris.

*Puppa* viridis aureo maculata, antice bicornis.

Conf. *Papilio Lucretia*. *Cram. Inf.* 4. tab. 45. fig. C. D.

452. P. N. P. alis dentatis fulvis, nigro maculatis, fascia *Procris*. alba, subtus variegatis.

*Papilio Procris*. *Cram. Inf.* 9. tab. 106. fig. E. F.

*Edw. Av.* tab. 299.

Habitat in China.

Magnitudo et statura omnino praecedentis. Alae supra fulvae, nigro maculatae. In medio fascia macularis alba, quae ad marginem crassiore macula magna atra terminatur. Margo posticus strigis duabus albis. Subtus basi virides, tunc albo fulvo nigroque variae.

453. P. N. P. alis dentatis atris, flavo maculatis, posticis *Micalia*. medio fulvo nigro maculato.

*Papilio Micalia*. *Cram. Inf.* 9. tab. 108. fig. C. D.

Habitat in America boreali.

Medius. Alae falcatae, anticae atrae linea bascos, macula magna media punctisque marginalibus flavis. Posticae atrae, in medio late fulvae, punctis quinque atris. Margo posticus ater punctis flavis. Subtus variegatae.

454. P. N. P. alis dentatis fuscis, anticis albo maculatis, posticis immaculatis. *Maia*.

*Papilio Maia* alis dentatis fuscis, anticis apice albo maculatis, posticis subtus cinereis, punctis quatuor virescentibus. *Syst. Ent.* 512. 295.

*Papilio Agathina*. *Cram. Inf.* 14. tab. 167. fig. E. F.

Habitat in Brasilia. *Mus. Dom. Banks.*

Femina alis supra coeruleo nitentibus maculisque albis obsoletis.

Conf. *Papilio Orphis*. *Cram. Inf.* 4. tab. 42. fig. E. F.  
huic nimis affinis, at alae posticae magis flavescentes  
punctaque approximata ocellata.

*Carinenta*. 455. P. N. P. alis falcato dentatis fuscis, flauo maculatis,  
anticis apice atris maculis quatuor albis.

*Papilio Carinenta*. *Cram. Inf.* 9. tab. 108. fig. E. F.  
Habitat Surinami.

Statura praecedentis. Corpus fuscum. Palpi porrecti  
acuminati. Alae anticae falcato dentatae, basi fuscae,  
apice atrae maculis quatuor albis. In medio maculae  
duae oblongae flavae. Subtus basi fulvae, medio fu-  
scae, albo maculatae; apice coerulescentes, nigro irro-  
ratae. Posticae supra fuscae macula magna flaua,  
subtus coerulescentes, nigro irroratae.

*Harpyia*. 456. P. N. P. alis falcato dentatis fuscis, fascia communi  
coerulescente nigro punctata, subtus brunneis, striga  
fusca.

Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*

Nimis affinis sequenti, at paullo maior. Alae anticae  
valde falcatae fuscae maculis aliquot ferrugineis ad  
basin marginis crassioris. Pone medium fascia lata  
coerulescens, nigro punctata, posticeque fuluo margi-  
nata. Versus apicem alae maculae aliquot albae. Sub-  
tus brunneae striga abbreviata fusca punctisque apli-  
cis albis. Posticae dentatae fuscae fascia lata coe-  
rulescente nigro punctata et postice fascia ferruginea  
terminata, subtus brunneae striga fusca punctisque  
aliquot subocellaribus minutissimis albis.

Alium vidi simillimum vix distinctum, at alis obs-  
curioribus fascia incarnata.

*Pelarga*. 457. P. N. P. alis dentatis nigris, fascia communi flaua,  
nigro punctata. *Syst. Ent.* 513. 296. \*

Habitat in Brasilia. *Mus. Dom. Banks.*

*Melissa*. 458. P. N. P. alis dentatis fuscis, posticis subtus griseis  
marmoratis. *Syst. Ent.* 513. 298. \*

Habitat in Insula Terre neuve Americae. *Mus. Dom. Banks.*

Conf. *Papilio Acanthea*. *Cram. Inf.* 12. tab. 134. fig. D. E.  
F. G.

CLASS. VI. GLOSSATA. Papilio. 105

9. P. N. P. alis dentatis albis, limbo fuluo, posticis *Herpilia*.  
subtus fuluis, fasciis duabus punctisque medio albis.

*Gen. Inf. Mant.* 265.\*

*Papilio Mardania. Cram. Inf.* 18. tab. 213. fig. F. G.

Habitat in America. *D. Schultz.*

10. P. N. P. alis dentatis fuscis, anticis fascia fulua, po- *Elsa.*  
sticis alba. *Syst. Ent.* 513. 300. *Linn. Syst. Nat.* 2.

781. 183. *Mus. Lud. Vlr.* 312.

Habitat in Indiis.

11. P. N. P. alis dentatis fuscis, disco communi cinereo *Amphice-*  
fusco vndato, subtus apice griseis, lunulis nigris. *da.*

*Papilio Amphiceda. Cram. Inf.* 13. tab. 146. fig. D. E.

Habitat in Guinea.

*Paruus.* Alae omnes supra basi margineque omni fu-  
scae, in medio cinercae strigis duabus valde vndatis,  
antere fufca, pofteriore atra, fubtus bafi palli-  
dae maculis rufis lineolisque vndatis nigris, apice  
grifeae ftriga lunularum nigrarum.

12. P. N. P. alis dentatis fuscis, anticis fupra fascia ful- *Mefente-*  
ua, fubtus maculis centralibus coeruleis. *ria.*

*Papilio Mefenteria. Cram. Inf.* 14. tab. 162. fig. B. C.

Habitat Surinami.

Alae omnes fupra obfcurae ftrigis obfcurioribus. Anti-  
cae fascia lata fulua macula transuerfa nigra. Sub-  
tus anticae bafi flauefcentes, in medio maculis fex  
coeruleis annulo atro cinctis, tunc rufae, apice fu-  
scae ftrigis obfcurioribus. Pofticae fupra immacula-  
tae fubtus bafi flauefcentes lineolis tribus transuerfis  
atris et pone has fufcae, ftriatae.

13. P. N. P. alis dentatis luteis, anticis apice nigris fa- *Niphe.*  
fcia albida, pofticis fubtus argentatis, ocellis quin-  
que. *Syst. Ent.* 514. 302. *Linn. Syst. Nat.* 2. 785.  
208.

*Papilio Niphe. Drury Inf.* 1. tab. 6. fig. 1.

*Papilio Niphe. Cram. Inf.* 2. tab. 14. fig. B. C.

*Papilio Hyperbius. Linn. Amoen. acad.* 6. 408. 75.

*Aub. mifcell.* tab. 93. fig. 1. 2.

Habitat in China.



106 CLASS. VI. GLOSSATA. Papilio.

Conf. *Drury Inf.* 1. tab. 6. fig. 2.

*Cram. Inf.* 2. tab. 14. fig. D. E.

*Lucina.* 464. P. N. P. alis dentatis fuscis testaceo maculatis, subtus fasciis duabus macularum albidarum. *Syst. Nat.* 514. 303. *Linn. Syst. Nat.* 2. 784. 203. *Fn. Sv.* 106.

*Papilio fritillarius minor. Raj. Inf.* 122. 12.

*Petiv. Gazoph.* tab. 16. fig. 10.

*Esp. pap.* 1. tab. 16. fig. 1.

*Schaeff. Icon.* tab. 172. fig. 12.

*Seb. Musf.* 4. tab. 3. B. 1-4.

Habitat in Europae graminibus.

*Cinxia.* 465. P. N. P. alis dentatis fulvis nigro maculatis, postice subtus fasciis tribus albidis nigro maculatis. *Syst. Nat.* 514. 304. *Linn. Syst. Nat.* 2. 784. 205. *Fn. Sv.* 1063. *Geoff. Inf.* 2. 45. 12.

*Papilio fritillarius fasciis subtus pallidis. Raj. Inf.* 121. 9.

*Petiv. Gazoph.* tab. 18. fig. 10.

*Wilk. pap.* 58. tab. 3. a. 8.

*Roef. Inf.* 4. tab. 13.

*Reaum. Inf.* 2. tab. 9.

*Schaeff. Elem.* tab. 1. fig. 9.

*Esp. papil.* 1. tab. 16. fig. 3.

— — — tab. 25. Suppl. 1. fig. 2.

— — — tab. 46. Suppl. 22. fig. 2.

*Schaeff. Icon.* tab. 217. fig. 3. 4.

*Degeer Inf.* 2. tab. 1. fig. 17. 18.

Habitat in Hieracio Pilosella.

*Larva* spinosa, atra, linea laterali e punctis albis.

*Puppa* fusca, subtus punctis fulvis lineata.

Plures huius vidi, an varietates aut species, determinabit metamorphosis.

*Hylla.* 466. P. N. P. alis dentatis fulvis, nigro punctatis, postice basi supra fuscis subtus albidis.

*Papilio Hyllus. Cram. Inf.* 4. tab. 43. fig. B. C.

Habitat in Oriente.

Maior P. *Cinxia*. Alae anticae vtrinque fulvae, nigro punctatae, supra margine fusco. Posticae nigro punctatae.

Gane

**CLASS. VI. GLOSSATA. Papilio.** 187

Ætatac, supra basi fuscae, apice fulvae, subtus basi albidæ, apice fulvae.

1. P. N. P. alis dentatis fulvis, nigro maculatis, posticis fasciis subtus fasciis flavis, strigis vndatis nigris. *Linn.*

*Syst. Nat.* 2. 784. 204. *Enl. Sv.* 1062.

*Petiv. Gazoph.* tab. 1. fig. 8.

*Raj. Inf.* 120. 8.

*Wilk. pap.* 58. tab. 2. a. 9.

habitat in Corylo, Erica, Scabiosa.

simis affinis P. Cinxiae, at paullo maior.

2. P. N. P. alis fulvis, nigro maculatis, posticis fasciis tribus virecentibus, fimbria punctata. *Syst. Ent.* 514. 304. *Dia.*

3. *apilio* *Dia* alis dentatis fuscis, testaceo maculatis, subtus maculis quatuor, ordinibusque punctorum duobus argenteis. *Linn. Syst. Nat.* 2. 785. 207.

4. *apilio* fritillaria fimbria punctata. *Raj. Inf.* 121. 11.

*Roes. Inf.* 4. tab. 13. fig. 1. 2. 3. 6. 7.

*Esp. Pap.* 1. tab. 16. fig. 4.

*Kleman. Inf.* 1. tab. 40.

habitat in Plantagine.

similimus P. Cinxiae, attamen distinctus.

5. *arua* grisea spinarum albarum ferruginearumque ordinibus alternis.

6. *uppa* flavescenti nigro variegata.

7. P. N. P. alis dentatis fuscis, rufo maculatis, apice fulvis, subtus maculis rufis albo cinctis. *Dryope.*

8. *apilio* *Dryope.* *Cram. Inf.* 7. tab. 78. fig. E. F.

habitat in Sierra Leon Africae.

9. *arvus.* Alae omnes fuscae, rufo maculatae, apice fulvae, margine fusco, striga posticarum fulva, subtus basi virecentes maculis rufis, linea alba cinctis, apice pallidiores striga vndata alba.

10. P. N. P. alis denticulatis variegatis subtus reticulatis, anticis supra maculis aliquot albis. *Syst. Ent.* 515. 306. *Levand.*

*Linn. Syst. Nat.* 2. 783. 201.

*Roes. Inf.* 1. pap. 1. tab. 9.

*Esp. pap.* 1. tab. 15. fig. 2.

*Bergstr.*

108 CLASS. VI. GLOSSATA. Papilio.

*Bergstr. Inf.* 1. tab. 35. fig. 9. | 10.

*Schaeff. Icon.* tab. 132. fig. 1. 2.

Habitat in Vrtica dioica.

*Larua* gregaria, spinosa, atra, pedibus rufis capite spinoso.

*Puppa* dentata, fusca, thorace subtus flavescente.

*Prorsa.* 471. P. N. P. alis dentatis fuscis, fascia vtrinque alba antice interrupta, subtus albo reticulatis. *Syst. Ent.* 515. 307. *Linn. Syst. Nat.* 2. 783. 202.

*Roef. Inf.* 1. pap. 1. tab. 8.

*Merian. Europ.* 88. fig. 1.

*Esp. pap.* 1. tab. 15. fig. 3.

*Bergstr. Inf.* 2. tab. 35. fig. 5. 6.

Habitat in Vrtica dioica.

*Larua* gregaria spinosa fusco fulvoque variegata, capite bispinoso.

*Puppa* dentata supra fusca, subtus pallescens.

*Paphia.* 472. P. N. P. alis dentatis fulvis, nigro maculatis, sub fasciis argenteis. *Syst. Ent.* 515. 308. *Linn. Syst. Nat.* 2. 785. 209. *Fn. Sv.* 1064. *Geoff. Inf.* 2. 42. 8.

*Papilio* maior, alis fulvis, maculis nigris supina parte pronata transuersis argenteis depicta. *Raj. Inf.* 119.

*Roef. Inf.* 1. pap. 1. tab. 7.

*Wilk. pap.* 57. tab. 2. a. 7.

*Schaeff. Elem.* tab. 94. fig. 2.

— — — *Icon.* tab. 97. fig. 3. 4.

*Esp. pap.* 1. tab. 17. fig. 1. 2.

*Bergstr. Inf.* 2. tab. 41. fig. 1-5.

*Petiv. Mus.* 321.

Habitat in Viola canina.

*Larua* solitaria, spinosa, fulva, linea dorsali fuscis spinis collaribus longissimis.

*Puppa* grisea, collo coarctato, dentibus sex punctis que quatuor anticis aureis.

*Cynara.* 473. P. N. P. alis dentatis fuscis atro flavoque variegatis subtus viridibus, maculis tribus fasciisque duabus argenteis. *Gen. Inf. Mant.* 265.\*

*Papilio Maja.* *Cram. Inf.* 3. tab. 25. fig. B. C.

Habitat in Oriente.

CLASS. VI. GLOSSATA. Papilio. 109

4. P. N. P. alis crenatis flavis, fascia communi argentea, *Cytheria*.  
 subtus fascia lanceolata argentea. *Syst. Ent.* 516. 309.  
*Linn. Syst. Nat.* 2. 785. 210. *Mus. Lud. Vlr.* 305.  
*Clerk. Icon.* tab. 39. fig. 3.  
*Seb. Mus.* 4. tab. 13. fig. 11. 12.  
 — — — tab. 31. fig. 5. 6.

Habitat in Indiis.

75. P. N. P. alis dentatis fulvis, nigro maculatis, subtus *Aglaja*.  
 maculis 21 argenteis. *Syst. Ent.* 516. 310. *Linn. Syst.*  
*Nat.* 2. 785. 211. *Fn. Sv.* 1065. *Geoff. Inf.* 2. 42. 9.  
*Papilio* maior alis fulvis, supina parte maculis nigris  
 crebris, prona etiam argenteis depictis. *Raj. Inf.*  
 119. 5.

*Papilio* fritillarius maior maculis subtus argenteis.

*Petiv. Mus.* 320.

*Wlk. pap.* 59. tab. 2. a. 12.

*Schaeff. Icon.* tab. 7. fig. 1. 2.

*Roef. Inf.* 4. tab. 25.

*Esp. papil.* 1. tab. 17. fig. 3.

Habitat in Europae Viola.

*Larva* solitaria spinosa, atra maculis quadratis latera-  
 libus ferrugineis.

*Puppa* fusca.

76. P. N. P. alis dentatis fulvis, nigro maculatis, subtus *Myrina*.  
 maculis 30 argenteis.

*Papilio Myrina. Cram. Inf.* 16. tab. 189. fig. B. C.

Habitat in America boreali.

Parvus. Maculae 9 argenteae in alis anticis et 21 in  
 posticis, et praeter has punctum centrale strigaeque  
 posticae et punctis atris ocellatis.

77. P. N. P. alis dentatis fulvis nigro maculatis, subtus *Cybele*.  
 maculis 34 argenteis. *Syst. Ent.* 516. 311.\*

*Papilio Daphnis. Cram. Inf.* 5. tab. 57. fig. E. F.

Habitat in America.

Fascia flava alae posticae ante marginem.

78. P. N. P. alis dentatis fulvis, nigro maculatis, subtus *Idalia*.  
 maculis 37 argenteis, posticis supra coeruleis, albo  
 punctatis. *Syst. Ent.* 516. 312.\*

*Papilio*

110. CLASS. VI. GLOSSATA. Papilio.

*Papilio Idalia. Drury Inf.* 1. tab. 13. fig. 1. 2. 3.

*Papilio Idalia. Cram. Inf.* 4. tab. 44. fig. D. E. F. G.

Habitat in America boreali.

Sexus alter margine alarum anticarum, albo maculato et puncta posticarum omnia flava.

*Diana.* 479. P. N. P. alis dentatis fulvis, disco communi nigro, posticis subtus maculis novem argenteis.

*Papilio Diana. Cram. Inf.* 2. tab. 98. fig. D. E.

Habitat in Virginia.

Magnus. Alae anticae basi nigrae, apice fulvae striga duabus punctorum nigrorum, subtus disco maculato. Posticae basi nigrae, apice fulvae striga una punctorum nigrorum. Subtus basi fuscae maculae duabus argenteis, apice flavescentes lunulis septem argenteis.

*Adippe.* 480. P. N. P. alis dentatis fulvis, nigro maculatis, subtus maculis 23 argenteis. *Syst. Ent.* 517. 313. *Linn. Syst. Nat.* 2. 786. 212.

*Degeer Inf.* 2. tab. 1. fig. 8. 9.

*Schaeff. Icon.* tab. 61.

*Bergstraeff. Inf.* 2. tab. 42. fig. 5. 6.

*Esp. pap.* 1. tab. 18. fig. 1.

— — — tab. 26. Suppl. 11. fig. 4.

— — — tab. 43. suppl. 19. fig. 2.

Habitat in Europae pratis.

*Lathonia.* 481. P. N. P. alis dentatis luteis nigro maculatis, subtus maculis 37 argenteis. *Syst. Ent.* 517. 314. *Linn. Syst. Nat.* 2. 786. 213. *Fn. Sv.* 1068. *Geoff. Inf.* 2. 120. 6.

*Papilio Rigenis minor aureus, maculis argenteis subtus perbelle notatus. Raj. Inf.* 120. 6.

*Hufn. pict.* tab. 11. fig. 11.

*Robert. Icon.* tab. 12.

*Merian. Europ.* 2. tab. 157.

*Roef. Inf.* 3. tab. 10.

*Esp. pap.* 1. tab. 18. fig. 2.

*Schaeff. Icon.* tab. 143. fig. 1. 2.

*Seb. Mus.* 4. tab. 1. H. 1 — 4.

Habitat in Europa.

*Larus*

CLASS. VI. GLOSSATA. Papilio. III.

*Larva* spinosa fusca nigro maculata lineaque dorsali alba.

*Puppa* crenata, antice fusca, postice virescens.

3. P. N. P. alis dentatis fulvis, nigro maculatis, subtus *Euphro-*  
maculis 9 argenteis. *Syst. Ent.* 517. 315. *Linn. Syst.* *lyne.*  
*Nat.* 2. 786. 214. *Fn. Sv.* 1069. *Geoff. Inf.* 2. 44. 11.

*Papilio* fritillarius maior alis fulvis, supernis maculis nigris tessellatis. *Raj. Inf.* 120. 7.

*Petiv. Mus.* 35. 382.

*Esp. Papil.* 1. tab. 18. fig. 3.

*Bergstr. Inf.* 2. tab. 42. fig. 1. 2.

*Degeer Inf.* 2. tab. 1. fig. 10. 11.

Habitat in Europae Viola montana.

3. P. N. P. alis dentatis fulvis, nigro maculatis, subtus *Niobe.*  
maculis pallidis punctisque quatuor argenteis ocella-  
tis. *Syst. Ent.* 517. 316. *Linn. Syst. Nat.* 2. 786. 215.  
*Fn. Sv.* 1067.

*Esp. pap.* 1. tab. 18. fig. 4.

-*Schaeff. Icon.* tab. 208. fig. 1. 2.

Habitat in Europa.

4. P. N. P. alis dentatis fulvis, nigro maculatis, post- *Bellona.*  
cis subtus apice argenteis, maculis ocellaribus fulvis  
sex. *Syst. Ent.* 517. 317.\*

Habitat in America boreali. *Dom. Ammiral.*

Conf. *Papilio Hegisia. Cram. Inf.* 18. tab. 209. fig. E.

F. vix tamen idem.

5. P. N. P. alis dentatis fulvis nigro maculis, post- *Phalan-*  
cis subtus apice glaucis, punctis ocellaribus fulvis *ta.*  
tribus. *Syst. Ent.* 518. 318.

*Papilio Phalantha. Drury Inf.* 1. tab. 21. fig. 1. 2.

Habitat in India orientali.

16. P. N. P. alis dentatis fulvis nigro maculatis, subtus *Vanillae.*  
maculatis 30 argenteis. *Syst. Ent.* 518. 319. *Linn.*  
*Syst. Nat.* 2. 787. 216. *Mus. Lud. Vlr.* 306.

*Clerk. Icon.* tab. 40. fig. 2.

*Aubert. miscell.* tab. 70. fig. 2. 3.

*Merian. Sur.* 25. tab. 26.

*Sloas.*

*Sloan. Iam.* 2. tab. 239. fig. 23. 24.

*Sulz. Hist. Inf.* tab. 18. fig. 4. 5.

*Cram. Inf.* 18. tab. 202. fig. A. B.

Habitat in Epidendro Vanilla Americae.

*Iuno.* 437. P. N. P. alis angulato dentatis oblongis fulvis limbo atro, posticis subtus atris, maculis fulvis argenteisque.

*Papilio Iuno. Cram. Inf.* 18. tab. 215. fig. B. C.

Habitat Surinami.

Alae magis oblongae, magis angulatae, quam in reliquis. Apex alae anticae ater, supra maculis fulvis, subtus flavis argenteisque.

### † Plebeji Rurales.

*Cupido.* 488. P. P. R. alis posticis sex dentato caudatis, subtus albis maculis argenteis. *Syst. Ent.* 318. 320. *Linna. Syst. Nat.* 2. 787. 217. *Mus. Lud. Vlr.* 313.

*Petiv. Gazoph.* tab. 10. fig. 9.

*Roef. Inf.* 4. tab. 3. fig. 7.

*Merian. Sur.* tab. 10.

*Aubert. miscell.* tab. 18. fig. 4. 5.

*Seb. Mus.* 4. tab. 33. fig. 5—8.

*Cram. Inf.* 14. tab. 164. fig. D. E. F. G.

Habitat in Americis Gossypio.

*Larva* alba nigro punctata.

*Helius.* 489. P. P. R. alis tricaudatis, anticis nigris, posticis coeruleis, subtus cinereis, strigis duabus brunneis occurrentibus, limbo duobus anguli ani.

Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*

Medius. Antennae nigrae, albo annulatae, apice fulvae.

Alae anticae nigrae margine tenuiori coeruleo, posticae coeruleae nitidae caudis tribus atris apice albis. Subtus omnes cinerae strigis tribus fuscis occurrentibus, limbo duobus atris iride rufa ad angulum ani.

*Ianias.* 490. P. P. R. alis tricaudatis coeruleis, limbo atro, subtus viridibus.

*Papilio Ianias. Cram. Inf.* 18. tab. 213. fig. D. E.

Habitat Surinami.

Macula atra subtus in alis posticis versus marginem.

191. P. P. R. alis tricaudatis fuscis, subtus disco variegato *Amor*.  
 strigaeque postica aurea. *Syst. Ent.* 518. 321.\*  
 Habitat in India orientali. *Mus. Tottianum*.
192. P. P. R. alis tricaudatis fuscis, subtus niucis, nigro *Montus*.  
 subfasciatis, posticis ocellis duobus auratis.  
 Habitat in Africa aequinoctiali. *Mus. Dom. Banks*.  
 Paruus. Alae omnes supra fuscae immaculatae, margi-  
 ne tamen posticarum albo strigoso. Caudae tres bre-  
 vissimae albae. Subtus albiissimae, anticis macula  
 transversali media, fascia apiceque nigris, posticis  
 punctis tribus bascos, fascia media interrupta, litu-  
 raque apicis nigris. Ocelli duo nigro inaurati iride  
 fulua ad marginem posticum.
193. P. P. R. alis tricaudatis fuscis, subtus albis strigis dua- *Faunus*.  
 bus fuluis. *Gen. Inf. Mant.* 266.\*  
*Papilio Faunus. Drury Inf.* 2. tab. 1. fig. 4. 5.  
*Cram. Inf.* 4. tab. 39. fig. B. C.  
 — — — 8. tab. 96. fig. F. G.  
 Habitat in Sierra Leon Africae.
194. P. P. R. alis tricaudatis coerulescentibus, apice nigris, *Aetæon*.  
 subtus aureis, atomis nigris. *Syst. Ent. App.* 829.\*  
 Habitat in — — — *Dom. Drury*.
195. P. P. R. alis tricaudatis fuscis, subtus albo maculatis, *Hymen*.  
 posticis supra albis. *Syst. Ent.* 519. 322.\*  
 Habitat in Sierra Leon Africae. *Mus. Dom. Banks*.
196. P. P. R. alis subtricaudatis coeruleis nitidissimis, po- *Gabriela*.  
 sticis subtus fuscis, ante apicem albo fasciatis.  
*Papilio Gabriela. Cram. Inf.* 1. tab. 6. fig. C. D.  
 Habitat in America meridionali.  
 Alae supra coeruleae, nitidissimae margine tenuissime  
 atro. Subtus posticae fuscae, apice ferrugineae, albo  
 fasciatae, angulo ani virefcenti.
197. P. P. R. alis subtricaudatis coeruleis, auro punctatis, *Pelion*.  
 limbo fusco. *Gen. Inf. Mant.* 267.\*  
*Papilio Pelion. Cram. Inf.* 1. tab. 6. fig. E. F.  
 Habitat in America meridionali.
198. P.



- Aunus.* 498. P. P. R. alis subtricaudatis coeruleis, limbo atro, fuscus nigris, albo fasciatis. *Gen. Inf. Mant.* 267.\*  
*Papilio Aunus. Cram. Inf.* 2. tab. 23. fig. E. F.  
Habitat Coracao Americae.
- Vulcanus.* 499. P. P. R. alis bicaudatis, supra fuscis, fulvo maculatus subtus strigis flauis fuluis argenteisque variegatus. *Syst. Ent.* 519. 323.\*  
*Papilio Etolus. Cram. Inf.* 18. tab. 208. fig. E. F.  
Habitat in India orientali.
- Ganymedes.* 500. P. P. R. alis bicaudatis coerulefcentibus, limbo fuscis subtus basi virescentibus punctis aureis. *Gen. Inf. Mant.* 267.\*  
*Papilio Ganymedes. Cram. Inf.* 4. tab. 40. fig. C. D.  
Habitat in India.
- Mars.* 501. P. P. R. alis bicaudatis fuscis, posticis macula rubra subtus punctis duobus fasciaque obliqua albis. *Gen. Inf. Mant.* 268.\*  
*Papilio Acis. Cram. Inf.* 15. tab. 175. fig. C. D.  
*Drury Inf.* 1. tab. 1. fig. 2.  
Habitat in America meridionali. *Mus. Dom. Spengler.*
- Celerio.* 502. P. P. R. alis bicaudatis albis, subtus fasciis fuscis alternis, angulo ani ocello triplici rubro. *Gen. Inf. Mant.* 268.\*  
*Papilio Celerio. Cram. Inf.* 3. tab. 31. fig. C. D.  
Habitat Surinami.
- Ematheon.* 503. P. P. R. alis bicaudatis coeruleo nitidis, subtus atris coeruleo irroratis.  
*Papilio Ematheon. Cram. Inf.* 14. tab. 163. fig. F. G.  
Habitat Surinami.  
Alae supra coeruleae, anticis macula apicibusque nigris subtus omnes atrae atomis plurimis coeruleis nitidis.
- Didymaon.* 504. P. P. R. alis bicaudatis fuscis, subtus basi rubris punctis ocellaribus albis.  
*Papilio Didymaon. Cram. Inf.* 12. tab. 134. fig. A.  
Habitat Surinami.  
Macula virescens subtus ad augulum anl.

CLASS. VI. GLOSSATA. Papilio: 117

5. P. P. R. alis bicaudatis coeruleis, subtus atris, viridi *Pholeus*  
coeruleoque fasciatis.

*Papilio Pholeus*. *Cram. Inf.* 14. tab. 193. fig. D. E.

Habitat Surinami.

Caudae terminantur puncto albo.

6. P. P. R. alis bicaudatis subtus viridibus auro rufoque *Endymion*  
irroratis, posticis striga atra fasciaque sanguinea. *on.*

*Papilio regalis*. *Cram. Inf.* 6. tab. 72. fig. E. F.

Habitat Surinami.

Alae supra coeruleae margine atro anguloque ani sanguineo. Fascia sanguinea terminatur alba sanguineo irrorato.

7. P. P. R. alis bicaudatis, subtus anticis viridibus, auro *Venus*  
irroratis, posticis aureis viridi atroque maculatis.

*Papilio imperialis*. *Cram. Inf.* 7. tab. 76. fig. E. F.

Habitat Surinami.

Alae supra cyanae, apice nigrae. Puncta duo ferruginea in centro alae anticae. Caudae atrae.

8. P. P. R. alis bicaudatis, supra fuscis, subtus pallenscentibus, fascia rufa ocelloque terminali rubro. *Echion*  
*Syst.*

*Ent.* 519. 325. *Linn. Syst. Nat.* 2. 788. 224.

*Kleeman. Inf.* 1. tab. 7. fig. 3. 4.

9. *Papilio Endymion* alis bicaudatis fuscis macula rufa, subtus fascia maculisque duobus terminalibus rufis.

*Syst. Ent.* 519. 324.

Habitat in America.

Conf. *Papilio Eryx*. *Cram. Inf.* 12. tab. 143. fig. D.

10. P. P. R. alis bicaudatis coeruleis nitidis, posticis *Telemus*  
subtus striga marginali angulata atra. *Gen. Inf.*

*Mant.* 268.\*

*Papilio Telemus*. *Cram. Inf.* 1. tab. 4. fig. D. E.

Habitat in America meridionali.

11. P. P. R. alis bicaudatis supra fuscis immaculatis, posticis subtus angulo ani fasciis tribus coeruleis. *Hemon*  
*Gen.*

*Inf. Mant.* 269.\*

*Papilio Hemon*. *Cram. Inf.* 2. tab. 20. fig. D. E.

Habitat Surinami.

*Sphinx.* 511. P. P. R. alis bicaudatis coerulescentibus, subtus  
reis, angulo ani macula gemina atra. *Syst. L.*  
520. 326.\*

*Papilio Dindymus.* *Cram. Inf.* 4. tab. 46. fig. F. G.  
Habitat in India orientali. *D. Koenig.*

*Acmon.* 512. P. P. R. alis bicaudatis coeruleis, posticis subtus  
apice strigis viridi aureis.

*Papilio Acmon.* *Cram. Inf.* 5. tab. 51. fig. C. D.  
Habitat Surinami.

Alae supra coerulae, nitidae, anticis macula centralis

*Marfyas.* 513. P. P. R. alis bicaudatis, supra coeruleis, subtus  
lescentibus, punctis atris. *Syst. Ent.* 520. 327. *L.*  
*Syst. Nat.* 2. 788. 223. *Mus. Lud. Vlr.* 315. *C.*  
*Icon.* tab. 41. fig. 1.

*Kleeman. Inf.* 1. tab. 5. fig. 1. 2.

*Edw. Av.* 31. tab. 81.

*Seb. Mus.* 4. tab. 5. fig. 9. 10.

— — — tab. 34. fig. 15. 16.

Habitat in America.

*Bathis.* 514. P. P. R. alis bicaudatis coeruleis ocello atro, sub  
fuscis, albo fasciatis, angulo ani rufo.

*Papilio Bathis.* *Cram. Inf.* 5. tab. 51. fig. E. F.

Habitat Surinami.

Ocellus ater iride cinerea in medio alae anticae.

*Cyanus.* 515. P. P. R. alis bicaudatis coeruleis nitidis, subtus  
albo.

*Papilio Cyanea.* *Cram. Inf.* 7. tab. 76. fig. C. D.

Habitat Surinami.

Subtus in alis posticis maculae septem coerulae pupae  
atra.

*Hyacin- 516. P. P. R. alis bicaudatis coeruleo nitidis, posticis*  
*cinthus.* tus medio atro, fascia alba.

*Papilio Hyacinthus.* *Cram. Inf.* 3. tab. 36. fig. R.

Habitat in India occidentali.

*Halefus.* 517. P. P. R. alis bicaudatis coeruleis margine nigro, posticis  
cis apice utrinque auro maculatis.

*Papilio Halefus.* *Cram. Inf.* 2. tab. 97. fig. B. C.

Habitat

Habitat in America boreali.

Macula ocellaris fusca, lunula rufa in medio alae anticae. Subtus omnes fuscae, anticis puncto rufo lineoque coerulea ad basin, posticis puncto bascos rufo aureoque.

P. P. R. alis bicaudatis fuscis, subtus flauescentibus, *Formosus*. sanguineo auroque maculatis.

*Papilio Formosus*. *Cram. Inf.* 10. tab. 118. fig. G.

Habitat Surinami.

Margo posticus subtus aureus.

P. P. R. alis bicaudatis albis, limbo fusco, anticis *Meton*. macula ocellari aurea.

*Papilio Meton*. *Cram. Inf.* 17. tab. 201. fig. C. D.

Habitat Surinami.

Alae subtus nigro fusco flauoque variegatae.

P. P. R. alis dentatis subbicaudatis nigricantibus, maculis fulvis, subtus maculis argenteis. *Syst. Ent.* 520. 328. *Linn. Syst. Nat.* 2. 787. 219. *Mus. Lud. Vlr.* 328.

Habitat in Africa.

P. P. R. alis caudatis fulvis, nigro maculatis, subtus *Harpax*. cinereis, punctis aureis. *Syst. Ent. App.* 829.

Habitat in America.

P. P. R. alis caudatis fuscis, subtus flauescentibus *Croesus*. maculis rubris auro foetis.

Habitat Surinami. *Mus. Dom. Banks.*

Ferus. Alae omnes supra fuscae, anticis macula media fenestrata, posticis immaculatis. Cauda parua nigra. Subtus omnes flauescentes maculis magnis difformibus rubris, in quibus maculae punctaque aurea. Versus marginem posticum striga aurea.

P. P. R. alis caudatis coerulefcentibus, limbo fusco *Centaureus*. subtus cinereis, maculis bascos ocellaribus. *Syst. Ent.* 326. 329.\*

Habitat in noua Hollandia, *Mus. Dom. Banks.*

P. P. R. alis caudatis nigris, macula communi transversa alba, anticis ocello, posticis geminato. *Gerontes*.

Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*

Magnus in hac familia. Alae supra nigrae macula una oblonga communi alba. Anticae ocello atronula pupillari argentea. Posticae ocello magno lunulis duabus argenteis iride communi ferruginea. Margo posticus lunulis argenteis caudisque crassibus diuergentibus, apice albis. Subtus concolores, pallidiores.

*Betulae.* 525. P. P. R. alis subcaudatis fuscis, subtus luteis, posticis strigis duabus albis. *Syst. Ent.* 520. 330.

*Papilio Betulae* alis subcaudatis fuscis, primoribuscula reniformi fulua, subtus luteis fascia fulua. *Linn. Syst. Nat.* 2. 787. 220. *Fn. Sv.* 1070.

*Papilio* supra fuscus macula fulua, subtus fuluus duplici transuersa albida, alis secundariis inaequaliter caudatis. *Geoff. Inf.* 2. 58. 27.

*Papilio* minor, alis exterioribus nigricantibus macula in medio lata arcuata fulua. *Raj. Inf.* 139. 10.

*Wilk. pap.* 61. tab. 1. a. 2.

*Roef. Inf.* 1. *pap.* 2. tab. 6.

*Huffn. Inf.* tab. 12. fig. 1.

*Alb. Inf.* tab. 5. fig. 7.

*Petiv. Gazoph.* tab. 11. fig. 11.

Habitat in Betula, Pruno spinosa.

Mas macula alarum fulua distinguitur.

*Larua* obesa, viridis, lineis duabus dorsalibus et quatuor obliquis transuersis lateralibus albis.

*Puppa* glabra ferruginea.

*Pruni.* 526. P. P. R. alis subcaudatis, supra fuscis, posticis fascia marginali fulua nigro punctata. *Syst. Ent.* 331. *Linn. Syst. Nat.* 2. 788. 221. *Fn. Sv.* 1071.

*Roef. Inf.* 1. *pap.* 2. tab. 7.

*Reaum. Inf.* 1. tab. 28. fig. 6. 7.

*Schaeff. Elem.* tab. 94. fig. 5.

— — *Icon.* tab. 14. fig. 1. 2.

Habitat in Pruno.

*Larua* obesa viridis, linea laterali albicante, dentato.

*Puppa* fusca capite albo.

*Quercus.* 527. P. P. R. alis subcaudatis coerulescentibus, subtus nercis striga alba punctoque ani gemino fuluo.

*Ent.* 521. 332. *Linn. Syst. Nat.* 2. 788. 222. *Fn. Sv.* 1072.

*Papilio* supra coeruleus, subtus fuscus, lineæ transuersali vndata albicante, alis secundariis infra macula duplici fulua et in imo caudatis. *Geoff. Inf.* 2. 57. 26.

*Papilio* alis latis externis versicoloribus e nigro et coeruleo imo margine nigro cum fimbria alba cinctis, *Raj. Inf.* 130. 9.

*Petiv. Gazoph.* tab. 11. fig. 9.

*Albin. Inf.* tab. 52. fig. a. b.

*Roef. Inf.* 1. pap. 2. tab. 9.

*Wilk. pap.* 61. tab. 1. a. 1.

Habitat in Quercu.

*Larua* obesa, supra rosea, lineis tribus punctorum viridium.

*Puppa* glabra, ferruginea, lineis tribus dorsalibus punctorum fuscorum.

328. P. P. R. alis subcaudatis fuscis, macula disci alba, anticeis apice atris.

*Symethus.*

*Papilio Symethus. Cram. Inf.* 13. tab. 149. fig. B. C.

Habitat in America.

Paruus. Alae anticae basi fuscae macula media magna alba, apice atrae, subtus griseae macula media alba limboque ferrugineo. Margo posticus striga punctorum nigrorum. Posticae subcaudatae fuscae macula disci alba, subtus variegatae margine nigro punctato.

329. P. P. R. alis caudatis coerulescentibus, subtus cinereis albo vndatis, angulo ani ocello gemino aurato. *Syst.*

*Boeticus.*

*Ent.* 522. 333. *Linn. Syst. Nat.* 2. 789. 226.

Habitat in India, *D. Koenig.* In Lusitania, *D. Gray.*

330. P. P. R. alis caudatis coeruleis, posticis subtus fuscis coeruleo variis angulo ani macula gemina aurata.

*Apidanus.*

*Papilio Apidanus. Cram. Inf.* 12. tab. 137. fig. F. G.

Habitat Surinami.

Alae omnes coeruleae margine omni atro. Subtus fuscae macula baseos sanguinea, posticae coeruleo variae maculis duabus anguli ani aureis.

331. P. P. R. alis caudatis fuscis, disco coerulescentibus, subtus cinereis, fascia alba, angulo ani ocello gemino

*Strephon.*

ruf. *Syst. Ent.* 522. 344.\*

Habitat in India orientali.

- Cleon*. 532. P. P. R. alis caudatis fuscis, subtus cinereis, antice striga, posticis fascia sanguinea, angulo ani ocelli gemino rufo. *Syst. Ent.* 522. 335.\*  
Habitat in Brasilia. *Mus. Dom. Banks.*
- Alcides*. 533. P. P. R. alis caudatis nigris coeruleo nitidis, subtus ferrugineis, striga flavescente.  
*Papilio Alcides*. *Cram. Inf.* 8. tab. 96. fig. D. E.  
Habitat in Sierra Leon Africae.  
Alae omnes supra fuscae, basi coeruleo nitidae. Subtus basi brunneae apice dilutiores caudis elongatis brunneis. Striga flavescent in alis anticis abbreviata.
- Eryx*. 534. P. P. R. alis caudatis nigris subtus viridibus, postice margine subdentato maculatoque. *Syst. Ent.* 522. 336.  
*Linn. Mant.* 1. 537.  
*Papilio Amyntor*. *Cram. Inf.* 4. tab. 48. fig. E.  
*Aubert. miscell.* tab. 71. fig. 4. 5.  
Habitat in China.
- Aeolus*. 535. P. P. R. alis caudatis cyaneis, anticis macula nigra, omnibus subtus fascia alba nigro striata. *Syst. Ent.* 522. 337.\*  
*Aubert. miscell.* tab. 18. fig. 6. 7.  
Habitat in Indiis.
- Timeus*. 536. P. P. R. alis caudatis fulvo fuscoque variis, antice subtus punctis ocellaribus atris.  
*Papilio Timeus*. *Cram. Inf.* 16. tab. 186. fig. E. F.  
Habitat Surinami.  
Nimis affinis P. Virgaureae, at caudatus. Alae antice fuscae disco fulvo nigro punctato, subtus fulvae punctis annulo albo cinctis, apice fuscis. Postice fuscae punctis quatuor coerulescentibus fascisque fere marginali fulvae, subtus subfuscae margine obscuriori.
- Simaethis*. 537. P. P. R. alis caudatis fuscis, subtus viridibus, striga argentea. *Syst. Ent.* 523. 338.\*  
*Papilio Simaethis*. *Drury Inf.* 1. tab. 1. fig. 3.  
Habitat in Indiis.

8. P. P. R. alis caudatis coeruleis, margine albo maculato, subtus albis, maculis quadratis nigris fasciaque rufa. *Sedi.*

*Naturf.* 9. tab. 1. fig. 4. 5.

Habitat in Sedo Telephio Germaniae.

Magnitudo et statura P. Argi.

*Larva* viridis, linea dorsali rubra.

*Puppa* obtusa, antice virescens, postice brunnea.

9. P. P. R. alis subcaudatis, supra fuscis, subtus viridibus. *Rubi.*  
*Syst. Ent.* 523. 339. *Linn. Syst. Nat.* 2. 791. 237. *Fn.*  
*Sv.* 1077.

*Papilio* alis rotundatis integerrimis, subtus viridibus immaculatis. *Geoff. Inf.* 2. 64.

*Papilio* parva, alis supinis pullis, pronis viridibus.  
*Raj. Inf.* 133. 22.

*Albin. Inf.* tab. 5. fig. 8.

*Wilk. pap.* 62. tab. 1. a. 2.

*Petiv. Gazoph.* tab. 2. fig. 11.

*Schaeff. Icon.* tab. 29. fig. 5. 6.

Habitat in Rubo.

10. P. P. R. alis caudatis? fuscis, subtus striga obliqua niuea, posticis fascia submarginali rufa. *Ixon.*  
*Syst. Ent.* 523. 340. \*

Habitat in India. *Mus. Tottianum.*

11. P. P. R. alis caudatis albis, supra limbo punctisque disci nigris, subtus angulo ani punctis tribus aureis. *Rosimon.*  
*Syst. Ent.* 523. 341. \*

Habitat in India. *Mus. Tottianum.*

12. P. P. R. alis caudatis albis, apice fuscis, subtus albo atroque fasciatis, angulo ani punctis tribus albis. *Linus.*  
*Gen. Inf. Mant.* 269. \*

Habitat Surinami. *D. Schulz.*

13. P. P. R. alis dentato caudatis atris, basi macula fulva, *Amyntor.*  
apice fascia flava.

*Papilio Caecilia.* *Cram. Inf.* 13. tab. 159. fig. D. E.

Habitat in India.



Alae omnes supra atrae macula magna bascos fulva, fasciaque apicis flava anticarum; maiori, quae tamen margines haud attingit. Subtus flavae fascia arcuata margineque postico atris. In posticis margo posticus ater albo punctatus.

- Narcissus.* 544. P. P. R. alis dentatis fuscis, disco coerulescente, subtus fasciis maculisque sanguineis, argenteo marginatis. *Syst. Ent.* 524. 342. \*  
Habitat in noua Hollandia. *Mus. Dom. Banks.*

- Apelles.* 545. P. P. R. alis dentatis fulvis, limbo fusco, posticis subtus fasciis rufis, argenteo marginatis. *Syst. Ent.* 524. 343. \*  
Habitat in noua Hollandia. *Mus. Dom. Banks.*  
Nimis praecedenti affinis.

- Thysbe.* 546. P. P. R. alis denticulatis subcaudatis fulvis, nigromaculatis, posticis subtus striatis aureoque maculatis. *Syst. Ent.* 524. 344. *Linn. Syst. Nat.* 2. 789. 12.  
*Mus. Lud. Vlr.* 333.  
Habitat in Africa.

- Nais.* 547. P. P. R. alis dentato subcaudatis, basi argentatis, aureo striatis.  
*Papilio Nais. Cram. Inf.* 4. tab. 47. fig. D. E.  
Habitat ad Cap. bon. sp.

Paruus. Alae apice auro maculatae, anticae fuscae, posticae fulvae. Subtus anticae fulvae punctis auratis annulo nigro cinctis. Posticae fuscae maculis auratis.

- Epulus.* 548. P. P. R. alis integerrimis fuscis, fuluo maculatis, subtus variegatis.  
*Papilio Epulus. Cram. Inf.* 5. tab. 50. fig. G. D.  
Habitat Surinami.

Magnitudo P. Arionis. Alae anticae fuscae, fuluo maculatae, posticae fuscae striga macularum fulvarum pupillis atris. Subtus omnes fusco, ferrugineo alboque variegatis.

- Arion.* 549. P. P. R. alis ecaudatis supra fuscis, disco coerulesco, maculis atris, subtus canis, punctis ocellaribus. *Syst.*

*Syst. Ent.* 524. 345. *Linn. Syst. Nat.* 2. 789. 230.  
*Fn. Sv.* 1073.

*Papilio minor coerulescens. Petiv. Gazoph.* tab. 35.  
fig. 2.

*Roef. Inf.* 3. tab. 45. fig. 3. 4.

Habitat in Europae pinetis.

350. P. P. R. alis ecaudatis, posticis subtus limbo ferrugineo ocellis coeruleo argenteis. *Syst. Ent.* 525. 346. *Linn. Syst. Nat.* 2. 789. 232. *Fn. Sv.* 1074. *Argus.*

*Papilio* alis rotundatis integerrimis fuscis fascia marginali fulva, subtus cinereis, ocellis numerosis. *Geoff. Inf.* 2. 63. 32.

*Papilio* parva alis superne pullis cum ordine macularum lutearum ad imum marginem. *Raj. Inf.* 131. 12.

*Degeer Inf.* 1. tab. 4. fig. 14. 15.

*Wilk. pap.* 63. tab. 1. a. 1.

*Merian. Europ.* tab. 153.

*Huffn. Inf.* 1. tab. 4.

*Robert. Icon.* tab. 17.

*Schaeff. Icon.* tab. 29. fig. 3. 4.

351. *Papilio* alis rotundatis coeruleis, subtus ocellis numerosis. *Geoff. Inf.* 2. 61. 30.

*Papilio* parva, alis superne coeruleis, subtus cinereis, maculis nigris, circulo purpurascēte cinctis punctisque nigris depictis. *Raj. Inf.* 131. 11.

352. *Papilio* alis rotundatis coeruleis, subtus ocellorum fascia solitaria. *Geoff. Inf.* 2. 63. 31.

*Papilio* minor alia supinis purpureo coeruleis, subtus ocellorum fascia solitaria. *Raj. Inf.* 132. 17.

*Roef. Inf.* 3. tab. 37. fig. 4.

Habitat in Europae Rhamno.

Rarissime alis posticis dentatis inuenitur.

353. P. P. R. alis caudatis supra coeruleis margine nigris, subtus albidis punctis nigris sparsis. *Syst. Ent.* 525. 347. *Linn. Syst. Nat.* 2. 790. 234. *Fn. Sv.* 1076. *Argiolus.*

*Robert. Icon.* tab. 30.

Habitat in Europae pratis.

354. P. P. R. alis integris fuscis coeruleo variis, posticis *Hebrus.* coeruleis, margine interiori albo.

*Papilio*

*Papilio Hebrus.* Cram. Inf. 5. tab. 50. fig. E. F.

Habitat Surinami.

Magnitudo et statura P. Argi.

*Auius.* 553. P. P. R. alis integris concoloribus fuscis, strigis duabus apicis coeruleis.

*Papilio Auius.* Cram. Inf. 8. tab. 92. fig. B.

Habitat in India occidentali.

*Ceneus.* 554. P. P. R. alis integris atris coeruleo vndatis, margine albo punctato.

*Papilio Ceneus.* Cram. Inf. 13. tab. 156. fig. F.

Habitat Surinami.

*Erinus.* 555. P. P. R. alis integerrimis fuscis, subtus cinereis, anticis angulo postico puncto gemino nigro. *Syst. Ent.* 525. 348. \*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

*Icarus.* 556. P. P. R. alis integris supra albis, fasciis fuscis, subtus fasciis albis nigrisque alternis. *Gen. Inf. Mant.* 269. \*

*Papilio Philemon.* Cram. Inf. 2. tab. 22. fig. G. H.

Habitat Surinami.

*Parsimon.* 557. P. P. R. alis integris fuscis, subtus cinereis albo vndatis, posticis basi punctis ocellaribus nigris apiceque ocello. *Syst. Ent.* 526. 349. \*

Habitat in Sierra Leon Africae. *Mus. Dom. Banks.*

Variat supra alis anticis macula media nigra et posticis lunulis apicis albis maculaque rufa.

*Damoetes.* 558. P. P. R. alis integerrimis fuscis, subtus cinereo vndatis, posticis ocello gemino aurato. *Syst. Ent.* 526. 350. \*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

*Hylax.* 559. P. P. R. alis integerrimis supra fuscis immaculatis, subtus cinereis nigro punctatis. *Syst. Ent.* 526. 351. \*

Habitat in Malabaria. *D. Koenig.*

*Thelephus.* 560. P. P. R. alis integerrimis nigris, anticis albo punctatis, posticis supra basi rufis, subtus coeruleis nigro variis.

*Papilio Thelephus.* Cram. Inf. 6. tab. 66. fig. E. F.

Habitat

Habitat in America.

Paruus. Alae posticae supra rufae margine nigro, subtus coeruleae nigro variae.

561. P. P. R. alis integerrimis nigris, albo punctatis, anti- *Emylus*.  
cis fascia flaua, omnibus subtus variegatis.

*Papilio Emylius*. *Cram. Inf.* 6. tab. 66. fig. G. H.

Habitat Surinami.

Subtus alae omnes flauescentes, nigro variegatae, posticae tamen pallidiores.

562. P. P. R. alis integris fuluis, disco communi albo, lim- *Caricae*.  
bo cinereo subocellato. *Syst. Ent.* 526. 352. *Linn.*

*Syst. Nat.* 1. 792. 244. *Mus. Lud. Vlr.* 324.

*Clerk. Icon.* tab. 20. fig. 2.

*Merian. Swr.* 40. tab. 40. fig. 1.

*Cram. Inf.* 15. tab. 170. fig. E.

Habitat in Carica Americae.

Conf. *Papilio Pelops*. *Cram. Inf.* 15. tab. 170. fig. F.  
vix fatis distinctus.

563. P. P. R. alis integerrimis coeruleis, anticis macula *Lagus*.  
gemina alba, subtus cinereis nigro lineatis.

*Papilio Lagus*. *Cram. Inf.* 10. tab. 117. fig. F. G.

Habitat Surinami.

Limbus alarum supra ater.

564. P. P. R. alis subangulatis fuscis, macula albida, sub- *Midas*.  
tus basi albis strigaeque rufa. *Syst. Ent.* 527. 353.\*

Habitat in Indiis. *Mus. Dom. Allemand.*

565. P. P. R. alis integerrimis, supra fuscis, macula alba, *Aesopus*.  
subtus albis immaculatis.

*Papilio Thetys*. *Drury Inf.* 2. tab. 9. fig. 3. 4.

Habitat in India orientali. *Mus. Dom. Banks.*

Alae omnes supra fuscae macula magna alba, subtus  
omnes albae, nitidae, immacolatae.

566. P. P. R. alis integerrimis fuluis, margine fusco, sub- *Phae-*  
tus albis immaculatis. *drus.*

Habitat in India orientali. *Mus. Dom. Banks.*

Magnitudo P. Virgaureae. Antennae extrorsum crassiores, fuscae. Alae omnes supra fulvae, nitidae, margine

gine inprimis anticarum fusco, subtus omnes albae nitidae.

**Phareus.** 567. P. P. R. alis integerrimis subconcoloribus rubro limbo nigro.

*Papilio Phareus.* Cräm. Inf. 15. tab. 170. fig. 6.

Habitat Surinami.

Paruus. Alae omnes rubrae nitidae limbo omni atro qui in anticis dentem exferit. Subtus concolores, multo pallidiores.

**Hippothoe.** 568. P. P. R. alis integris, margine albo, subtus cinereis punctis ocellaribus numerosis. *Gen. Inf. Mant.* 276.

*Papilio Hippothoe* alis integerrimis, supra fulvis immaculatis, subtus cinerascens, punctis ocellaribus numerosis. *Linn. Syst. Nat.* 2. 793. 254. *Fn. Sv.* 1046.

*Roef. Inf.* 3. tab. 37. fig. 6. 7.

*Degeer Inf.* 2. tab. 2.

Habitat in Europae pratis.

Alae anticae colore variant, at margo tamen semper tenuissime albus.

**Virgaurea.** 569. P. P. R. alis subangulatis fulvis, margine fusco, subtus punctis nigris albisque. *Syst. Ent.* 527. 354.

*Papilio Virgaureae* alis angulatis fulvis margine fusco punctis atris sparsis. *Linn. Syst. Nat.* 2. 793. 255. *Fn. Sv.* 1079.

*Geoff. Inf.* 2. 65. 35.

*Schaeff. Icon.* tab. 97. fig. 7.

Habitat in Solidagine Virgaurea.

Variat alis supra maculatis et immaculatis.

**Phlaeus.** 570. P. P. R. alis subintegris fulvis, nigro punctatis, subtus canescentibus. *Syst. Ent.* 527. 355. *Linn. Syst. Nat.* 2. 793. 252. *Fn. Sv.* 1078.

*Papilio* rotundatis fulvis, vtrinque punctis nigris. *Geoff. Inf.* 2. 65. 36.

*Papilio* parua alis exterioribus circa margines nigris media parte rufis maculis nigris pictis. *Raj. In.* 125. 20.

*Merian. Europ.* tab. 164.

*Roef. Inf.* 3. tab. 45. fig. 5. 6.

Habitat in Europae nemoribus.

1. P. P. R. alis integris fulvis, nigro maculatis, posticis *Pentheus*.  
apice albis, nigro punctatis.

*Papilio Penthea. Cram. Inf.* 13. tab. 143. fig. E.

Habitat Surinami.

Statura P. Phlaeas. Alae anticae fulvae, nigro maculae; posticae basi fulvae, nigro maculatae, apice albae macula magna marginali fusca punctisque quatuor submarginalibus atris.

2. P. P. R. alis integris coeruleo nitidis, subtus basi *Vlrica*.  
brunneis, anticis ocello tripupillato.

*Papilio Vlrica. Cram. Inf.* 9. tab. 100. fig. E. F.

Habitat Surinami.

Alae subtus brunneae, apice oliuaceae, fascia brunnea.

3. P. P. R. alis integris, disco fuluo nigro punctato, *Thyra*.  
subtus purpurascens, maculis argenteis. *Syst. Ent.*  
527. 356. *Linn. Syst. Nat.* 2. 789. 227. *Mus. Lud.*  
*Vlr.* 329.

Habitat in Guinea.

4. P. P. R. alis integris, disco fuluo, nigro maculato, *Zenoxo*.  
anticis subtus argenteo maculatis. *Syst. Ent.* 527. 357.\*  
*Linn. Syst. Nat.* 2. 789. 231. *Mus. Lud. Vlr.* 331.

Habitat in Africa.

5. P. P. R. alis integerrimis subtestaceis, supra anticis *Lara*.  
vniocellatis, posticis biocellatis. *Syst. Ent.* 528. 358.  
*Linn. Syst. Nat.* 2. 791. 238. *Mus. Lud. Vlr.* 320.

Habitat in Africa.

Ocelli nigri iride alba.

6. P. P. R. alis integris fuscis, subtus coeruleo fasciatis, *Cephus*.  
anticis ocello vnico, posticis quatuor. *Syst. Ent.*  
528. 359.\*

Habitat in Indlis. *Mus. Dom. Gronovii*.

7. P. P. R. alis integerrimis atris, albo punctatis, macu- *Phlegeus*.  
laque bascos oblonga fulua.

*Papilio Phlegea. Cram. Inf.* 17. tab. 197. fig. F.

*Seb. Mus.* 4. tab. 34. fig. 7. 8.

Habitat Surinami.

Affinis

gine inprimis anticarum fusco, subtus omnes albae nitidae.

**Phareus.** 567. P. P. R. alis integerrimis subconcoloribus rubris limbo nigro.

*Papilio Phareus*. *Cräm. Inf.* 15. tab. 170. fig. 6.

Habitat Surinami.

Paruus. Alae omnes rubrae nitidae limbo omni atque qui in anticis dentem exferit. Subtus concolores, multo pallidiores.

**Hippothoe.** 568. P. P. R. alis integris, margine albo, subtus cineris punctis ocellaribus numerosis. *Gen. Inf. Mant.* 270.

*Papilio Hippothoe* alis integerrimis, supra fulvis immaculatis, subtus cinerascens, punctis ocellaribus numerosis. *Linn. Syst. Nat.* 2. 793. 254. *Fn. Sv.* 1046.

*Roef. Inf.* 3. tab. 37. fig. 6. 7.

*Degeer Inf.* 2. tab. 2.

Habitat in Europae pratis.

Alae anticae colore variant, at margo tamen semper tenuissime albus.

**Virgaureae.** 569. P. P. R. alis subangulatis fulvis, margine fusco, subtus punctis nigris albisque. *Syst. Ent.* 527. 354.

*Papilio Virgaureae* alis angulatis fulvis margine fusco punctis atris sparsis. *Linn. Syst. Nat.* 2. 793. 253. *Fn. Sv.* 1079.

*Geoff. Inf.* 2. 65. 35.

*Schaeff. Icon.* tab. 97. fig. 7.

Habitat in Solidagine Virgaurea.

Variat alis supra maculatis et immaculatis.

**Phlaeus.** 570. P. P. R. alis subintegris fulvis, nigro punctatis, subtus canescentibus. *Syst. Ent.* 527. 355. *Linn. Syst. Nat.* 2. 793. 252. *Fn. Sv.* 1078.

*Papilio* alis rotundatis fulvis, utrinque punctis nigris. *Geoff. Inf.* 2. 65. 36.

*Papilio* parua alis exterioribus circa margines nigris media parte rufis maculis nigris pictis. *Raj. In.* 125. 20.

*Merian. Europ.* tab. 164.

*Roef.*

*Roef. Inf.* 3. tab. 45. fig. 5. 6.

Habitat in Europae nemoribus.

571. P. P. R. alis integris fulvis, nigro maculatis, posticis *Penthæus*.  
apice albis, nigro punctatis.

*Papilio Penthæa. Cram. Inf.* 13. tab. 143. fig. E.

Habitat Surinami.

Statura P. Phlaeas. Alae anticae fulvae, nigro macula-  
tae, posticae basi fulvae, nigro maculatae, apice albae  
macula magna marginali fusca punctisque quatuor  
submarginalibus atris.

572. P. P. R. alis integris coeruleo nitidis, subtus basi *Vlricus*.  
brunneis, anticis ocello tripupillato.

*Papilio Vlrica. Cram. Inf.* 9. tab. 100. fig. E. F.

Habitat Surinami.

Alae subtus brunneae, apice oliuaceae, fascia brunnea.

573. P. P. R. alis integris, disco fulvo nigro punctato, *Thyra*.  
subtus purpurascens, maculis argenteis. *Syst. Ent.*  
527. 356. *Linn. Syst. Nat.* 2. 789. 227. *Mus. Lud.*  
*Vlr.* 329.

Habitat in Guinea.

574. P. P. R. alis integris, disco fulvo, nigro maculato, *Zenox*.  
anticis subtus argenteo maculatis. *Syst. Ent.* 527. 357.\*

*Linn. Syst. Nat.* 2. 789. 231. *Mus. Lud. Vlr.* 331.

Habitat in Africa.

575. P. P. R. alis integerrimis subtestaceis, supra anticis *Lara*.  
vniocellatis, posticis biocellatis. *Syst. Ent.* 528. 358.

*Linn. Syst. Nat.* 2. 791. 238. *Mus. Lud. Vlr.* 320.

Habitat in Africa.

Ocelli nigri iride alba.

576. P. P. R. alis integris fuscis, subtus coeruleo fasciatis, *Cephus*.  
anticis ocello vnico, posticis quatuor. *Syst. Ent.*

528. 359.\*

Habitat in Indlis. *Mus. Dom. Gronovii*.

577. P. P. R. alis integerrimis atris, albo punctatis, macu- *Phlegæus*.  
laque bascos oblonga fulva.

*Papilio Phlegæa. Cram. Inf.* 17. tab. 197. fig. F.

*Seb. Mus.* 4. tab. 34. fig. 7. 8.

Habitat Surinami.

Affinis



128 CLASS. VI. GLOSSATA. Papilio.

Affinis sequenti. Macula fulva alae anticac multo mai  
maculisque tribus atris pupilla alba.

*Aëtoris.* 578. P. P. R. alis integerrimis concoloribus fuscis, et  
punctatis.

*Papilio Aëtoris.* *Cram. Inf.* 8. tab. 93. fig. D.

Habitat Surinami.

Alae omnes fuscae, albo punctatae.

*Cassius.* 579. P. P. R. alis integerrimis, supra albis, anticis line  
fusco, posticis simbria punctata. *Gen. Inf. Mo*  
270. \*

*Papilio Cassius.* *Cram. Inf.* 2. tab. 20. fig. C. D.

Habitat Surinami.

*Talus.* 580. P. P. R. alis integerrimis, fusco coeruleoque fuscis  
subtus fascia sanguinea abbreviata.

*Papilio pygmaeus.* *Cram. Inf.* 1. tab. 7. fig. C. D.

Habitat Surinami.

Fascia sanguinea nec marginem anteriorem alae antice  
nec angulum ani attingit.

*Noleus.* 581. P. P. R. alis integerrimis concoloribus nigris, antice  
punctis decem albis, posticis disco toto albo. *Sy*  
*Ent.* 528. 361. *Linn. Syst. Nat.* 2. 792. 246. *M*  
*Lud. Vir.* 260.

*Clerk. Icon.* tab. 45. fig. 2.

*Seb. Musf.* 4. tab. 30. fig. 11. 12.

Habitat in Indiis.

*Athemon.* 582. P. P. R. alis integerrimis subfuscis, disco communi  
albo, fascia postica dimidiata fusca. *Syst. Ent.* 3  
528. *Linn. Syst. Nat.* 2. 792. 243. *Musf. Lud. V*  
323.

*Clerk. Icon.* tab. 37. fig. 2.

— — tab. 46. fig. 3.

*Seb. Musf.* 4. tab. 27. fig. 25. 26.

Habitat in America.

*Talaus.* 583. P. P. R. alis integerrimis concoloribus fuscis, antice  
punctis albis basique sanguineis, posticis disco albo  
*Linn. Syst. Nat.* 2. 792. 247. *Amoen. acad.* 6. 407.

CLASS. VI. GLOSSATA. Papilio.

129

*Clerk. Icon. tab. 45. fig. 1.*

*Seb. Musf. 4. tab. 57. fig. 18.*

Habitat in Indiis.

4. P. P. R. alis integerrimis atris, fasciis tribus flauiscentibus, postica abbreviata. *Hisbon.*

*Papilio Hisbon. Cram. Inf. 7. tab. 83. fig. C.*

Habitat in India occidentali.

Fascia prima bascos communis marginem exteriorem haud attingit, secunda in medio communis, integra, tertia abbreviata.

5. P. P. R. alis integerrimis atris, macula communi transuersali fulua. *Sagaris.*

*Papilio Sagaris. Cram. Inf. 7. tab. 83. fig. D.*

*Seb. Musf. 4. tab. 42. fig. 23. 24.*

Habitat Surinami.

Alae subtus immaculatae.

6. P. P. R. alia integris atris, macula coerulea, subtus medio rufis, fasciis albis. *Euriteus.*

*Papilio Euriteus. Cram. Inf. 13. tab. 152. fig. D. E.*

Habitat Surinami.

Alae omnes supra nigrae macula magna coerulea. Subtus omnes basi flavae, tunc rufae fasciis tribus albis, quarum vltima in alis posticis striga atra flexuosa terminatur. Posticae apice flavae macula atra dimidiato alba.

7. P. P. R. alis integerrimis concoloribus atris, anticis fascia lineari rubra. *Phereclus.*

*Syst. Ent. 529. 364. Linn. Syst. Nat. 2. 792. 248. Musf. Lud. Vlr. 326.*

*Clerk. Icon. tab. 45. fig. 4.*

*Cram. Inf. 15. tab. 178. fig. D.*

Habitat in America.

Conf. *Papilio Agyrtas. Cram. Inf. 11. tab. 123. fig. B.*

C. Differt fascia anticarum arcuata omnibusque subtus fascia flaua.

8. P. P. R. alis angulatis fuscis, omnibus striga rubra, subtus cinereo punctatis. *Lysippus.*

*Papilio alis subangulatis vtrinque atris, omnibus fascia lineari*

130 CLASS. VI. GLOSSATA. Papilio.

lineari rubra. *Linn. Syst. Nat.* 2. 793. 250. *Muf. Lud. Vlr.* 332.

*Papilio Lyfippus. Drury Inf.* 1. tab. 2. fig. 2.

*Clerk. Icon.* tab. 22. fig. 2.

*Papilio Minerea. Cram. Inf.* 8. tab. 94. fig. D. R?

Habitat in America.

Conf. *Papilio Micalia. Cram. Inf.* 8. tab. 94. fig. M  
at alae vtrunque albo punctatae.

*Thisbe.* 589. P. P. R. alis integerrimis concoloribus atris, fascia  
fulva.

Habitat Caiennae. *Muf. Dom. Blomfield.*

Statura omnino praecedentis, at distinctus alis magis  
atris coeruleo micantibus, immaculatis, excepta fascia  
fulva latiori.

*Pyramus.* 590. P. P. R. alis integerrimis fuscis coeruleo micantibus  
macula fulva, posticis subtus griseis.

Habitat Caiennae. *Muf. Dom. Teats.*

Alae omnes fuscae coeruleo micantes macula magna in  
medio fulva. Anticae subtus concolores, posticae  
griseae s. cinereo fuscoque variae.

*Priafus.* 591. P. P. R. alis integerrimis concoloribus nigris, fascia  
cis fascia duplici fulva. *Syst. Ent.* 529. 366. *Linn.*

*Syst. Nat.* 2. 793. 251. *Muf. Lud. Vlr.* 319.

Habitat in Indiis.

Macula minor fulva fascias vniens.

*Philius.* 592. P. P. R. alis ecaudatis fuscis subtus maculatis, posticis  
postice supra ocellis duobus, subtus vnico. *Syst. Ent.* 529. 367. *Linn. Syst. Nat.* 2. 790. 233.

Habitat in Africa.

*Gyas.* 593. P. P. R. alis integerrimis fuscis, subtus fulvis strigis  
punctatis aureis.

*Papilio Gyas. Cram. Inf.* 3. tab. 28. fig. F. G.

Habitat Surinami.

Paruus supra immaculatus, subtus strigis e punctis aureis.

*Aegon.* 594. P. P. R. alis integerrimis fuscis, fasciis plurimis molecularibus fulvis.

Habitat

Habitat in Iamaica Insula. *Mus. Dom. Blomfield.*

Minutus. Alae omnes supra fuscae fasciis plurimis inaequalibus macularibus fulvis. Alae posticae subtus cinerascetes.

**\*\* Plebeji Vrbicolae.**

5. P. P. V. alis integris diuaticis fuscis, anticis lineo- *Exclama-*  
la punctoque flauescens. *Syst. Ent.* 530. 373. \* *tionis.*

Habitat in Coromandel. *D. Koenig.*

6. P. P. V. alis integerrimis diuaticis fulvis lineola *Comma.*  
nigra, subtus punctis albis. *Syst. Ent.* 531. 374. *Linn.*  
*Syst. Nat.* 2. 793. 256. *Fn. Sv.* 1080.

*Papilio* alis diuaticis fulvis limbo neruisque nigris,  
primariis macula oblonga nigra. *Geoff. Inf.* 2. 66. 37.

*Petiv. Gazoph.* tab. 24. fig. 7. 8. 9.

*Merian. Europ.* 15. tab. 48.

Habitat in Europa.

*Larva* viridis albo punctata.

*Puppa* elongata, cylindrica, fusca.

7. P. P. V. alis diuaticis fulvis, fascia obliqua margi- *Augias.*  
neque postico nigris. *Syst. Ent.* 531. 375. *Linn.*  
*Syst. Nat.* 2. 794. 257. *Amoen. acad.* 4. 410. 80.

Habitat in India.

8. P. P. V. alis diuaticis fulvis, macula media margi- *Colon.*  
neque striato fuscis. *Syst. Ent.* 531. 376. \*

Habitat in India. *Mus. Tottianum.*

9. P. P. V. alis integerrimis diuaticis fuscis, flauo ma- *Paniscus.*  
culatis. *Syst. Ent.* 531. 377. \*

*Naturf.* 12. tab. 2. fig. 11. 12.

Habitat Lipsiae.

Conf. *Papilio Salius. Cram. Inf.* 6. tab. 68. fig. E.

10. P. P. V. alis integerrimis diuaticis fuscis, margine *Helirius.*  
nigro, anticis puncto centrali flauo.

*Papilio Helirius. Cram. Inf.* 5. tab. 60. fig. D.

Habitat Surinami.

Præcedentibus maior. Antennae vncinatae.

132 CLASS. VI. GLOSSATA. Papilio.

- Phineus*. 601. P. P. V. alis subangulatis diuvaricatis fuscis, fuscis  
ua anticarum interrupta.  
*Papilio Phineus*. *Cram. Inf.* 15. tab. 176. fig. E.  
Habitat Surinami.  
Antennae vncinatae. Abdomen atrum cingulis fluis.
- Helius*. 602. P. P. V. alis subdiuvaricatis fuscis, posticis margin  
interiori fuluis.  
*Papilio Helius*. *Cram. Inf.* 17. tab. 198. fig. B.  
Habitat Surinami.  
Magnitudo P. Coruma. Macula magna ouata fulua i  
margine interiori alae posticae.
- Metis*. 603. P. P. V. alis integerrimis fuscis, maculis fuluis spar  
fis, posticis subtus immaculatis. *Syst. Ent.* 528. 360.  
*Linna. Syst. Nat.* 2. 792. 245. *Mus. Lud. Vlr.* 325.  
*Drury Inf.* 2. tab. 16. fig. 3. 4.  
*Cram. Inf.* 14. tab. 162. fig. G.  
Habitat ad Promontorium bon. sp.
- Sileneus*. 604. P. P. V. alis caudatis fuscis, subtus brunneo, strig  
flauescente. *Syst. Ent.* 531. 378. \*  
Habitat in Sierra Leon Africae. *Mus. Dom. Banks.*
- Orion*. 605. P. P. V. alis caudatis fuscis, anticis maculis fenest  
tis, posticis caudis albis.  
*Papilio Orion*. *Cram. Inf.* 13. tab. 155. fig. A. B.  
*Clerk. Icon.* tab. 42. fig. 5. 6.  
Habitat Surinami.  
Alae valde caudatae caudis albis.
- Proetus*. 606. P. P. V. alis caudatis fuscis, maculis fenestratis, in  
tennis vncinatis. *Syst. Ent.* 532. 379. *Linna. Syst. Nat.*  
2. 794. 259. *Mus. Lud. Vlr.* 333.  
*Clerk. Icon.* tab. 42. fig. 1. 3.  
*Merian. Sar.* tab. 63. fig. x.  
Habitat in Americae gramine.  
Maculae alarum variant.
- Tityrus*. 607. P. P. V. alis subcaudatis fuscis, anticis vtrinque  
fiscia flaua, posticis subtus niuea. *Syst. Ent.* 532. 382. \*  
*Papilionoides caroliniana* fusca, signatura aurea pelluci  
da, subtus albida. *Petiv. Gazoph.* 51. tab. 32. fig. 5.  
*Papilio*

*Papilio Clarus*. *Cram. Inf.* 4. tab. 41. fig. E. F.  
Habitat in America.

8. P. P. V. alis subcaudatis fuscis, anticis fascia flava, *Thyrfa*.  
subtus omnibus purpureis flavo, irroratis. *Syst. Ent.*  
532. 383.\*

*Papilio Hiarbas*. *Cram. Inf.* 2. tab. 18. fig. F.?  
Habitat in America.

9. P. P. V. alis subcaudatis atris, fascia sanguinea, sub- *Alpheus*.  
tus variegatis.

*Papilio Alpheus*. *Cram. Inf.* 16. tab. 182. fig. E. F.  
Habitat ad Cap. bon. sp.  
Fascia alae anticae abbreviata.

10. P. P. V. alis subcaudatis atris, viridi maculatis, sub- *Pretus*.  
tus subferrugineis flavo maculatis.

*Papilio Pretus*. *Cram. Inf.* 16. tab. 182. fig. C. D.  
Habitat ad Cap. bon. sp.

11. P. P. V. alis subcaudatis fuscis fenestrato maculatis, *Amyntas*.  
posticis subtus punctato baseos atro. *Syst. Ent.* 533.  
384.\*

*Papilio Phocas*. *Cram. Inf.* 14. tab. 162. fig. F.?  
Habitat in America. *Mus. Tottianum*.

12. P. P. V. alis subcaudatis fulvis, posticis margine exte- *Rhetus*.  
riori coerulesco atris.

*Papilio Midas*. *Cram. Inf.* 6. tab. 63. fig. G.  
Habitat Surinami.  
Alae omnes fulvae. Margo alae anticae exterior re-  
flexus.

13. P. P. V. alis subcaudatis atris, anticis fascia niuea, *Clonias*.  
posticis subtus coerulescentibus, ano sanguineo.

*Papilio Clonias*. *Cram. Inf.* 7. tab. 80. fig. C. D.  
Habitat Surinami.  
Caudae hircus, diuergentes.

14. P. P. V. alis subcaudatis atris, anticis fascia niuea, po- *Acastus*.  
sticis subtus virescentibus, strigis abbreviatis flavis.

*Papilio Acastus*. *Cram. Inf.* 10. tab. 111. fig. D. E.  
Habitat Surinami.

Affinis praecedenti, Alae posticae subtus basi virescetes strigis quatuor abbreviatis flauescentibus.

*Palemon*. 615. P. P. V. alis subcaudatis coerulescentibus, anticis puncto medio, posticis angulo ani rufis.

*Papilio Palemon*. *Cram. Inf.* 11. tab. 131. fig. F.

Habitat Surinami.

Antennae vncinatae. Palpi strigaeque postica capitis sanguinei. Margo posticus alarum albicat.

*Coridon*. 616. P. P. V. alis subcaudatis fuscis, anticis flavo maculatis, posticis subtus fusco glaucoque variis. *Syst. Ent.* 533. 385. \*

*Papilio Talus*. *Cram. Inf.* 15. tab. 176. fig. D.

Habitat in Iamaica. *Mus. D. Hunter*.

*Arinus*. 617. P. P. V. alis ecaudatis concoloribus atris, anticis macula transuersa niuea, ano sanguineo.

*Papilio Arinus*. *Cram. Inf.* 9. tab. 100. fig. D.

Habitat Surinami.

Alae posticae subemarginatae margine albido.

*Aunus*. 618. P. P. V. alis ecaudatis fuscis, anticis fascia, posticis margine albis.

Habitat in Africa aequinoctiali. *Mus. Dom. Banks*.

Paruus. Alae omnes fuscae, vti totum corpus. Fascia obliqua alba in alis anticis et margo posticus posticorum albus.

*Alexis*. 619. P. P. V. alis ecaudatis nigricantibus, posticis subtus fascia coerulescente. *Syst. Ent.* 533. 387. \*

Habitat in India. *Mus. Tottianum*.

*Thrax*. 620. P. P. V. alis ecaudatis fuscis, maculis aliquot fensistratis, antennis vncinatis. *Syst. Ent.* 533. 338. *Lin.* *Syst. Nat.* 2. 794. 260.

*Papilio Fulgorator*. *Natursf.* 17. tab. 1. fig. 2. a. 6.

*Papilio Salus*. *Cram. Inf.* 6. tab. 68. fig. E.

Habitat in India.

Conf. *Papilio Erythus*. *Cram. Inf.* 5. tab. 59. fig. G. a. maior, vti et *P. Daunus*. *Cram. Inf.* 11. tab. 126. fig. F.

et *P. Orchamus*. *Cram. Inf.* 13. tab. 155. fig. E. F.

11. P. P. V. alis ecaudatis fuscis, anticis fenestrato maculatis, posticis subtus disco albo, nigro punctato. *Fleus.*  
Habitat in Africa aequinoctiali. *Muf. Dom. Banks.*  
Affinis praecedenti. Antennae vncinatae, nigrae.
12. P. P. V. alis dentatis fulvis nigro maculatis maculisque aliquot fenestratis. *Salus.*  
*Papilio nobilis. Cram. Inf. 9. tab. 108. fig. A. B.*  
Habitat Surinami.  
Corpus fulvum abdomine atro annulato.
13. P. P. V. alis ecaudatis flavo maculatis, posticis subtus punctis sex niueis. *Yacchus.*  
*Syst. Ent. 533. 389.\**  
Habitat in noua Hollandia. *Muf. Dom. Banks.*
14. P. P. V. alis dentatis subconcoloribus atris, anticis maculis tribus fenestratis, posticis coeruleo fasciatis. *Gnetus.*  
*Aubert. miscell. tab. 18. fig. 8. 9.*  
Habitat in India. *Muf. Dom. Banks.*  
Magnus. Antennae vncinatae, nigrae. Abdomen cingulis atris coeruleisque. Alae anticae atrae maculis tribus hyalinis interiore maiore. Posticae atrae fasciis coeruleis. Margo posticus, albo maculatus.
15. P. P. V. alis ecaudatis concoloribus fuscis, posticis apice flauis. *Anaphus.*  
*Papilio Anaphus. Cram. Inf. 15. tab. 178. fig. F.*  
Habitat Surinami.  
Antennae vncinatae nigrae. Abdomen fuscum annulis flauis. Alae fuscae, apice obscuriores, posticae apice flavae.
16. P. P. V. alis rotundatis fuscis basi virentibus, posticis subtus fascia lactea. *Bixae.*  
*Syst. Ent. 534. 390. Linn.*  
*Syst. Nat. 2. 595. 264. Muf. Lud. Vtr. 335.*  
*Clerk. Icon. tab. 42. fig. 4.*  
*Cram. Inf. 17. tab. 199. fig. C. D.*  
*Merian. Sur. tab. 44.*  
Habitat in America.  
Varietas forte P. Tityri.
17. P. P. V. alis subcaudatis viridi fuscis, anticis macula alba, subtus maculis fulvis argenteo marginatis. *Polycletus.*  
*Syst.*



136 CLASS. VI. GLOSSATA. *Papilio*.

*Syst. Ent.* 534. 391. *Linn. Syst. Nat.* 2. 795. 263.  
*Mus. Lud. Vlr.* 336.

*Clerk. Icon.* tab. 17. fig. 2.

*Sphinx Polycletus. Cram. Inf.* 13. tab. 159. fig. F. G.  
Habitat in Indiis.

*Aracini-* 628. P. P. V. alis rotundatis integerrimis fuscis, posticis  
*thus.* subtus griseis, maculis ocellaribus albis. *Gen. M.*  
*Mant.* 371. \*

Habitat in Austria. *D. Schulz.*

*Hemes.* 629. P. P. V. alis rotundatis integerrimis atris, anticis  
macula, posticis fascia albis.

*Papilio Hemes. Cram. Inf.* 9. tab. 103. fig. F.

Habitat Surinami.

Antennae vncinatae, nigrae. Alae concolores.

*Conf. Papilio Phorcus. Cram. Inf.* 13. tab. 156. fig. D.  
at maior, et *Papilio Phyllus. Cram. Inf.* 15. tab. 176.  
fig. B.

*Eumelus.* 630. P. P. V. alis rotundatis integerrimis, anticis atris  
bo maculatis, posticis fulvis nigro venosis.

*Papilio Eumelus. Cram. Inf.* 13. tab. 156. fig. E.

Habitat Surinami.

Statura omnino praecedentis. Macula parva sub  
basin alae anticae.

*Philemon.* 631. P. P. V. alis rotundatis integerrimis atris immacu-  
tis, antennis vncinatis. *Syst. Ent.* 534. 392. \*

Habitat in Americae meridionalis insulis.

*Phidias.* 632. P. P. V. alis integris atris nitentibus margine albi-  
ore anoque rubris. *Syst. Ent.* 534. 393. *Linn. Syst.*

*Nat.* 2. 795. 263. *Mus. Lud. Vlr.* 334.

*Papilio Acastus. Cram. Inf.* 4. tab. 41. fig. C. D.

— — 17. tab. 199. fig. E.

*Clerk. Icon.* tab. 44. fig. 1 — 4.

*Petiv. Gazoph.* tab. 43. fig. 15.

Habitat in Asia.

*Lycagus.* 633. P. P. V. alis rotundatis integerrimis viridi coerules  
immaculatis, ano sanguineo.

*Papilio*

*Papilio Lycagus. Cram. Inf. 15. tab. 176. fig. G.*

Habitat Surinami.

Reliquis maior.

4. P. P. V. alis rotundatis integris nigro alboque variis, posticis subtus cinereis, strigis undatis nigris. *Syrich-tus.*

*Syst. Ent. 534. 395.\**

Habitat in America.

5. P. P. V. alis integris albis, margine maculisque nigris. *Maimon.*  
*Syst. Ent. 534. 395.\**

Habitat in Asia.

Conf. *Papilio Nicaus. Cram. Inf. 2. tab. 22. fig. C.*

6. P. P. V. alis integerrimis fuscis, margine ferrugineis, *Mimas.*  
subtus striga punctorum nigrorum.

*Papilio Mimas. Cram. Inf. 5. tab. 52. fig. E. F.*

Habitat Surinami.

Antennae subuncinatae. Alae supra fusco nebulosae  
marginis postico ferrugineo anticisque punctis tribus  
approximatis albis. Subtus omnes fuscae marginis  
ferrugineo et in hoc striga punctorum nigrorum.

17. P. P. V. alis integris concoloribus flauis, limbo atro, *Gentius.*  
aplicarum fasciis flauis.

*Papilio Gentius. Cram. Inf. 15. tab. 179. fig. C.*

Habitat Surinami.

Corpus flauum. Alae omnes flauae limbo lato atro,  
anticae fasciis duabus abbreviatis punctoque inter fa-  
scias flauis in limbo.

Conf. *Papilio Procas. Cram. Inf. 15. tab. 179. fig. D.*

8. P. P. V. alis dentatis diuaticatis nigris albo macula- *Maluae.*  
tis. *Syst. Ent. 535. 396. Linn. Syst. Nat. 2. 795. 267.*

*Fn. Sv. 1081.*

*Papilio Morio. Scop. carn. 461.*

*Geoff. Inf. 2. 67. 38.*

*Petiv. Gazoph. tab. 36. fig. 6.*

*Merian. Eur. 1. tab. 38.*

*Reaum. Inf. 1. tab. 11. fig. 6. 7.*

*Rosf. Inf. 1. pap. 2. tab. 10.*

*Wlk. pap. 54. tab. 2. c. 1.*

138 CLASS. VI. GLOSSATA. *Papilio*.

*Huffnag. Inf. tab. 2.*

*Schaeff. Elem. tab. 94. fig. 9.*

Habitat in Malua, Althaca.

*Larva* grisea, capite nigro, collo punctis quatuor f  
phureis.

*Puppa* gibba, coerulefcens.

*Menalcas.* 639. P. P. V. alis integerrimis concoloribus albis, macula  
ne nigro. *Syst. Ent.* 535. 397. \*  
Habitat in Indiis.

*Menippus.* 640. P. P. V. alis integerrimis striatis fuscis, fuscis  
alba. *Gen. Inf. Mant.* 272. \*  
Habitat Surinami. *D. Scholtz.*

*Virbius.* 641. P. P. V. alis integerrimis nigris, anticis punctatis  
bus, posticis apice albis.  
*Papilio Virbius. Cram. Inf.* 12. tab. 143. fig. F.  
Habitat Surinami.  
Apex alae posticae terminatur macula magna alba.

*Phocion.* 642. P. P. V. alis integerrimis, anticis fuscis, posticis  
flavis.  
Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*  
*Parvus.* Antennae nigrae, apice flavae. Corpus fuscum  
Alae anticae, supra fuscae, subtus punctis albis innot  
tae. Posticae supra flavae margine tenuiori fuscae  
subtus flavae punctis ocellaribus albis.

*Tages.* 643. P. P. V. alis denticulatis diuaticatis fuscis, obsolete  
albo punctatis. *Syst. Ent.* 535. 398. *Linn. Syst. Nat.*  
2. 795. 268. *Fn. Sv.* 1082.  
*Papilio* alis diuaticatis cinereis punctorum alborum  
rie duplici transuersa. *Geoff. Inf.* 2. 68. 39.  
Habitat in Europae pratis.

*Nifo.* 644. P. P. V. alis integerrimis subreuerfis fuscis, anticis  
supra punctis quatuor albis sparsis. *Syst. Ent.* 535.  
399. *Linn. Syst. Nat.* 2. 796. 270. *Mus. Lud. Vh*  
339.  
Habitat in America. *Dom. v. Rohr.*

CLASS. VI. GLOSSATA. Sphinx. 139

P. P. V. alis integerrimis reuersis nigricantibus, vndique albo maculatis. *Syst. Ent.* 535. 400. *Linn. Syst. Nat.* 2. 796. 271. *Mus. Lud. Vlr.* 330. *Spio.*  
Habitat in India.

P. P. V. alis integerrimis fuscis immaculatis. *Syst. pygmaeus.*  
*Ent.* 536. 401. \*  
Habitat in India. *D. Koenig.*

142. SPHINX.

S. alis angulatis, posticis rufis ocello coeruleo. *Syst. ocellata.*  
*Ent.* 536. 1.

*Sphinx ocellata* alis repandis, posticis ocellatis. *Linn. Syst. Nat.* 2. 796. 1. *Fn. Sv.* 1083.

*Sphinx* elinguis alis angulatis, superioribus fuscis, inferioribus rubris, ocello coerulescente. *Geoff. Inf.* 2. 79. 1.

*Phalaena* maior, corpore crasso, alis angulis interioribus macula ophthalmoides insignibus. *Raj. Inf.* 361. 58.

*Phalaena* alis inferioribus macula ophthalmoides insignibus. *Alb. Inf.* tab. 8. fig. 2.

*Phalaena* subulicornis, elinguis, alis inferioribus ocello nigro, iride fusca, pupilla coerulea. *Leche nov. Inf. sp.* 58. fig. 8.

*Drury Inf.* 2. tab. 25. fig. 2. 3.

*Sepp. Inf.* 3. pag. 11. tab. 2.

*Ammis. Inf.* tab. 1.

*Harr. Aur.* tab. 5. fig. g. h.

*Roef. Inf.* 1. phal. 1. tab. 1.

*Merian. Eur.* 2. tab. 87.

*Sulz. Inf.* tab. 15. fig. 89.

*Schaeff. Icon.* tab. 99. fig. 5. 6.

Habitat in Europae, Americae Spiraea, Salice, Pomona.

*Larva* solitaria, caudata, scabra, viridis, striis obliquis lateralibus albis punctisque ocellaribus flauis.

*Eruca* maxima elegans, ligustrinae similis, cauda cornuta. *Raj. Inf.* 148. 2.

*Puppa* fusca, dorso atro.

- Nessus*. 2. S. alis angulatis, atro fasciatis, posticis rufis marginato, abdomine rufo flauoque maculato.  
*Sphinx Nessus*. *Cram. Inf.* 9. tab. 107. fig. D.  
Habitat in America boreali.  
Corpus obscurum abdomine cingulo baseos punctisque utrinque tribus pohe medium rufis. Inter haec puncta maculae duae dorsales flavae. Alae anticae subfuscae fasciis variis atris. Posticae rufae macula baseos grisea margineque postico atro.
- Quercus*. 3. S. alis angulato dentatis flavescentibus fusco subfasciatis, posticis fascia rubra.  
*Sphinx Quercus*. *Wien. Verzeichn.* 41. tab. 1. a. fig. 1.  
a. 1. b.  
— — — — — tab. 1. b. fig. 1.  
Habitat in Germaniae Quercu.  
*Larva* solitaria, caudata, viridis, striis obliquis lateralibus albis, stigmatibus rufis.  
*Puppa* brunnea marginibus rufis.  
*Imago* corpore flavescente, immaculato.
- lugubris*. 4. S. alis angulatis brunneis, anticis puncto subocellari atro. *Syst. Ent.* 537. 2.  
*Sphinx lugubris* alis angulatis ferrugineis fuscis, puncto atro. *Linn. Mant.* 537.  
*Sphinx lugubris*. *Drury Inf.* tab. 28. fig. 2.  
Habitat in Americae meridionalis insulis.
- lusca*. 5. S. alis angulatis obscuris, puncto atro, posticis atris, fascia fulva. *Gen. Inf. Mant.* 272.\*  
Habitat in Americae meridionalis insulis. *D. Spengler.*
- afiliformis*. 6. S. alis eroso dentatis, anticis cinereis, fascia obscuriore, puncto nigro, posticis rubris, margine nigro. *Syst. Ent. App.* 830.\*  
Habitat in India. *Mus. D. Edler.*
- fenestrina*. 7. S. alis eroso dentatis fusco aureis, maculis duabus fenestratis. *Gen. Inf. Mant.* 272.\*  
*Sphinx fenestrina*. *Wien. Verz.* p. 44.  
*Fuesl. Magazin.* 1. tab. 1.  
Habitat in Austria. *D. Schutz.*

CLASS. VI. GLOSSATA. Sphinx. 141

alis dentatis reuersis griseis, posticis basi ferrugineis, *Populi.*

anticis puncto albo. *Syst. Ent.* 537. 3. *Linn. Syst. Nat.* 2. 797. 2. *Fn. Sv.* 1084. *Mus. Lud. Vr.* 342. *Phinx* elinguis, alis ferratis cinereo fuscis, anticis fasciis obscurioribus transuersis, inferioribus basi macula fulua. *Geoff. Inf.* 2. 81. 3.

*Sepp. Inf.* 3. pag. 3. tab. 1.

*Roes. Inf.* 3. tab. 30.

*Merian. Europ.* 3. tab. 37.

*Wilk. pap.* 11. tab. B, C.

*Ammiral.* tab. 10.

*Degeer Inf.* 1. tab. 8. fig. 5.

*Alb. Inf.* tab. 37. fig. C.

*Schaeff. Icon.* tab. 100. fig. 6.

abitat in Populo, Salice.

*arua* solitaria, caudata, scabra, viridis, vtrinque linea strilisque transuersalibus obliquis albis. *arua* obscure grisea, postice ferruginea.

alis angulatis virescenti nebulosis, saturatius fasciatis, posticis supra luteo testaceis. *Syst. Ent.* 537. 4. *Linn. Syst. Nat.* 2. 797. 3. *Fn. Sv.* 1085. *Mus. Lud. Vr.* 343.

*Phinx* elinguis, alis laceris superioribus cinereo virefcentibus, fascia obscuriori transuersa inaequali fusco aurantiis. *Geoff. Inf.* 2. 80. 2.

*Merian. Europ.* 2. tab. 24.

*Wilk. pap.* 10. tab. 1. B. 4.

*Alb. Inf.* tab. 10.

*Roes. Inf.* 1. phal. 1. tab. 2.

*Schaeff. Elem.* tab. 116. fig. 1.

— — *Icon.* tab. 100. fig. 1. 2.

abitat in Europae Tilia.

*arua* solitaria, scabra, caudata, antice attenuata, viridis, strilis lateralibus obliquis sanguineis flauisque. *arua* obscure fusca.

8. alis eroso dentatis virefcentibus, fascia saturatiore, posticis fuluis apice atris. *Oenotherae.*

*Phinx Oenotherae.* *Wien. Verz.* 239. Tab. titul. praef.

*Fuesl. Magazin.* 2. tab. 1. fig. 9. 10.

*Phinx Proserpina.* *Pall. Spicik. fasc.* 9. tab. 2. fig. 7.

Habitat

Habitat in Germania.

Ad *Zygaenas* refert *Fuesly*, an rite? Affinis videtur  
Tiliae, at minor. Antennae nigrae, apice albae. Corpus viride. Alae croso dentatae, anticae viri  
fascia media saturatiore punctoque ocellari  
Posticae fulvae, apice atrae, margine tamen ipso

- Nerii.* 11. S. alis subangulatis viridibus, fasciis variis pallidis  
bus saturatioribus flavescentibusque. *Syst. Ent.*  
5. *Lin.* *Syst. Nat.* 2. 798. 5.  
*Roef. Inf.* 3. tab. 16.  
*Frisch. Inf.* 7. tab. 3.  
*Schaeff. Icon.* tab. 100. fig. 3. 4.

Habitat in Nerio.

*Larva* solitaria, caudata, albo punctata, collo &  
vtrinque coeruleo, cauda diflexa subarticulata.  
*Puppa* flava, linea dorsali punctisque lateralibus  
nigris.

- Apulus.* 12. S. alis dentatis, anticis fulvis, maculis duabus sub  
posticis niueis, antennis plumosis.  
*Sphinx Apulus. Cram. Inf.* 8. tab. 88. fig. E.  
Habitat Surinami.

Magnitudo S. *Nerii*. Antennae plumosae singulare  
hoc genere. Caput et thorax nigra. Abdomen  
guineum annulis atris. Alae anticae apice em  
natae, fulvae strigis duabus vndatis atris et inter  
maculae duae transversae aureae. Posticae dent  
alvae margine fulvo.

- Pylas.* 13. S. alis croso dentatis variegatis, posticis fulvis  
flavis apiceque nigris.

*Papilio Pylas. Cram. Inf.* 18. tab. 206. fig. A.  
Habitat Surinami.

Corpus paruum, brunneum, immaculatum. Alae  
sticæ fascia sesqui altera marginali atra, margine  
men ipso tenuissime flavescente.

- obscura.* 14. S. alis dentatis, posticis rubris, margine nigro,  
domine supra pallido immaculato. *Syst.*  
538. 6.\*  
Habitat in America.

15. S. alis dentatis nigris, strigis tribus approximatis pal- *Cacus.*  
lidis, posticis flavis nigro striatis. *Gen. Inf. Mant.*  
273.\*  
*Sphinx Cacus. Cram. Inf. 4. tab. 46. fig. E.*  
Habitat Surinami.
16. S. alis dentatis, posticis fuscis striga alba, abdomine *dentata.*  
albo annulato.  
*Sphinx dentata. Cram. Inf. 11. tab. 125. fig. G.*  
Habitat in Asia.  
Caput et thorax coerulescentia, fusco irrorata. Abdo-  
men fuscum, annulis albis. Alae anticae coerule-  
scentes, fusco irroratae fasciis fuscis, media maiori  
dentata.
17. S. alis subdentatis, posticis rufis, margine nigris, ab- *Ello.*  
domine pallido, cingulis atris. *Syst. Ent. 538. 7.*  
*Sphinx Ello alis integris, posticis rufis margine crassio-*  
*re nigris, abdomine pallido, cingulis atris. Linn.*  
*Syst. Nat. 2. 800. 13. Mus. Lud. Vlr. 351.*  
*Sphinx Ello. Drury Inf. 1. tab. 27. fig. 3.*  
Habitat in Indiis.
18. S. alis subdentatis, posticis nigris, basi rufis fasciaque *Iatro-*  
fenestrata, capite bicorni. *Syst. Ent. 538. 8.\** *phae.*  
*Sphinx Antaeus. Drury Inf. 2. tab. 25. fig. 1.*  
*Sphinx Hydaspus. Cram. Inf. 10. tab. 118. fig. A.*  
*Merian. Surin. tab. 38.*  
Habitat in Americae Iatropa gossypifolia.  
Larva viridis, cauda moniliformi.  
Puppa brunnea, cauda cylindrica, inflexa.
19. S. alis subdentatis, anticis viridibus, posticis fulvis, *Medea.*  
margine nigro.  
Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*  
Parua. Antennae extrorsum crassiores, fuscae. Cor-  
pus viride. Alae anticae virides puncto medio mar-  
ginalibusque fuscis, subtus pallidiores. Posticae  
fulvae margine dentato nigro, subtus virescentes  
margine fusco.
20. S. alis emarginatis nigris, apice pallidioribus strigis *Pholus.*  
nigris posticis brunneis.  
*Sphinx*



*Sphinx Phobus. Cram. Inf. 3. tab. 87. fig. B.*

Habitat in India occidentali.

Corpus medium, griseo fuscum. Alae anticae basi  
grae macula centrali pallida, apice fuscae macula  
fa strigisque duabus nigris. Posticae brunneae  
carneae margine late emarginato.

*carolina.* 21. S. alis integris, omnibus margine postico albo  
Stath., abdomine ocellis sex parium fulvis. S.  
Ent. 539. 9. Linn. Syst. Nat. 2. 798. 7. Mus. L.  
Vir. 346.

*Sphinx Carolina. Drury Inf. 1. tab. 25. fig. 1.*

Merian. Surin. 57. tab. 57.

Brown. Jamaic. 438. tab. 43. fig. 17.

Habitat in Americae Plidio.

*strigilis.* 22. S. alis integerrimis griseis, posticis fulvis strigis ful  
Syst. Ent. 539. 10.\*

*Sphinx strigilis* alis integris, superioribus supra ne  
losis puncto fusco, inferioribus luteis, strigis ful  
Linn. Mant. 1. 538.

*Sphinx strigilis. Drury Inf. 1. tab. 28. fig. 4.*

Cram. Inf. 9. tab. 106. fig. B.

Habitat in Americae meridionalis insulis.

*Atropos.* 23. S. alis integris, posticis luteis, fasciis fuscis abdomi  
Auteo, cingulis nigris. Syst. Ent. 539. 11. Linn. S.  
Nat. 2. 799. 9. Mus. Lud. Vir. 348.

*Sphinx spiritlinguis*, alis superioribus fusco nebula  
inferioribus abdomineque fasciis transuersis lute  
thorace maculis caput mortuum referentibus. G.  
Inf. 2. 85. 8.

Alb. Inf. tab. 6.

Reaum. Inf. 1. tab. 14.

— — — 2. tab. 24.

Roef. Inf. 3. tab. 1. 2.

Alt. Bresl. Sect. 13. et 37.

Sulz. Inf. tab. 15. fig. 88.

Wilk. pap. 9. tab. 1. B. 1.

Hasselqu. Itin. 417. 104. 105.

Schaeff. Icon. tab. 99. fig. 1. 2.

Cram. Inf. 7. tab. 78. fig. A.

Habi

Habitat in Iasmino, Solano tuberoso Europae australis, Aegypti.

*Larua* solitaria, caudata, flava nigropunctata, lineis lateralibus transversis, dimidiato coeruleis viridibusque, cauda diflexa.

*Puppa* brunnea stigmatibus vtrinque quinque nigris.

• *S. alis* integris, fusco coeruleoque variis, posticis fulvis, margine nigro. *Erotus*.

*Sphinx Erotus*. *Cram. Inf.* 9. tab. 104. fig. B.

Habitat — — — —

Caput et thorax fusca linea laterali pallida. Abdomen brunneum. Alae anticae fuscae macula magna bascos coerulecente, in medio punctum fulvum et margo posticus coerulecens. Posticae fulvae margine nigro.

• *S. alis* integris, obscure viridibus, lineolis coeruleis, posticis atris, maculis duabus fulvis. *Phorbas*.

*Sphinx Phorbas*. *Cram. Inf.* 5. tab. 55. fig. B.

Habitat in Indiis.

Lineolae aliquot coeruleae in alis anticis et macula obsoleta obscure ferruginea in medio.

• *S. alis* fusco nebulosis, posticis fascia nigra, anguloque ani albo. *Syst. Ent.* 540. 13. *Linn. Syst. Nat.* 2. 800. 15. *Mus. Lud. Vlr.* 353.

*Clerk. Icon.* tab. 49. fig. 2.

*Drury Inf.* 2. tab. 26. fig. 1.

*Merian. Sur.* tab. 33.

Habitat in America.

*Larua* viridis flavo lineata capite caudaque nigris.

*Puppa* brunnea.

• *S. alis* dentatis cinereis fusco variis, abdomine subtus albo, punctis quatuor ferrugineis. *Syst. Ent.* 542. 14. \* *Tetrio*.

*Sphinx Tetrio* alis denticulatis cano nebulosis abdomine subtus maculis quatuor nigris. *Linn. Mant.* 1. 538.

Habitat in America meridionali.

• *S. alis* variegatis, puncto medio albo, abdomine ocellis trium parium fulvis. *Syst. Ent.* 540. 15. \* *rustica*.

*Fabr. Sp. Inf.* T. II.

K

*Sulz.*

146 CLASS. VI. GLOSSATA. Sphinx.

*Swiz. Hist. Inf.* tab. 20. fig. 2.

Habitat in America.

*pagana.* 29. S. alis integris, cinereo nebulosis, posticis supra ferrugineo.

Habitat in India orientali. *Mus. Dom. Banks.*

Corpus totum griseum. Caput sub antennis fuscum. Linea dorsalis thoracis fusca. Alae omnes cinereo fusco nebulosae, posticis tamen disco ferrugineo. Subtus omnes cinerae, immacolatae. Femora carnea.

*Pinafiri.* 30. S. alis integris canis, anticis lineolis tribus confusis nigris, abdomine fusco, cingulis albis. *Syst.* 541. 16.

*Sphinx Pinafiri* alis integris canis, margine postico albo maculato, abdomine fusco, cingulis albis. *Linn. Syst. Nat.* 2. 802. 22. *Fn. Sv.* 1083.

*Phalaena subulicornis* fusco cinerea, thorace nigro griseo, lineis duabus nigris longitudinalibus. *G. Diff.* 28.

*Drury Inf.* 1. tab. 27. fig. 2.

*Roef. Inf.* 1. phal. 1. tab. 6.

*Sepp. Inf.* 3. p. 23. tab. 5.

*Degeer Inf.* 1. tab. 10. fig. 1. 2. 3.

*Reaum. Inf.* 1. tab. 13. fig. 8.

Habitat in Pino.

*Larva* caudata virescens, linea dorsali ferruginea, terali flava, stigmatibus ferrugineis subocellaribus. *Puppa* brunnea.

*plebeja.* 31. S. alis integris griseis, puncto medio albo, abducentibus lateribus albis nigro marginatis. *Gen. Mant.* 273.\*

Habitat in Americae meridionalis insulis. *Speng.*

*Euphorbiae.* 32. S. alis integris griseis, fasciis duabus virescentibus posticis rufis basi strigaeque nigris, antennis nigris.

*Sphinx Euphorbiae* alis integris fuscis, vitta antenarum pallida, posticis rubra. *Syst. Ent.* 541. 17. *Syst. Nat.* 2. 802. 19. *Fn. Sv.* 1086. *Mus. Vr.* 356.

*Sf.*

CLASS. VI. GLOSSATA. Sphinx. 147

*hinx spirilingius*, alis viridi fulvo purpureoque variegatis et maculatis, subtus purpureis. *Geoff.*

*Inf.* 2. 87. 11.

*Drury Inf.* 1. tab. 29. fig. 3.

*Roej. Inf.* 1. phal. 1. tab. 3.

*Frijch. Inf.* 2. tab. 11.

*Reaum. Inf.* 1. tab. 13. fig. 1. 4. 5. 6.

*Degeer Inf.* 1. tab. 8. fig. 6 — 11.

*Schaeff. Icon.* tab. 99. fig. 3. 4.

abitat in Europae Euphorbia.

*trua* caudata, nigra, albo punctata, linea dorsali sanguinea, laterali punctisque flavicantibus.

*ppa* brunnea, stigmatibus nigris.

. alis integris virescentibus, vitta alba, posticis pal- *Galii.*  
lidis basi strigae atris, antennis fuscis. *Gen. Inf.*

*Mant.* 273.\*

*hinx Galii. Wien. Verz.* pag. 42.

*hinx Galii. Naturf.* 7. 107.

*hinx sexta. Schaeff. Icon.* tab. 78. fig. 1. 2.

*hinx francofurtana. Pall. spicil. Zoolog.*

*Degeer. Inf.* 2. 236.

*Berl. Magazin.* 2. tab. 2. 4.

abitat in Germaniae Galio vero.

*trua* caudata virescens nitida, punctis vtrinque decem ocellaribus, ano sanguineo.

*ppa* brunnea.

. alis virescentibus, fascia striisque albis, posticis ni- *lineata.*  
gris, fascia rubra. *Syst. Ent.* 541. 18.\*

*hinx Daucus. Cram. Inf.* 11. tab. 125. fig. D.

abitat in America, in Lusitania. *D. Gray.*

. alis integris virescentibus, fascia vittaque albis, *Vitis.*  
posticis cinereis, fascia nigra margineque rufo. *Syst.*  
*Ent.* 542. 19.

*hinx Vitis* alis cinereis fulco variis, posticis margi-  
ne interiori sanguineis. *Linn. Syst. Nat.* 2. 801. 16.

*Mus. Lud. Vlr.* 354.

*Drury Inf.* 1. tab. 28. fig. 1.

*Wien. Verz.* 238. tab. titulo praef.

*Merian. Surin.* tab. 47. fig. 1.

abitat in Americae Vite, Magnolia glauca.

*satellitica.* 36. S. alis integris griseo nebulosis, anticis puncto ocellari sesqui altero, posticis basi cinereis. *Syst. Ent.* 542. 20. *Linn. Mant.* 1. 539.  
*Sphinx satellitia.* *Drury Inf.* 1. tab. 29. fig. 1. 2.  
Habitat in Iamaica.

*Oldenlandiae.* 37. S. alis integerrimis griseis, vitta argentea puncto medio fusco, abdomine lineis tribus punctorum argenteorum. *Syst. Ent.* 542. 21. \*  
Habitat in Indiae Oldenlandiis. *D. Koenig.*

*Lycetus.* 38. S. alis integris cinerascens, abdomine griseo vitta que lineis tribus aureis.  
*Sphinx Lycetus.* *Cram. Inf.* 5. tab. 61. fig. D.  
Habitat in India orientali.  
Nimis affinis sequenti. Thorax vtrinque linea vitta abdomen tribus aureis.

*Boerhaviae.* 39. S. alis integris cinereis, posticis rubris, apice fuscis abdomine griseo vtrinque linea aurea. *Syst. Ent.* 542. 22.\*  
*Sulz. Hist. Inf.* tab. 20. fig. 3.  
Habitat in Indiae Boerhaviis. *D. Koenig.*

*Pluto.* 40. S. alis integris virescentibus, posticis fuscis, fuscis fulva, abdomine subtus aureo. *Gen. Inf. Mus.* 274.\*  
Habitat in America meridionali. *Dom. Spengler.*

*didyma.* 41. S. alis integris fuscis, anticis punctis duobus approximatis albis. *Syst. Ent.* 543. 23.\*  
*Sphinx Penaeus.* *Cram. Inf.* 8. tab. 88. fig. D.  
Habitat in America. *Mus. Dom. Banks.*  
Conf. *Sphinx Morpheus.* *Cram. Inf.* 13. tab. 149. fig. I  
et *Sphinx Bubastus.* *Cram. Inf.* 13. tab. 149. fig. E.

*Parce.* 42. S. alis integris, anticis griseis, lunula media argentea, posticis rufescentibus. *Syst. Ent.* 543. 24.\*  
Habitat in Brasilia. *Mus. Dom. Banks.*

*Elpenor.* 43. S. alis integris, viridi purpureoque variis, posticis rubris basi atris. *Syst. Ent.* 543. 25. *Linn. Syst. Nat.* 2. 801. 17. *Fn. Sv.* 1049. *Mus. Lud. Vr.* 35  
*Sphinx*

**CLASS. VI. GLOSSATA, Sphinx. 149**

**Sphinx spirilingnis**, alis viridi purpureoque fasciatis, fasciis linearibus transuersis. *Geoff. Inf.* 2. 86. 10.

**Phalaena maior**, cauda acuta, alis angustis acutis, ex viridi fuluo et purpureo colore variis. *Raj. Inf.* 145. 2.

*Frisch. Inf.* 13. 4. tab. 2.

*Deger Inf.* 1. 694. tab. 9. fig. 8. 9.

*Petiv. Gazoph.* tab. 40. fig. 11. 12. 17.

*Wilk. pap.* 11. tab. 1. B.

*Harr. Aurel.* tab. 7. fig. g. h.

*Alb. Inf.* tab. 9. fig. 13.

*Roef. Inf.* 1. phal. 1. tab. 4.

*Merian. Europ.* 2. tab. 33. fig. 73.

Habitat in Epilobio angustifolio, Impatiente, Conuoluulo.

**Larua caudata**, fusco maculata, ocellis collaribus vtrinque duobus coeruleis.

**Puppa** antice obscure grisea, postice fusca, stigmatibus obscurioribus.

. S. alis integris, flauo purpureoque variis, abdomine *Porcellus*. subtus sanguineo albo punctato. *Syst. Ent.* 544. 26.

**Sphinx Porcellus** alis integris, margine rubris, posticis basi fuscis. *Linn. Syst. Nat.* 2. 801. 18. *Fn. Sv.* 1090.

**Sphinx spirilinguis**, alis viridi purpureoque fasciatis, fasciis serratis transuersis. *Geoff. Inf.* 2. 83. 12.

*Roef. Inf.* 1. phal. 1. tab. 5.

*Alb. Inf.* tab. 9.

*Merian. Europ.* 8. tab. 22.

Habitat in Epilobio, Impatiente.

**Larua ecaudata fusca**, ocellis vtrinque tribus collaribus coeruleis.

**Puppa nigricans**.

. S. alis integris, anticis fuscis apice variegatis, posticis atris albomaculatis. *Hylaeus*.

**Papilio Hylaeus**. *Cram. Inf.* 9. tab. 107. fig. C.

**Papilio Hylaeus**. *Drury Inf.* 2. tab. 26. fig. 9.

Habitat in America boreali.

Corpus fuscum lineis thoracis punctisque abdominis lateralibus albis. Alae anticae basi fuscae strigis se-

scutellis nigris, apice atro alboque variae, pedibus atris maculis quinque albis.

*Conuolutuli.*

46. S. alis integris nebulosis, posticis subfasciatis, abdomine cingulis rubris atris albisque. *Syst. Ent.* 544. 27.

*Sphinx Conuolutuli* alis integris, posticis nigro fasciatis margine postico albo punctatis, abdomine rubro cingulis atris. *Linn. Syst. Nat.* 2. 798. 6. *Mus. L.* 345.

*Sphinx spirilinguis*, alis omnibus fuscis fasciis dentatis obscurioribus, abdomine fasciis transuersis rubris. *Geoff. Inf.* 2. 86. 9.

*Drury Inf.* 1. tab. 25. fig. 1.

*Roef. Inf.* 1. phal. 1. tab. 7.

*Sepp. Inf.* 3. p. 19. tab. 4.

*Merian. Europ.* 39. tab. 75. fig. 2.

*Reaum. Inf.* 1. tab. 13. fig. 8.

*Wilk. pap.* 10. tab. 1. B. 2.

*Goed. Inf.* 3. tab. 5.

Habitat in Europae Conuolutulo.

*Larua* caudata vtrinque lineis lateralibus obliquis punctisque subocellatis.

*Puppa* brunnea, cornu reflexo inuoluta.

- Ligustri.* 47. S. alis integris, posticis rufis, fasciis tribus nigris, abdomine rubro, cingulis nigris, *Syst. Ent.* 545. 28. *Linn. Syst. Nat.* 2. 799. 8. *Fn. Sv.* 1087.

*Sphinx spirilinguis*, alis superioribus fuscis, inferioribus abdomineque fasciis transuersis rubris. *Geoff. Inf.* 2. 84. 7.

*Phalaena maxima*, cauda acuta, alis angustis longius acutis, abdomine roseo, areolis nigris. *Raj. Inf.* 144. 1.

*Swammerd. Bibl.* tab. 29. fig. 1. 2. 3.

*Roef. Inf.* 3. tab. 5.

*Degeer Inf.* 1. tab. 1. fig. 6.

*Reaum. Inf.* 2. tab. 20. fig. 1-4.

*Wilk. pap.* 10. tab. 1. B. 3.

*Harr. Aursl.* tab. 2. fig. a-f.

*Schaeff.*

*Schaeff. Elem.* tab. 116. fig. 2.

*Alb. Inf.* tab. 7. fig. 10.

Habitat in Europae Ligustro, Syringa, Fraxino, Sambuco.

*Larva* caudata viridis, striis lateralibus obliquis, antice incarnatis postice albis, quiescit corpore antico erecto, pedibusque eleuatis.

*Puppa* fusca, cauda quadridentata.

S. alis integris, posticis nigro fasciatis basi sanguineis, margine albo punctatis, abdomine cingulis sanguineis atrisque. *Syst. Ent.* 545. 29.\*

Habitat in America. *Dom. v. Rohr.*

S. alis integris fuscis, posticis rufis nigro striatis abdomine cinereo, annulis atris. *Caicus.*

*Sphinx Caicus. Cram. Inf.* 11. tab. 125. fig. F.

Habitat Surinami.

Caput et thorax fusca. Abdomen annulis cinereis atrisque alternis. Alae anticae fuscae striis aliquot palidioribus. Posticae rubrae striis abbreviatis atris.

S. alis integris griseis, striis albis posticis fuscis, maculis sex rubris. *Syst. Ent.* 545. 30. *Celerio.*

*Sphinx Celerio* alis integris griseis, lineola albo nigra, inferioribus basi rubris, maculis sex. *Linn. Syst.*

*Nat.* 2. 800. 12.

*Roef. Inf.* 3. tab. 8.

*Frisch. Inf.* 13. tab. 1. fig. 2.

*Cram. Inf.* 3. tab. 25. fig. 2.?

*Raj. Inf.* 159. 9.

*Petit. Gazoph.* tab. 12. fig. 9.

Habitat in Vite Europae australioris.

*Larva* caudata fusca, lineis duabus lateralibus albis, ocellisque vtrinque duobus collaribus.

*Puppa* antice fusca, postice brunnea.

S. alis integris, fuscis atro maculatis, posticis rubris, *Crantor.* margine atro.

*Sphinx Crantor. Cram. Inf.* 9. tab. 104. fig. A.

Habitat in India orientali.



Corpus cinereum. Alae anticae fuscae macula basali pallida. Maculae duae atrae ad marginem inferiorem, et duae ad marginem exteriorem, anterie lineari. Posticae rubrae marginae postico nigro angulum ani tantum maculato.

- Aleto*. 52. S. alis integris, anticis supra griseis, posticis rubris margineque atris. *Syst. Ent.* 544. 31. *Lin. Syst. Nat.* 2. 802. 20. *Mus. Lud. Vtr.* 357.  
*Drury Inf.* 2. tab. 27. fig. 4.  
*Cram. Inf.* 12. tab. 137. fig. D.  
 Habitat in India.

- Gnomia*. 53. S. alis integris cinereis, posticis supra vitta nigra. *Syst. Ent.* 546. 32.\*  
*Sphinx Butus. Cram. Inf.* 13. tab. 152. fig. A.  
 Habitat in India. *D. Koenig.*

- Hespera*. 54. S. alis integris, anticis viridi nebulosis, posticis scis immaculatis. *Syst. Ent.* 546. 33.\*  
 Habitat in India orientali. *Mus. Tottianum.*

- Anabus*. 55. S. alis integris, anticis subferrugineis, posticis atris striga punctorum albidorum.  
*Sphinx Anabus. Cram. Inf.* 11. tab. 123. fig. 6.  
 Habitat Surinami.  
 Alae anticae subferrugineae puncto paruo medio a Versus marginem strigae aliquot maculaeque maculae fuscae. Posticae atrae striga e punctis septem distinctis. Margo ipse griseus.

- Neckus*. 56. S. alis integris, anticis viridibus, striga testacea, strigis nigris, maculis bascos fuscis et flavis.  
*Sphinx Neckus. Cram. Inf.* 15. tab. 172. fig. B.  
 Habitat in America.

Corpus viride. Antennae albae. Thorax lineari alba. Abdomen lineis duabus dorsalibus et lineis nigrorum lateribusque ferrugineis. Femora alae posticae fere macularis.

- Labrusca*. 57. S. alis subfuscis, anticis subtus puncto albo, abdominis lateribus punctis quinque albis. *Syst. Ent.*

34. . *Linn. Syst. Nat.* 2. 800. 14. *Mus. Lud. Vlr.* 352.

*Sphinx Clotho* alis integerrimis, antich viridibus, posticis disco coerulescente, abdomine subtus vtrinque quinque niueis. *Syst. Ent.* 540. 12. \*

*Sphinx Labruscae.* *Cram. Inf.* 16. tab. 184. fig. A.  
*Clerk. Icon.* tab. 47. fig. 3.

*Merian. Sur.* tab. 34.

Habitat in Americis Vite.

*Larua* caudata brunnea, albo maculata, postice ocello albo.

8. S. alis integris, posticis subtus luteis, abdomine cinereo, cingulis atris. *Syst. Ent.* 547. 35. *Linn. Syst. Nat.* 2. 800. 11. *Mus. Lud. Vlr.* 350.

*Clerk. Icon.* tab. 47. fig. 1.

*Merian. Sur.* tab. 64. fig. 1. 2.

Habitat in Carica Americis.

*Larua* flauo viridique variegata.

*Puppa* testacea.

9. S. alis integris griseis puncto fusco, posticis nigris basi fasciaeque maculari albis. *Syst. Ent.* 547. 36. *terfa.*

*Sphinx terfa* alis integerrimis glaucescentibus fusco lineatis, subtus testaceis fusco punctatis. *Linn. Mant.* 1. 358.

*Sphinx terfa.* *Drury Inf.* 1. tab. 28. fig. 3.

Habitat in America meridionali.

10. S. alis ferrugineis, vitta nigricante. *Syst. Ent.* 547. *Thyelis.* 37. *Linn. Syst. Nat.* 2. 803. 24. *Mus. Lud. Vlr.* 360.

*Clerk. Icon.* tab. 46. fig. 4.

Habitat in Indiis.

*Parua.* Thorax lineis longitudinalibus fuscis.

## 143. S E S I A.

1. S. abdomine barba, segmento tertio niueo. *Syst. Tantalus.* *Ent.* 547. 1.

154 CLASS. VI. GLOSSATA. Sesia.

*Sphinx Tantalus*. Linn. *Syst. Nat.* 2. 803. 25. 1  
Lud. *Vlr.* 361.

Habitat in Europa australi, Aegypto.

*Ixion*. 2. S. abdomine barbato, striga alba rufaque, alis an  
variegatis, punctis tribus albis.

*Sphinx Tantalus*. *Cram. Inf.* 6. tab. 68. fig. F.

Habitat Surinami.

Distincta a praecedente. Alae antecae coerulesco, ni  
fuscoque variegatae punctis tribus albis. Post  
atrae margine albo. Thorax fuscus. Abdomen  
rufescens fascia bascos alba strigaque apicis rufa.

*Melas*. 3. S. alis fenestratis, margine atro, abdomine cyaneo  
si atro, cingulis tribus flavis. *Gen. Inf. Mant.* 27

*Sphinx Melas*. *Cram. Inf.* 4. tab. 45. fig. G.

Habitat Surinami.

Magnitudo sequentis. Antennae subfiliformes. C  
flaum. Thorax niger lineis duabus thoracem  
gentibus flavis. Abdomen basi atrum fasciis tri  
flavis, apice cyaneum segmentorum marginibus a  
Alae omnes fenestratae margine postico nigro.

*Hylas*. 4. S. alis fenestratis, abdomine barbato viridi, cing  
purpureo. *Syst. Ent.* 547. 2. \*

*Sphinx Hylas* alis fenestratis, abdomine fascia pur  
rascente. *Linn. Mant.* 1. 539.

*Sphinx Picus*. *Cram. Inf.* 13. tab. 148. fig. B.

Habitat in China.

*Fadus*. 5. S. abdomine barbato, alis fuscis, anticis fasciis du  
abbreviatis albis.

*Sphinx Fadus*. *Cram. Inf.* 5. tab. 61. fig. 6.

Habitat Surinami.

Statura et magnitudo S. Hylae. Corpus fuscum bi  
ani obscuriore.

*stellatarum*. 6. S. abdomine barbato, lateribus albo nigroque var  
alis posticis ferrugineis. *Syst. Ent.* 548. 3.

*Sphinx stellatarum*. *Linn. Syst. Nat.* 2. 803. 27.

*Sv.* 1091.

*Sph*

*Sphinx spirilinguis* alis superioribus fuscis nebulosis, inferioribus ferrugineis. *Linn. Syst. Nat.* 2. 83. 6. tab. 11. fig. 5.

*Papilio* velocissima, alis albis breuibus, corpore crasso inter volitandum stridorem edens. *Raj. Inf.* 133. 1.

*Roesl. Inf.* 1. phal. 1. tab. 8.

*Bradl. nat.* tab. 26. fig. 1. A.

*Reaum. Inf.* 1. tab. 12. fig. 5. 6.

*Merian. Europ.* 2. 33. tab. 29.

*Schaeff. Elem.* tab. 116. fig. 3.

— — *Icon.* tab. 16. fig. 1.

♂. *Sphinx Belis* abdomine barbato, supra nebuloso, subtus flavescente, alis posticis basi flauis. *Linn. Syst. Nat. Ed.* 10. 493. 31.

*Sphinx Belis. Cram. Inf.* 8. tab. 94. C.

Habitat in Galio, Rubo reliquisque stellatis.

*Larva* caudata, albo punctata, cauda subulata coerulescens apice ferruginea.

*Puppa* fusca.

Conf. *Sphinx Ciculus. Cram. Inf.* 13. tab. 146. fig. G.

Vix specie differt, at alae posticae atrae fascia flaua.

♂. alis brunneis immaculatis, posticis emarginatis. *Brunnus.*

*Sphinx Brunnus. Cram. Inf.* 13. tab. 147. fig. C.

Habitat Surinami.

Corpus paruum totum brunneum.

♂. alis atris, maculis duabus anticarum flauis, posticarum albis. *Syst. Ent. Append.* 830. \* *gmaculata.*

Habitat in India. *Mus. D. Hunter.*

♂. alis variegatis, posticis atris, macula bascos hyalina. *Marica.*

*Sphinx Marica. Cram. Inf.* 2. tab. 20. fig. F. G.

Habitat Surinami.

*Larva.* Alae anticae albo nigroque variegatae. Posticae atrae macula bascos hyalina. Subtus omnes atrae albo maculatae. Abdomen atrum ano barbato.

♂. abdomine barbato, postice fusco, alis nigris, macula hyalina. *Thysbe.*

7. *Sphinx Thysbe. Syst. Ent.* 548. 4. \*

Habitat in America. *Dem. Drury.*

Sp

II. S.

- fuciformis.* 11. S. abdomine barbato nigro, fascia flavescente, alis fenestratis, margine nigro. *Syst. Ent.* 548. 5.  
*Sphinx fuciformis.* Linn. *Syst. Nat.* 2. 803. 28. Fa. Sv. 1092.  
*Sphinx elinguis*, alis lanceolatis vitreis, venis nigris, superioribus macula limboque nigris fuluo nebulis. *Geoff. Inf.* 2. 82.  
*Roef. Inf.* 3. tab. 38.  
— — 4. tab. 34. fig. 1 — 4.  
*Bradl. nat.* 26. fig. 1. B.  
*Sulz. Inf.* tab. 15. fig. 90.  
*Poda Inf.* tab. 2. fig. 6.  
*Schaeff. Icon.* tab. 16. fig. 1.  
Habitat in Lonicera, Scabiosa.  
*Larua* viridis linea laterali flava cornuque rufo.  
*Puppa* folliculata nigra antice strigis flavis.
- apiformis.* 12. S. alis fenestratis, abdomine flavo, incisuris atris, thorace nigro, maculis duabus flavis. *Syst. Ent.* 349. 6.  
*Sphinx apiformis.* Linn. *Syst. Nat.* 2. 804. 29. Fa. Sv. 1093.  
*Clerk. Icon.* tab. 9. fig. 2.  
*Harr. Inf. angl.* tab. 3. fig. 7.  
*Fuesl. Magaz.* tab. 1. fig. A.  
*Sulz. Hist. Inf.* tab. 20. fig. 6.  
Habitat imago in Europae floribus, larua in Populi truncis. *Harris.*
- haemorrhoidalis.* 13. S. alis hyalinis, margine strigaeque nigris, ano barbato sanguineo. *Syst. Ent.* 549. 7. \*  
*Sphinx Leucaspis.* Cram. *Inf.* 5. tab. 52. fig. C. D. ?  
Habitat in Indiis.
- culiciformis.* 14. S. alis hyalinis margine fasciaque nigris, abdomine barbato, cingulo fuluo. *Syst. Ent.* 549. 8.  
*Sphinx culiciformis.* Linn. *Syst. Nat.* 2. 804. 30. Fa. Sv. 1094. *Amoen. acad.* 5. tab. 3. fig. 204.  
*Clerk. Icon.* tab. 9. fig. 3.  
*Fuesl. Magaz.* tab. 1. fig. B.  
Habitat in Europae floribus.  
Antennae ante apicem albae.

CLASS. VI. GLOSSATA. *Zygaena*. 157

8. alis feneſtratis, margine faſciaque nigris, abdomine *tipuliſor-*  
barbato nigro, inciſuris alternis margine flauis. *Syſt. mis.*  
*Ent.* 549. 9.

*Sphinx tipuliſiformis.* *Linn. Syſt. Nat.* 2. 804. 32. *Fn.*  
*Sv.* 1096.

*Clerk. Icon.* tab. 9. fig. 31.

*Fueſl. Magaz.* tab. 1. fig. 6.

*Harr. Inf. angl.* tab. 3. fig. 8.

Habitat in Europae floribus.

Variat interdum abdomine toto atro.

Alarum apex niger, aureo ſtriatus.

144. Z Y G A E N A.

Z. alis anticis cyaneis, punctis ſex rubris, poſticis ru- *Filipen-*  
bris, margine cyaneo. *Syſt. Ent.* 550. 1. *dulae.*

*Sphinx Filipendulae.* *Linn. Syſt. Nat.* 2. 805. 34. *Fn.*  
*Sv.* 1097.

*Sphinx ſpirilinguis*, alis ſuperioribus ſubcoeruleis pun-  
ctis ſex rubris, inferioribus rubris. *Geoff. Inf.* 2.  
88. 13.

*Papilio Phalaena Aries* antennis ſimplicibus, alis ſu-  
perioribus viridi coeruleis nitidis, maculis rubris,  
inferioribus rubris. *Degeer Inf.* 7. 591. 1. tab. 44.  
fig. 1.

*Papilio pratensis* alis pendulis minor, corpore craſſo  
nigro, alis externis coeruleo nigricantibus cum ma-  
culis ſanguineis. *Raj. Inf.* 134. 2.

*Papilionoides virescens* maculis quinque miniatis orna-  
tus. *Petiv. Muſ.* 36. 330.

*Leopardus ſylueſtris.* *Albin. Inf.* tab. 82. fig. C. D.

*Mouff. Inf.* 97. fig. infim.

*Merian. Europ.* 2. 27. tab. 17. fig. 67.

*Reaum. Inf.* 1. tab. 12. fig. 14 — 17.

— — 2. tab. 2. fig. 2.

*Roef. Inf.* 1. phal. 2. tab. 57.

*Harr. Aur.* tab. 1. fig. c — h.

*Satz. Inf.* tab. 15. fig. 91.

*Wilks pap.* 46. tab. 1. a. 1.

*Schaeff. Icon.* tab. 69. fig. 4. 5.

Goed.

*Goed. Inf.* 2. tab. 31.

*Fuesl. Magaz.* tab. 1. fig. 2. 3.

Habitat in *Spiraea Filipendula*.

*Larva* obesa, caudata, sulphurea, lineis quatuor pectorum nigrorum.

*Puppa* fusca, medio sulphureo, stigmatibus fuscis.

*Pythia*. 2. *Z. atra*, alis anticis viridibus, maculis tribus oblongis approximatis sanguineis, posticis rubris. *Gen. In Mant.* 275. \*

*Sphinx Scabiosae. Naturf.* 10. 97. 7.

*Sphinx Minos. Wien. Verz.* 45.

*Fuesl. Magaz.* tab. 1. fig. 5. 6.

Habitat in Germania australi. *D. Schulz.*

Variat rarius disco alae anticae toto rubro.

*Fulvia*. 3. *Z. cyanea*, alis anticis viridibus, punctis quinque rubris, posticis sanguineis immaculatis. *Gen. Inf. Mant.* 275. \*

*Sphinx quinta. Schaeff. Icon.* tab. 71. fig. 1.

*Fuesl. Magaz.* tab. 1. fig. 1.

Habitat in Austria. *D. Schulz.*

*Phegea*. 4. *Z. viridi atra*, alis punctis fenestratis, anteriorum sex posteriorum duobus, abdomine cingulo luteo. *Syst. Ent.* 550. 2.

*Sphinx Phegea. Linn. Syst. Nat.* 2. 805. 35. *Mus. Lud. Vlt.* 364.

*Frösch. Inf.* 6. 33. tab. 15.

*Fuesl. Magaz.* tab. 1. fig. E.

Habitat frequens in Germaniae Quercu.

*Larva* fusca, capite pedibusque rufis, dorso fasciculato.

*Ephialtes*. 5. *Z. cyanea*, alis anticis punctis sex rubris, posticis viridibus, abdomine cingulo rubro. *Syst. Ent.* 551. 3.

*Sphinx Ephialtes. Linn. Syst. Nat.* 2. 806. 36.

*Fuesl. Magaz.* tab. 1. fig. F.

*Naturf.* 10. tab. 2. fig. 7.

Habitat in Europae australioris Medicagine.

Variat punctis albis, bascos tamen semper aut rubros aut flavos.

1. *Z. nigra*, alis hyalino punctatis, anticis atris, posticis cyaneis. *Agrins.*

Habitat Surinami. *Mus. Dom. Teats.*

Corpus nigrum abdominis lateribus fulvis. Alae anticae supra atrae, in medio marginis crassioris punctis duobus approximatis posticeque striga punctorum hyalinorum. Subtus concolores, at cyaneo nitidae. Posticae cyaneae nitidae punctis duobus approximatis hyalinis.

7. *Z. nigra*, alis anticis maculis sex, posticis basi macula annulata, que flavis, abdomine flavo annulato. *Syst. Ent.* 551. 4.\*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

Variat maculis alarum flavis et hyalinis.

8. *Z. atra*, alis anticis cyaneis, punctis quinque sanguineis subocellatis, posticis rubris limbo nigro. *Gen. Inf.* *ca.*

*Mant.* 275.\*

*Sphinx carniolica.* *Scop. Ent.* 478.

*Sphinx septima.* *Schaeff. Icon.* tab. 80. fig. 4. 5.

*Fuesl. Magaz.* tab. 1. fig. G.

Habitat in Austria.

9. *Z. nigra*, alis fuscis, punctis quinque sanguineis, posticis rubris, margine fusco. *caffra.*

*Zygaena caffra* nigra, alis anticis fusco cinereis, punctis quinque sanguineis, posticis rubris. *Syst. Ent.* 551. 5. *Linn. Syst. Nat.* 2. 806. 37. *Mus. Lud. Vlr.* 362.

Habitat ad Promontorium bon. sp. *Mus. Dom. Teats.*

Differt manifeste a praecedente. Caput nigrum palpis orbitaque oculorum sanguineis. Thorax niger punctis utrinque duobus sanguineis. Puncta alae anticae sanguinea, annulo nigro cincta.

10. *Z. alis fuscis*, maculis duabus albis, abdomine cyaneo, cingulo fuluo. *Syst. Ent.* 551. 6.\* *guineensis.*

Habitat in Guinea. *D. Schloffer.*

11. *Z. alis nigris*, punctis fenestratis, anticarum sex posticarum vnico, abdomine cingulis cyaneis sanguineisque alternis. *Passalis.*

Habitat



Habitat - - - - - *Mus. Dom. Teats.*

Caput cyaneum. Thorax medio rufus, antice lateribusque cyaneus. Abdomen cylindricum cingulis sex cyaneis, quinque sanguineis. Alae omnes basi perurum aureo nitentes, anticis punctis sex posticis vnicis hyalinis.

*Cerbera*. 12. *Z. alis nigris punctis fenestratis, anticarum sex, posticarum duabus, abdomine cingulis sanguineis. Syst. Ent. 551. 7.*

*Sphinx Cerbera. Linn. Syst. Nat. 2. 808. 38. Mus. Lud. Vlr. 363.*

*Drury Inf. 1. tab. 26. fig. 2.*

*Cram. Inf. 7. tab. 83. fig. F.*

*Sulz. Hist. Inf. tab. 20. fig. 8.*

Habitat in Aethiopia.

*Thetis*. 13. *Z. coerulea, humeris rubro punctatis, alis nigris anticis apice, posticis disco hyalinis. Syst. Ent. 552. 1.\**

*Sphinx Thetis. Linn. Mant. 1. 539.*

*Drury Inf. 1. tab. 26. fig. 4.*

Habitat in Iamaica.

*Phlegmon*. 14. *Z. alis atris albo punctatis, abdomine flavo nigro lineato. Gen. Inf. Mant. 276.\**

*Sphinx Phlegmon. Cram. Inf. 3. tab. 35. fig. H.*

Habitat Surinami.

*fenestrata*. 15. *Z. alis fulcis, maculis anticarum quatuor, posticarum unica hyalinis, abdomine flavo nigroque annulato. Syst. Ent. 552. 9.\**

*Sphinx fenestrata. Drury Inf. 2. tab. 28. fig. 5.*

*Sphinx Thelebas. Cram. Inf. 13. tab. 151. fig. D.*

Habitat in China.

*Cassandra*. 16. *Z. nigra, abdomine vtrinque maculis quinque coeruleis, ano, ore basique femorum abdominisque coccineis. Syst. Ent. 552. 10.\**

*Sphinx Cassandra. Linn. Syst. Nat. 2. 806. 41.*

Habitat in America.

*Hypparchus*. 17. *Z. nigra viridi maculata, alis anticis maculis tribus posticis unica aureis.*

*Sphinx*

**CLASS. VI. GLOSSATA. *Zygaena*. 161**

*Sphinx Hypparchus*. *Cram. Inf.* 17. tab. 197. fig. 6.

Habitat in Sierra Leon Africae.

Abdomen nigrum cingulis viridibus.

1. *Z. alis virescenti atris, maculis aureis, posticis violaceis basi aureis.* *Argynnis.*

Habitat in Brasilia. *Mus. Dom. Teats.*

Porte mera praecedentis varietas. Caput et thorax viridi nitentia. Abdomen atrum ano sanguineo. Alae viridi atrae, nitidae maculis aliquot viridibus nitidis marginis exterioris maculisque tribus aureis in disco. Posticae violaceae, nitidae basi aureae.

2. *Z. atra, alis hyalinis, margine omni nigro, abdomine sanguineo, linea dorsali cingulisque atris.* *Myrtus.*

Habitat in Brasilia. *Mus. Dom. Teats.*

Caput atrum. Antennae defunt. Thorax ater lateribus coeruleis nitidis linea carnea. Abdomen cylindricum, coccineum, supra linea dorsali segmentorum marginibus atris, subtus immaculatum. Pedes atri tarsis basi albis. Margo alarum sinuato niger.

3. *Z. atra, alis hyalinis, margine fasciaque nigris, ano sanguineo.* *Syft. Ent. Append.* 830. \* *Andromacha.*

*Sphinx Leucaspes*. *Cram. Inf.* 5. tab. 52. fig. C. D. ?

Habitat in America. *Dom. Drury.*

4. *Z. ferruginea, alis hyalinis, margine omni nigro.* *Gen. Bromus. Inf. Mant.* 276. \*

*Sphinx Bromus*. *Cram. Inf.* 3. tab. 35. fig. G.

Habitat Surinami.

5. *Z. alis atris, anticis basi aureis, macula media abdomineque subtus sanguineis.* *Syft. Ent.* 552. 11. \* *Eryx.*

*Sphinx inaurata*. *Cram. Inf.* 12. tab. 140. fig. E. F.

*Sulz. Hist. Inf.* tab. 20. fig. 4.

Habitat in America meridionali. *Mus. Dom. Teats.*

6. *Z. alis hyalinis, anticis margine omni flavo.* *Egeon.*

*Sphinx Egeon*. *Cram. Inf.* 5. tab. 39. fig. B.

Habitat in China.

*Fabr. Sp. Inf. T. II.*

L

An-

162 CLASS. VI. GLOSSATA. *Zygaena*.

Antennae pectinatae. Alae anticae hyalinae margine flauo. Apex flauus macula hyalina. Posticae hyalinae margine postico nigro striga hyalina. Thorax niger albo maculatus. Abdomen nigrum dorso flauo linea longitudinali nigra.

*Achemon*. 24. *Z. flaua*, alis hyalinis apice nigris, anticis basi flauis. Habitat in Iamaica Insula. *Muf. Dom. Teats*.

Parua. Caput nigrum. Thorax flauus, immaculatus. Abdomen flauum lineis duabus dorsalibus punctorum nigrorum. Anus niger. Alae anticae hyalinae basi flauae lunula media margineque omni nigris, posticae hyalinae apice nigrae.

Variat interdum macula magna flaua ad angulum anterioris alae anticae.

*Meliffa*. 25. *Z. alis atris*, disco hyalino, abdomine nigro, lineis tribus aureis, ano fuluo. *Syst. Ent.* 553. 12. \*  
Habitat - - - - *Muf. Dom. Teats*.

*Polymena*. 26. *Z. nigra*, alis maculis luteis, anticarum tribus, posticarum duabus, abdomine cingulis duobus coccineis. *Syst. Ent.* 553. 13.

*Sphinx Polymena*. *Linn. Syst. Nat.* 2. 806. 40.

*Drury Inf.* 1. tab. 26. fig. 1.

*Cram. Inf.* 2. tab. 13. fig. D.

*Papilio parua*, alis nigris, maculis fuluis variis. *Raj. Inf.* 135.

Habitat in China.

*Lethe*. 27. *Z. alis nigris*, maculis duabus luteis, abdomine cingulis rubris nigris cyaneis alboque. *Syst. Ent.* 553. 14. \*

*Sphinx Eumolphos*. *Cram. Inf.* 17. tab. 127. fig. D.

Habitat in Africa. *Dom. Drury*.

Conf. *Sphinx Sperchius*. *Cram. Inf.* 13. tab. 146. fig. 6.

*Helymus*. 28. *Z. atra*, abdomine lineis tribus aureis. *Gen. Inf. Mant.* 276. \*

*Sphinx*

CLASS. VI. GLOSSATA. *Zygaena*. 163

*Sphinx Helymus*. *Cram. Inf.* 1. tab. 2. fig. D. E.  
Habitat Surinami.

9. *Z. alis concoloribus rubris, maculis nigris margine nigro connexis.* *Syst. Ent.* 553. 15. *fausta*.

*Sphinx fausta*. *Linn. Syst. Nat.* 2. 807. 42.

*Sulz. Hist. Inf.* tab. 20. fig. 7.

*Fuesl. Magaz.* tab. 1. fig. H.

Habitat in Europa australi.

Maculae alarum anticae subocellatae. Alae posticae rubrae, immacolatae. Abdomen cyaneo nigrum cingulo rubro.

10. *Z. alis fuscis, posticis interius sanguineis, antennis pectinatis.* *Syst. Ent.* 554. 16. *infausta*.

*Sphinx infausta*. *Linn. Syst. Nat.* 2. 807. 43.

Habitat in Europa australi.

11. *Z. virescens, alis posticis atris, fascia alba, antennis pectinatis.* *Gen. Inf. Mant.* 276. \* *Eriphia*.

Habitat Surinami. *D. Schulz.*

12. *Z. alis oblongis atris, posticis subcaudatis basi virescentibus.* *Syst. Ent. App.* 831. \* *flabellicornis*.

*Phalaena Rhodope*. *Cram. Inf.* 3. tab. 30. fig. 6.

Habitat in America meridionali. *Muf. D. Hunter.*

13. *Z. cyanea albo punctata, alis rubris, margine postico fusco.* *Syst. Ent.* 554. 17. *Pugione*.

*Sphinx Pugione*. *Linn. Syst. Nat.* 2. 807. 45.

*Sphinx Lichas*. *Cram. Inf.* 4. tab. 45. fig. B.

Habitat in America.

Antennae pectinatae apice rubrae.

14. *Z. alis apice fuscis, striga punctorum alborum, antennis pectinatis.* *Eurocilia*.

*Sphinx Eurocilia*. *Cram. Inf.* 15. tab. 178. fig. C.

Habitat in America.

Statura *Z. Pugione*. Caput nigrum albo maculatum.

Thorax testaceus margine antico punctoque flauis.

Abdomen testaceum striga dorsali punctorum alborum lateribusque atris stria flaua. Alae anticae ba-

fi testaceae macula magna media flava et in hac macula fusca, apice fuscae punctis sex albis. Posticae basi sanguinea apice atrae striis ferrugineis punctisque albis.

- pectinicornis*. 35. *Z.* subfusca, alis violaceis, fasciis duabus albis subinterruptis, collari sanguineo. *Syst. Ent.* 554. 18.  
*Sphinx pectinicornis*. *Linna. Syst. Nat.* 2. 807. 44.  
*Phalaena tiberina*. *Cram. Inf.* 3. tab. 32. fig. C. D.  
*Edw. Av.* 5. tab. 226.  
 Habitat in America.

- Ergotis*. 36. *Z.* alis atris, anticis fascia apicis alba, thorace fulvo. Habitat in Madera Insula. *Mus. Dom. Teats*.  
 Affinis sequenti. Caput nigrum, albo punctatum. Thorax fulvus, immaculatus. Alae anticae atrae margine bascos vtrinque fulvo. Fascia alba versus apicem, quae tamen margines haud attingit. Posticae nigrae, immaculae. Abdomen nigrum. Variat sexu antennis pectinatis alisque hyalino nigri.

- Glaucopis*. 37. *Z.* alis nigricantibus, anticis fascia obliqua niuea. *Zygaena Pylotis*. *Syst. Ent.* 554. 19. \*  
*Phalaena Glaucopis*. *Drury Inf.* 2. tab. 6. fig. 4.  
 Habitat in Carolina.

- Erynnis*. 38. *Z.* alis atris, anticis fascia obliqua testacea. *Gen. Inf. Mant.* 277. \*  
 Habitat Surinami. *D. Schulz*.

- Sypilus*. 39. *Z.* alis atris, anticis vitta interrupta, posticis basi testaceis, abdomine punctato.  
*Sphinx Sypilus*. *Cram. Inf.* 9. tab. 99. fig. A.  
 Habitat in America.  
 Abdomen atrum basi cingulis testaceis, apice punctis coeruleis.

- Scyton*. 40. *Z.* alis albidis, abdomine atro flavo fasciato, ano rubro.  
*Sphinx Scyton*. *Cram. Inf.* 9. tab. 99. fig. B.  
 Habitat in America.

Alae anticae albae litura media frigate postica fuscis. Posticae obscuriores disco hyalino. Corpus albidum fronte fulva.

11. *Z. nigra*, lateribus pilosis sanguineis, alis fenestratis *Auge*.  
postice nigris, antennis pectinatis.

*Zygaena Auge* sanguineo cyaneoque variegata, alis fenestratis, marginibus nigris. *Syst. Ent.* 554. 29.

*Sphinx Auge*. *Linn. Syst. Nat.* 2. 807. 46.

*Sphinx Eugrus*. *Cram. Inf.* 17. tab. 198. fig. C.

Habitat in America.

Macula niuea vtrinque ad basin abdominis.

12. *Z. alis nigris*, anticis fasciis duabus, posticis disco fenestratis, abdomine sanguineo, dorso nigro, *Syst. Ent.* 554. 21. \*

Habitat in America. *Mus. Tottianum*.

13. *Z. alis flavo hyalinis apice fuscis*, corpore auro maculato. *Gen. Inf. Mant.* 277. \*

*Sphinx coarctata*. *Cram. Inf.* 1. tab. 4. fig. F. G.

*Drury Inf.* 2. tab. 27. fig. 2. femina.

*Seb. Mus.* 4. tab. 25. fig. 17. femina.

*Pallas Spicil. Zoolog.* 9. tab. 2. fig. 8. mas.

Habitat in America meridionali.

14. *Z. alis fuscis*, anticis maculis quinque fenestratis, posticis brevissimis, macula flava. *Syst. Ent.* 555. 22. \*

Habitat in India orientali. *Mus. Tottianum*.

15. *Z. alis hyalinis*, macula fusca, posticis brevissimis filiformibus nigris. *Syst. Ent.* 555. 23. \*

Habitat in Brasilia. *Mus. Dom. Banks*.

16. *Z. alis anticis fuscis*, posticis hyalinis, pedibus posticis elongatis hirsutissimis. *Syst. Ent.* 555. 24. \*

*Sphinx tibialis*. *Drury Inf.* 2. tab. 28. fig. 2.

Habitat in Africa.

17. *Z. atra*, alis anticis puncto bascos, abdomine maculis transversalibus flavis, pedibus posticis elongatis. *Ladis*.

- Sphinx Ladis.* *Cram. Inf.* 7. tab. 83. fig. E.  
Habitat Surinami.  
Abdomen subtus atrum albo punctatum. Antennæ pectinatae.
- Capys.* 48. *Z. nigra*, alis anticis apice albis.  
*Sphinx Capys.* *Cram. Inf.* 5. tab. 56. fig. E.  
Habitat Surinami.  
Corpus totum nigrum alis anticis folis apice albis.
- Pholus.* 49. *Z. atra*, alis basi luteis. *Syst. Ent.* 555. 25. \*  
*Sphinx Pholus.* *Drury Inf.* 2. tab. 28. fig. 3.  
Habitat in Africa.
- Statice.* 50. *Z. viridi coerulca*, alis posticis fuscis. *Syst. Ent.* 555. 26.  
*Sphinx Statice.* *Linn. Syst. Nat.* 2. 808. 47. Fn. 8. 1098.  
*Sphinx viridis nitida*, antennis pectinatis clavatis, alis posticis fuscis. *Degeer Inf.* 2. t. 183. 2. tab. 3. fig.  
*Phalaena pectinicornis spirilinguis viridi coerulca*, alis posticis fuscis. *Geoff. Inf.* 2. 129. 40.  
*Papilio parua*, alis pendulis, corpore et alis vndique viridibus aut coeruleis. *Raj. Inf.* 134. 3.  
*Robert. Icon.* tab. 30. fig. 1.  
*Papilionoides pratensis viridis minor.* *Petiv. Mus.* 329.  
*Schaeff. Icon.* tab. 1. fig. 9.  
Habitat in Europae Pruno spinosa.
- Achæron.* 51. *Z. cyanea*, alis fuscis, ano rufo. *Syst. Ent.* 556. 27.  
Habitat in noua Hollandia. *Mus. Dom. Banks.*
- Styx.* 52. *Z. corpore alisque cyaneis*, thorace rufo. *Syst. Ent.* 556. 29. \*  
Habitat in Africa. *Dom. Drury.*
- Thoos.* 53. *Z. alis anticis fuscis*, posticis atris, disco subhyali collari sanguineo.  
Habitat - - - *Mus. Dom. Teats.*  
Statura præcedentis. Caput et thorax nigra collari sanguineo. Abdomen cyaneum. Alae anticae

scæ nitidæ apice parum albicantes, posticæ atræ  
disco subhyalino.

145. B O M B Y X.

\* Alis patulis.

alis patentibus falcatis luteo variis, macula fenestra- *Atlas*.  
ta, anticis sesquialtera. *Syst. Ent.* 556. 1.

*Phalaena Atlas*. *Linn. Syst. Nat.* 2. 808. 1. *Muf.*  
*Lud. Vlr.* 366.

*Cram. Inf.* 1. tab. 9. fig. A.

*Merian, Sur.* tab. 52.

*Petiv. Gazoph.* tab. 8. fig. 7.

*Valent. Muf.* 2. 168. tab. 54.

*Knorr. Del.* tab. C. 4. fig. 1.

*Seb. Muf.* 4. tab. 57. fig. 1. 3.

— — — tab. 58. fig. 6. 10.

abitat in Americae, Asiae Citrø.

*Arua* verticillata, verrucis pilosis, net folliculos gran-  
des tenaces vix extricandos.

*Inf. Phalaena Aurota*. *Cram. Inf.* 1. tab. 8. fig. A.  
Vix distincta videtur.

alis patentibus falcatis luteo variis, macula fenestra- *Hesperus*.  
ta, posticis rotundatis. *Syst. Ent.* 557. 2.

*Phalaena Hesperus*. *Linn. Syst. Nat.* 2. 809. 2. *Muf.*  
*Lud. Vlr.* 267.

*Merian. Sur.* 65. tab. 65.

*Seb. Muf.* 4. tab. 57. fig. 4. 6.

— — — tab. 58. fig. 12. 13.

*Cram. Inf.* 6. tab. 68. fig. A.

*Aubert. Icon.* tab. 66. fig. 1.

*Sulz. Hist. Inf.* tab. 21. fig. 2.

abitat in Americae Citrø.

*Arua* tota glabra.

alis patentibus griseis, fascia fulva, anticis ocello *Cecropia*.  
subfenestrato ferrugineo. *Syst. Ent.* 557. 3.

*Phalaena Cecropia*. *Linn. Syst. Nat.* 2. 809. 3. *Muf.*  
*Lud. Vlr.* 368.

*Drury Inf.* 1. tab. 18. fig. 2.

*Cram. Inf.* 4. tab. 42. fig. A. B.

L 4

Clerk.



*Clerk. Icon. tab. 49. fig. 1.*

*Seb. Mus. 4. tab. 58. fig. 4. 5.?*

*Catesb. Carol. 86. tab. 86.*

Habitat in America boreali.

*Paphia.* 4. B. alis patentibus falcatis concoloribus flauis, strigis rufis ocelloque fenestrato. *Syst. Ent. 557. 4.\**

*Phalaena Paphia pectinicornis*, elinguis flaua, albifalcatis concoloribus ocello fenestratis. *Lin. Syst. Nat.*

3. 809. 4. *Mus. Lud. Vr. 369.*

*Petiv. Gazoph. tab. 29. fig. 3.*

*Rumph. Amb. 3. tab. 75.*

Habitat in Asia.

*Polyphemus.* 5. B. alis patentibus falcatis griseo carneis, fascia atra ocelloque fenestrato posticarum maiori.

*Phalaena Polyphemus. Cram. Inf. 1. tab. 5. fig. A. B.*

Habitat in America boreali, Iamaica.

Corpus ferrugineum capite abdomineque cinereis. Alae anticae supra basi apiceque carnae, in medio cinereae, qui color antice a striga alba, postice a fascia atra strigae albae adnatae includitur. Ocellus parvus fenestratus annulo flauo nigroque cinctus in disco. Posticae cinereae apice carnae fascia atra strigae albae adnata ante apicem. In disco ocellus magnus ater pupilla fenestrata, iride flaua. Subtus omnia obscuriores ocello minori.

*Cytheres.* 6. B. alis patentibus concoloribus griseis, strigis cinereis ocelloque fenestrato. *Syst. Ent. 557. 5.\**

*Sulz. Hist. Inf. tab. 21. fig. 1.*

Habitat ad Cap. bon. sp. *Mus. Dom. Banks.*

*Myliitta.* 7. B. alis patentibus falcatis flauis, striga ferruginea ocelloque fenestrato diuiso. *Syst. Ent. 558. 6.\**

*Phalaena Myliitta. Drury Inf. 2. tab. 5. fig. 1.*

*Phalaena Paphia. Cram. Inf. 13. tab. 146. fig. A.*

*Seb. Mus. 4. tab. 23. fig. 5. 6.*

Habitat in lamboliferis Indiae orientalis. *D. Koenig.*

*Promethea.* 8. B. alis patulis subfalcatis margine griseis, anticis utrinque ocello atro. *Syst. Ent. 558. 7.*

*Pha-*

*Phalaena Promethea*. Drury Inf. 2. tab. 11. fig. 1. 2. mas.

— — — tab. 12. fig. 12. fem.

*Cram. Inf.* 7. tab. 75. fig. A. B. fem.

— — — tab. 76. fig. A. B. mas.

Habitat in America boreali.

*Larva* viridis nigro punctata, antice spinis quatuor rubris.

9. B. alis patulis fuscescentibus, striga vndata pallida. *Erythri-*

*Phalaena Armida*. *Cram. Inf.* 17. tab. 197. fig. A. *uae.*

*Phalaena Cassandra*. *Cram. Inf.* 17. tab. 197. fig. B.

*Merian. Sur.* 11. tab. 11.

*Seb. Musf.* 4. tab. 21. fig. 1. 2. 5. 6.

Habitat in Americae meridionalis Erythrini.

*Larva* flavescent spiraculis atris, spinis collaribus quatuor, caudalibus atris.

20. B. alis patulis, anticis variegatis, subtus ocello atro, *Ianus.*  
posticis sanguineis, ocello atro.

*Phalaena Ianus*. *Cram. Inf.* 6. tab. 64. fig. A. B.

Habitat Surinami.

Magna. Alae anticae supra obscurae, variegatae, sub-  
tus griseae ocello magno atro pupilla oblonga alba.  
Posticae supra sanguineae, in medio ocello magno  
atro pupilla lunata alba, apice pallidae, subtus ci-  
nereae.

21. B. alis patulis subfalcatis fuscis, margine pallidioribus, *Hippoda-*  
striga alba anticarum interrupta. *mia.*

*Phalaena Hippodamia*. *Cram. Inf.* 11. tab. 126. fig. B.

Habitat Surinami.

Affinis B. Prometheae, paullo maior. Alae omnes fuscae  
margine imprimis anticarum pallidiore. Striga alba  
colores distinguit anticarum apice striga vndata fer-  
ruginea terminatur et pone hanc punctum subocella-  
re coeruleum.

22. B. alis patentibus fusco incarnatis, posticis ocello fer- *missitans.*  
rugineo, pupilla fenestrata. *Syst. Ent.* 558. 8.\*

Habitat in Africa. *Musf. Dom. Banks.*

Femina maior ocello alae posticae atro.

170 CLASS. VI. GLOSSATA. Bombyx.

- Semiramis.* 13. B. alis patentibus caudatis versicoloribus, punctis fenestrato, caudis longissimis. *Gen. Inf. Mant.* 276.  
*Phalaena Semiramis. Cram. Inf.* 2. tab. 13. fig. A.  
 Habitat in Americae australis Saccharo.
- Boreas.* 14. B. alis patulis caudatis cinereis fusco variis, antennis punctis duobus, posticis unico fenestratis.  
*Phalaena Boreas. Cram. Inf.* 6. tab. 70. fig. B.  
 Habitat in America.  
 Alae anticae falcatae, posticae caudatae.
- Luna.* 15. B. alis patentibus caudatis concoloribus virescentibus oculo disci lunato. *Syst. Ent.* 558. 9.  
*Phalaena Luna. Linn. Syst. Nat.* 2. 810. 5. *Mus. Lul.* Vlr. 370.  
*Clerk. Icon.* tab. 52. fig. 1.  
*Drury Inf.* 1. tab. 124. fig. 1.  
*Cram. Inf.* 1. tab. 2. fig. A.  
 — — 3. tab. 31. fig. A. B.  
*Petiv. Gazoph.* tab. 14. fig. 5.  
*Catesb. Carol.* 2. 84. tab. 84.  
 Habitat in America boreali.
- Epimethea.* 16. B. alis patulis caudatis subfuscis, striga alba, posticis oculo disci fuluo.  
*Phalaena Epimethea. Drury Inf.* 2. tab. 13. fig. 1.  
*Cram. Inf.* 15. tab. 176. fig. A.  
 Habitat in Guinea.
- Argus.* 17. B. alis patentibus caudatis pallide ferrugineis, punctis ocellaribus fenestratis numerosis, caudis longissimis.  
 Habitat — — — *Mus. Dom. Banks.*  
 Corpus paruum, pallide ferrugineum, antennis pectinatis, atris. Alae omnes pallide ferrugineae. Anticae punctis sex in medio approximatis fenestratis annulo fuluo nigroque cinctis. Posticae punctis quinque sparsis eiusdem coloris. Caudae corporis triplo longiores, ferrugineae, apice pallidae.
- fenestra.* 18. B. alis patentibus concoloribus flauis, anticis fenestra duplici, posticis simplici. *Syst. Ent.* 559. 10.

*Phalae*

CLASS. VI. GLOSSATA. Bombyx. 171

*Phalaena fenestra.* Linn. *Syst. Nat.* 2. 811. 9. *Muf.*  
Lud. *Vlr.* 372.

Clerk. *Icon.* tab. 55. fig. 1.

Habitat in Indiis.

9. B. alis patentibus flavescentibus, fusco irroratis, ocel- *Penelope.*  
lo centrali fenestrato. *Gen. Inf. Mant.* 278.\*

*Phalaena Penelope.* Cram. *Inf.* 4. tab. 45. fig. A.

Habitat in India.

10. B. alis patentibus griseis, strigis albis, ocello centrali *Tyrrhea.*  
atro, pupilla fenestrata. *Gen. Inf. Mant.* 278.\*

*Phalaena Tyrrhea.* Cram. *Inf.* 4. tab. 46. fig. A.

Habitat ad Cap. Bon. Spei.

11. B. alis patentibus fuscis, anticis fascia fenestrata breui. *perspicua.*  
*Syst. Ent.* 559. 11.

*Phalaena perspicua.* Linn. *Syst. Nat.* 2. 811. 10. *Muf.*

Lud. *Vlr.* 373.

Habitat in Indiis.

12. B. alis patulis concoloribus luteis, apice maculisque *militaris.*  
violaceis, anticis extus albo maculatis. *Syst. Ent.*  
559. 12.

*Phalaena militaris.* Linn. *Syst. Nat.* 2. 811. 12. *Muf.*

Lud. *Vlr.* 375.

Roef. *Inf.* 4. tab. 6. fig. 3.

Cram. *Inf.* 3. tab. 29. fig. B.

Aub. *Miscell.* tab. 67. fig. 1.

Habitat in Asia, China.

13. B. alis patentibus rotundatis medio albis, anticis ocel- *Castalia.*  
lo, posticis puncto fusco. *Syst. Ent.* 559. 13.\*

Habitat in noua Hollandia. *Muf. Dom. Banks.*

14. B. alis patentibus rotundatis griseo nebulosis subfa- *pauonia.*  
sciat, ocello nictitante subfenestrato. *Syst. Ent.*  
559. 14.

*Phalaena pauonia.* Linn. *Syst. Nat.* 2. 810. 7. *Fn.*

Sv. 1099.

1. *Phalaena pectinicornis* elinguis, alis cinereo fu- *minor.*  
scis planiusculis, singulis ocello fasciaque fuscis.

*Geoff. Inf.* 2. 100. 3.

*Phalae.*

*Phalaena maior pulcra*, maculis ophthalmoidibus in singulis alis singulis. *Raj. Inf.* 146. 1.

*Wilk. pap.* 15. tab. 2. a. 3.

*Roef. Inf.* 1. phal. 2. tab. 5.

*Alb. Inf.* tab. 37.

*Schaeff. Elem.* tab. 98. fig. 2.

— *Icon.* tab. 89. fig. 2 — 5.

*Merian. Europ.* tab. 13. fig. 23.

*Degeer Inf.* 1. tab. 19. fig. 1 — 12.

*maior.* b. *Phalaena pectinicornis* elinguis, alis cinereo fuscis planiusculis, singulis ocello maior. *Geoff. Inf.* 100. 1.

*Roef. Inf.* 4. tab. 15. 16. 17.

*Reaum. Inf.* 1. tab. 47. 48. 49.

*Knorr. Delic.* tab. 6. 2. fig. 2.

*Seb. Mus.* 4. tab. 60. fig. 9 — 14.

Habitat in Erica, Pomonae arboribus.

Mas antennis magis pectinatis alisque posticis basi fuscis uescentibus.

*Larva* gregaria, viridis, verticillata, verrucis rubris flavisque piliferis, pilis setaceis.

*Puppa* nigricans, folliculata, folliculo apice foramine elastico dehiscente.

*Achelous.* 25. B. alis patulis ferrugineis, omnibus fascia anticorum puncto albis.

*Phalaena Achelous. Cram. Inf.* 10. tab. 101. fig. A.

Habitat in America.

Magna. Pone fasciam albam fascia postice dentata fuscis

*argulata.* 26. B. alis patulis eroso dentatis, posticis truncatis obtusissimis.

*Phalaena argulata. Cram. Inf.* 6. tab. 61. fig. E. F.

Habitat Surinami.

Color alarum anticarum supra basi fuscus, apice ferrugineus, posticarum subferrugineus, subtus basi cinereus rufus irroratus, apice rufus.

*Ton.* 27. B. alis patulis testaceis, ocello subviolaceo, pupa haustata alba. *Syst. Ent.* 560. 15.

*Phalaena Tuu. Linn. Syst. Nat.* 2. 811. 8. *Fr. Sp. 1mo.*

*Aff. Holm.* 149. 130. tab. 4. fig. 8.

*Schaeff.*

CLASS. VI. GLOSSATA. Bombyx. 173

*Schaeff. Icon. tab. 85. fig. 4 — 6.*

*Roef. Inf. 4. tab. 7. fig. 3. 4.*

Habitat in Betula.

Ocellus in alis anticis vtrunque, in posticis subtus pupilla hastata alba absque iride.

Larva viridis, striis obliquis lateralibus albis, dorso nodoso.

Puppa brunnea pilosa. *Roef. Inf. 3. tab. 68.*

B. alis patulis flavis, anticis subtus, posticis supra ocello atro, pupilla alba. *Syst. Ent. 560. 16.\**

Habitat in America. *Mus. D. Hunter.*

B. alis patulis fuscis, posticis cinereis ocello rufo. *Abas.*

*Phalaena Abas. Cram. Inf. 7. tab. 77. fig. A. B.*

Habitat Surinami.

Ocellus magnus rufus iride atra pupillaque alba in medio alae posticae.

B. alis patulis, anticis fuscis striga atra, posticis rufis ocello nigro, lunula alba. *Salmonia.*

*Phalaena Salmonia. Cram. Inf. 14. tab. 162. fig. A.*

Habitat Surinami.

Magna. Corpus fuscum annulis abdominalibus rufis.

Alae anticae fuscae in medio lineolis duabus transversis rufis strigaque postica atra. Posticae rufae ocello magno atro iride flava pupillaque lunata alba. Pone ocellum striga tenuissima, latiorque ferruginea. Subtus omnes puncto centrali albo.

B. alis patulis rotundatis nigris, fascia alba, macula subocellari nigra. *Proserpina.*

*Syst. Ent. 561. 17.\**

*Phalaena Maja. Drury Inf. 2. tab. 24. fig. 3.*

*Cram. Inf. 9. tab. 98. fig. A.*

Habitat in America. *Mus. D. Hunter.*

\* \* Alis reuerfis.

B. alis reuerfis dentatis ferrugineis ore tibiisque nigris. *quercifolia.*

*Syst. Ent. 561. 19.*

*Phalaena quercifolia* clinguis, alis reuerfis semitectis dentatis ferrugineis, margine postico nigro. *Linn.*

*Syst. Nat. 2. 812. 18. F. Sv. 110.*

*Phalas-*

*Phalaena pectinicornis* elinguis, tota rufa, alae margine ferrato. *Geoff. Inf.* 2. 110.

*Phalaena* tota obscure rufa. *Raj. Inf.* 141. 1.

*Alb. Inf.* 1. tab. 16.

*Merian. Europ.* tab. 1. fig. 3.

*Frisch. Inf.* 3. tab. 1. fig. 3.

*Roef. Inf.* 1. phal. 2. tab. 41.

*Reaum. Inf.* 2. tab. 23.

*Wilk. pap.* 27. tab. 3. b. 1.

*Sulz. Inf.* tab. 16. fig. 93.

*Poda.* tab. 2. fig. 7.

*Schaeff. Icon.* tab. 71. fig. 4. 5.

Habitat in Pyro, Gramine, Salice.

*Larva* subcaudata, pilosa, ferruginea, collaribus ruleis.

*Larva* maxima ex albo et nigro coloribus variata.

*Raj. Inf.* 342. 2.

*Puppa* folliculata fusca, fasciis rubris.

*ilicifolia.* 33. B. alis reuerfis semitectis ferratis griseis, margine stico albo punctato. *Syst. Ent.* 562. 20.

*Phalaena ilicifolia.* *Linn. Syst. Nat.* 2. 813. 19. *Sv.* 1180.

*Phalaena* elinguis, antennis pectinatis, alis dentis reuerfis brunneis, strigis obscurioribus. *Degeer V. Germ.* 2. 1. 212. 1.

*Degeer Inf.* 1. tab. 14. fig. 7. 9.

Habitat in Salice.

*Promula.* 34. B. alis reuerfis? subdentatis fuscis immaculatis, aldomine brunneo.

*Phalaena Promula.* *Cram. Inf.* 6. tab. 72. fig. D.

Habitat in Iaua.

Antennae pectinatae, brunneae. Alae, ut videtur, reuerfae fuscae, immaculatae posticis subdentatis.

*capensis.* 35. B. alis reuerfis fuluolis, anticis strigis duabus repandis, posteriore nigra inducta. *Syst. Ent.* 562. 24.

*Phalaena capensis.* *Linn. Syst. Nat.* 2. 818. 20.

Habitat in Africa.

*Alneo.* 36. B. alis reuerfis fuscis, apice cinereis.

Habitat ad Cap. Bon. Spei.

**CLASS. VI. GLOSSATA. Bombyx. 179**

Corpus magnum, minime nitidum. Caput et thorax cinereo fusca. Alae anticae fuscae margine postico late cinereo. Subtus omnes ferrugineae atomis numerosissimis cinereis. Abdomen ferrugineo hirtum. Pedes hirti plantis albo annulatis.

B. alis reuerfis fuluolis, posticis basi subtus ferrugineis. *Syst. Ent. 562. 22.\** *Australasiae.*  
Habitat in noua Hollandia. *Mus. Dom. Banks.*

B. alis reuerfis rufo cinereis, fasciis duabus lineari- *Fagi.*  
bus luteis flexuosis. *Syst. Ent. 562. 23.*  
*Phalaena Fagi. Linn. Syst. Nat. 2. 816. 30. Fn.*  
*Sv. 1113.*

*Albin. Inf. tab. 58.*

*Aët. Holmenf. 1749. 132. tab. 4. fig. 10. 14.*

*Roes. Inf. 3. tab. 12.*

Habitat in Fago, Corylo, Betula.

*Larua* brunnea, dorso dentato, pedibus sex anterioribus elongatis, cauda reflexa corniculis duobus. Quiescit capite caudaque eleuatis, pedibus anticis pendentibus.

*Puppa* folliculata, brunnea.

B. alis reuerfis ferrugineis, striga flaua punctoque al- *Quercus.*  
bo. *Syst. Ent. 562. 24.*

*Phalaena Quercus. Linn. Syst. Nat. 2. 814. 25. Fn.*  
*Sv. 1106. Geoff. Inf. 2. 111. 13.*

*Phalaena* maxima fulua, alarum exteriorum superioritate intensius colorata, cum macula in medio alba, inferiore dilutiore. *Raj. Inf. 142. 2.*

*Merian. Europ. 1. tab. 10.*

*Petiv. Gazoph. tab. 45. fig. 5.*

*Albin. Inf. tab. 18. fig. 25.*

*Roes. Inf. 1. phal. 2. tab. 35.*

*Goed. Inf. 1. 51. tab. 7.*

*Reaum. Inf. 1. tab. 35.*

*Wilk. pap. 22. tab. 3. a. 11. 12.*

*Ammiral. Inf. tab. 31.*

*Schaeff. Icon. tab. 82. fig. 1 — 3.*

Habitat in Quercu, Salice.

*Larua* pilosa, grisea, nigro annulata alboque maculata.

*Puppa*



176 CLASS. VI. GLOSSATA. Bombyx.

*Puppa* dense folliculata, viridis antice fusca.  
Mas alis brunneis, femina pallidioribus.

*figma.* 40. B. alis reuerfis testaceis fusco irroratis punctoque centrali niueo. *Syst. Ent.* 563. 25.\*  
Habitat in America meridionali. *Mus. D. Hunter.*

*Pruni.* 41. B. alis reuerfis dentatis flauis, strigis duabus fusculis punctoque albo. *Syst. Ent.* 563. 26.\*  
*Phalaena Pruni.* *Linn. Syst. Nat.* 2. 813. 22.  
*Roes. Inf.* 1. phal. 2. tab. 36.  
Habitat in Pruno.

*Larua* laevis, cinerea, coerulesco lineata, collo lateribusque fasciculato pilosis, pedibus posticis portecti distantibus.

*Puppa* folliculata, antice nigra, postice brunnea.

*Amphimone.* 42. B. alis subreuerfis integris cinerascens, strigis duabus nigris, anticis puncto medio fuluo. *Syst. Ent.* 563. 27.\*  
Habitat in Terra del Fuego. *Mus. Dom. Banks.*

*potatoria.* 43. B. alis reuerfis subdentatis flauis, striga fulva repandaque, punctis duobus albis. *Syst. Ent.* 564. 28.

*Phalaena potatoria.* *Linn. Syst. Nat.* 2. 813. 23.  
*Phalaena maxima* alis e fuluo flauicantibus. *Raj. Inf.* 143. 3.

*Goed. Inf.* 1. tab. 12.

*Albin. Inf.* 1. tab. 17.

*Merian. Europ.* 27. tab. 16.

*Wilk. pap.* 27. tab. 3. b. 2.

*Sepp. Inf.* 4. 37. tab. 8.

*Roes. Inf.* 1. phal. 2. tab. 11.

*Schaeff. Icon.* tab. 67. fig. 10. 11.

Habitat in Europae graminibus.

*Larva* caudata, cristata pilosa, lateribus albo maculatis.

*Puppa* folliculata, brunnea.

*Oua* oblonga, plumbea, vtrinq̃ annulo punctoque medio viridibus.

CLASS. VI. GLOSSATA. Bombyx. 177

4. B. alis subreuerſis albis, punctis ocellaribus nigris *ocularia*.  
numerofis. *Syſt. Ent.* 564. 29.\*  
Habitat in America. *Muf. D. Hunter.*
  
5. B. alis ſubreuerſis concoloribus luteis, anticis ſtrigis *Hibiſci*.  
duabus, poſticis vnica. *Syſt. Ent.* 564. 30.\*  
Habitat in Hibifco populneo Indiae orientalis. *Muf.*  
*Tottianum.*
  
6. B. alis reuerſis? luteis, ſtrigis duabus annulisque tri- *Cynira*.  
bus anticarum connatis fufcis.  
*Phalaena Cynira. Cram. Inf.* 13. tab. 152. fig. C.  
Habitat Surinami.  
Magna. Alae omnes flauae ſtrigis duabus fufcis. An-  
nuli tres connati fuſci in centro alae anticae.
  
7. B. alis reuerſis luteis, ſtrigis duabus punctoque me- *Ceraſt.*  
dio fuſcis poſticoque albo. *Syſt. Ent.* 564. 31.\*  
Habitat in Pruno Ceraſo Angliae.
  
8. B. alis reuerſis grifeis, fascia ferruginea punctoque *Pini*.  
triangulari alba. *Syſt. Ent.* 565. 32.  
*Phalaena Pini. Linn. Syſt. Nat.* 2. 814. 24. *Fa. Sv.* 1104.  
*Merian. Europ.* tab. 22.  
*Friſch. Inf.* 10. tab. 10.  
*Wilk. pap.* 29. tab. 3. 6. 5.  
*Roef. Inf.* 1. phal. 2. tab. 59.  
*Schaeff. Icon.* tab. 86. fig. 1—3.  
Habitat in Europae Pino.  
*Larua* ſubcaudata, albo, grifeo fuſcoque variegata, col-  
laribus coeruleis punctis vtrinque ruſis.  
*Puppa* folliculata, fuſca.
  
9. B. alis reuerſis fuſcis, anticis puncto fascia margine- *dumeti*.  
que poſtico luteis. *Syſt. Ent.* 565. 33.  
*Phalaena dumeti. Linn. Syſt. Nat.* 2. 815. 26. *Fa.*  
*Sv.* 1107.  
*Petiv. Gazoph.* tab. 45. fig. 13.  
*Naturf.* 6. tab. 3. fig. 1—4.  
*Sulz. Hiſt. Inf.* tab. 21. fig. 3.  
Habitat in Europae Lactucae aliisque polyphaga.  
*Larua* fuſca flauo ruſoque maculata.  
*Puppa* brunnea, cauda dentata.  
*Fabr. Spec. Inf. T. II.* M 30. B.

178 CLASS. VI. GLOSSATA. Bombyx.

*versicolora*. 50. B. alis reuerfis griseis, strigis nigro albis, thorace antice albo. *Syst. Ent.* 565. 34.

*Phalaena versicolora*. *Linn. Syst. Nat.* 2. 817. 32. *Fn. Sv.* 1111.

*Phalaena* alis lineis albis et nigris vndatis. *Gadd. tag.* 82.

*Roef. Inf.* 3. tab. 39. fig. 3.

*Wilk. pap.* 45. tab. 1. a. 1.

*Sulz. Hist. Inf.* tab. 21. fig. 4.

Habitat in Europae Alno, Carpino, Betula.

*Larva* subcaudata, virefcens, flauo punctata lineisque obliquis flauis. *Fuesl. Magaz.* 2. tab. 1. fig. 4.

*Rubi*. 51. B. alis reuerfis ceruinis, strigis duabus albidis, fuscus nullis. *Syst. Ent.* 565. 35.

*Phalaena Rubi*. *Linn. Syst. Nat.* 2. 813. 21. *Fn. Sv.* 1112.

*Wilk. pap.* 25. tab. 3. a. 19.

*Ammiral. Inf.* tab. 32.

*Roef. Inf.* 3. tab. 49.

Habitat in Rubo, Salice.

*Larva* pilosa, subtus nigra, supra ferruginea, annulis nigris. Iunior velutina nigra annulis lacte luteis.

*Puppa* folliculata, nigricans, annulis tribus luteis.

*Vinula*. 52. B. alis subreuerfis, fusco venosis striatisque, corpore albo nigro punctato. *Syst. Ent.* 566. 36.

*Phalaena Vinula*. *Linn. Syst. Nat.* 2. 815. 29. *Fn. Sv.* 1112.

*Phalaena* pectinicornis elinguis, alis deflexis albidis diaphanis, vasis obscuris. *Geoff. Inf.* 2. 104. 5.

*Phalaena* alis amplis, exterioribus cinereis, maculis et lineis nigris eleganter depictis. *Raj. Inf.* 153. 5.

*Goed. Inf.* 1. tab. 65.

— 2. tab. 37.

*Merian. Europ.* tab. 39. fig. 140.

*Alb. Inf.* 11. tab. 5.

*Frisch. Inf.* 6. tab. 8.

*Roef. Inf.* 1. phal. 2. tab. 19.

*Reaum. Inf.* 2. tab. 21.

*Degeer Inf.* 1. tab. 23. fig. 12.

*Sepp. Inf.* 4. tab. 5.

*Wilk. pap.* tab. 19. fig. 1. c. 1.

Hab.

**CLASS. VI. GLOSSATA. Bombyx. 179**

Habitat in Salice, Populo.

*Larva* solitaria, gibba, viridis, dorso fusco, cauda bifeta, e rima sub capite humorem acrem espuit.

*Eruca* bicauda elegantissima Vinula dicta. *Raj. Inf.*

153. 3.

*Puppa* folliculata, fusca.

3. B. alis reuerfis ferrugineis, striga alba, anticis pun- *laneftris.*  
cto basique albis. *Syst. Ent.* 566. 37.

*Phalaena laneftris.* Linn. *Syst. Nat.* 2. 815. 28. *Fn.*

*Sv.* 1105.

*Roef. Inf.* 1. phal. 2. tab. 62.

*Schaeff. Icon.* tab. 88. fig. 10. 11.

*Wilk. pap.* 25. tab. 3. a. 18.

Habitat in Tilia, Pruno spinosa, Salice.

*Larva* gregaria, pilosa, nigra, segmento singulo punctis tribus albis inter maculas duas rubras penicillatas.

*Puppa* folliculata, sulphurea.

*Oua* lana densa cinerea tecta.

4. B. fusca antice pallida, alis reuerfis fuscescentibus, stri- *Populi.*  
ga selqui altera repanda albida. *Syst. Ent.* 566. 38.

*Phalaena Populi.* Linn. *Syst. Nat.* 2. 818. 34. *Fn.*

*Sv.* 1101.

*Wilk. pap.* 23. tab. 3. a. 13.

*Roef. Inf.* 1. phal. 2. tab. 60.

Habitat in Populo, Pomonae arboribus.

*Larva* cinerea, dorso obscuriore, segmentis singulis punctis duorum parium rubris.

*Puppa* folliculata antice fusca, postice rufa.

55. B. alis reuerfis ferrugineis unicoloribus, puncto albo. *Catax.*  
*Syst. Ent.* 567. 39.

*Phalaena Catax.* Linn. *Syst. Nat.* 2. 815. 27. *Fn.*

*Sv.* 1108.

*Roef. Inf.* 4. tab. 34. fig. 2. b.

— — 3. tab. 71. fig. 2. 1. a. 2. a. 3.

Habitat in Europae Quercu.

*Larva* cinerea, lineis duabus dorsalibus nigris rufo punctatis.

*Puppa* arcte folliculata, antice brunnea postice fusca.

M 2

56. B.

180 CLASS. VI. GLOSSATA. Bombyx.

*processionea*. 56. B. alis reuerfis fuscescentibus, striga obscuriore. *Syst. Ent.* 567. 40.

*Phalaena processionea*. *Linn. Syst. Nat.* 2. 819. 37.

*Reaum. Inf.* 2. tab. 11.

Habitat in Quercu.

*Larua* gregaria pilosa, fusco cinerea, dorso nigricanti, verrucis luteis. Incedunt per paria numero inaequalia. Exuviae tactu inflammationes excitant.

*Mori*. 57. B. alis reuerfis pallidis, strigis tribus obsoletis fuscis. *Syst. Ent.* 567. 41.

*Phalaena Mori*. *Linn. Syst. Nat.* 2. 817. 33. *Amon. acad.* 4. 563.

*Phalaena* pectinicornis elinguis, tota alba, alis deflexis, Bombyx dicta. *Geoff. Inf.* 2. 116. 18.

*Merian. Europ.* 1. tab. 1.

*Blank. Inf.* 60. tab. 9.

*Albin. Inf.* tab. 12. fig. 16.

*Bradt. nat.* tab. 27. fig. 1.

*Reaum. Inf.* 2. tab. 5. fig. 2.

*Roef. Inf.* 3. tab. 7. 8.

*Ammir. Inf.* tab. 9.

*Goed. Inf.* 112. tab. 42.

Habitat in moro Chinae, Persiae. Hospitatur in Europa australiori, tempore Iustiniario introducta.

*Larua* caudata, nuda, albida.

*Puppa* dense folliculata, brunnea.

*neustria*. 58. B. alis reuerfis griseis, strigis duabus ferrugineis, subtus vnica. *Syst. Ent.* 567. 42.

*Phalaena neustria*. *Linn. Syst. Nat.* 2. 818. 35.

*Phalaena* pectinicornis elinguis, alis deflexis pallidis, fascia alarum transversali saturatiore. *Geoff. Inf.* 2. 114. 16.

*Phalaena* media tota cinerea. *Raj. Inf.* 214. 8.

*Alb. Inf.* 19. fig. 27.

*Frisch. Inf.* 1. tab. 2.

*Reaum. Inf.* 2. tab. 4. fig. 1—11.

*Roef. Inf.* 1. phal. 2. tab. 6.

*Goed. Inf.* 1. 57. tab. 10.

*Wilk. pap.* 21. tab. 3. a. 10.

Habitat in Pomonae arboribus polyphaga.

*Larua*

CLASS. VI. GLOSSATA. Bombyx. 181

*Larva* gregaria, glauca, lineis vtrinque tribus rubris dorsaliq[ue] alba.

*Puppa* folliculata, fusca.

2. B. alis reuerfis obscuris, fasciis duabus pallidis. *Syst. castrensis*.  
Ent. 568. 43.

*Phalaena* castrensis elinguis, alis reuerfis griseis, strigis duabus pallidis, subtus vnica. *Linn. Syst. Nat.*

2. 818. 36.

*Frisch. Inf.* 10. tab. 8.

*Degeer Inf.* 1. tab. 13. fig. 4—6.

*Roef. Inf.* 4. tab. 14.

Habitat in Alchemilla, Millefolio polyphaga.

*Larva* gregaria, pilosa coerulea, lineis rubris nigro maculatis.

*Puppa* folliculata, obscura.

0. B. alis subreuerfis atro fuliginosis. *Syst. Ent. Hieracii*.  
568. 44.

*Phalaena* atra elinguis tota atra. *Linn. Syst. Nat.* 2.

823. 49.

Habitat in Hieracio, Delphinio.

*Larva* pilosa atra, linea dorsali sanguinea.

*Puppa* folliculata ferrugineo fusca.

\* \* \* alis deflexis.

1. B. alis deflexis flauescentibus, atomis strigisque duabus fuscis, pedibus anticis porrectis hirsutissimis. *lagopus*.

*Syst. Ent.* 568. 45.\*

Habitat in China. *Mus. D. Monson*.

1. B. alis flauis fusco maculatis, omnibus macula sub-*imperia-*  
ocellari ferruginea. *Syst. Ent.* 569. 46.\* *lis.*

*Phalaena imperialis*. *Drury Inf.* 1. tab. 9. fig. 1. 2.

Habitat in India.

3. B. alis deflexis cinereis, atomis strigisque vndatis fuscis. *crassicornis*.  
*Syst. Ent.* 569. 47.\*

Habitat in India. *D. Drury*.

14. B. alis deflexis coeruleis, anticis flauo maculatis. *Hyphinoe*.  
*Phalaena Hyphinos*. *Cram. Inf.* 13. tab. 154. fig. B. *noe*,

Habitat in Amboina.

Magna. Corpus obscure coeruleum. Alae anticae coeruleae maculis margineque postico flauis. Posticae coeruleae margine tenuissime flauo.

*Cossus*. 65. B. alis deflexis nebulosis, thorace postice fascia atra, antennis lamellatis. *Syst. Ent.* 569. 48.

*Phalaena Cossus*. Linn. *Syst. Nat.* 2. 827. 63. Fn. Sv. 1114.

*Phalaena pectinicornis* elinguis, alis albo cinereis, striis transuersis nebulosis nigris, abdomine annulis albis. *Geoff. Inf.* 2. 102. 4.

*Phalaena grandis*, alis cinerascens, lineolis creberrimis nigris. *Raj. Inf.* 150. 2.

*Phalaena* antennis filiformibus squamatis, elinguis, dorso cristato nigro lineato collarique albo alis cinereis, strigis numerosis nigris. *Degeer Inf. Vers. Germ.* 2. 1. 268. 1.

*Gronov. Zooph.* 204. 837.

*Goed. Inf.* 2. tab. 33.

*Petiv. Gazoph.* tab. 51. fig. 9.

*Merian. Europ.* tab. 36.

*Alb. Inf.* tab. 35. fig. 56.

*Frisch. Inf.* 7. tab. 1.

*Roef. Inf.* 1. phal. 2. tab. 18.

*Reaum. Inf.* 1. tab. 17. fig. 1—5.

*Lyonett. Less.* tab. 1. fig. 17—22.

*Wilks pap.* 15. tab. 2. a. 2.

*Schaeff. Icon.* tab. 61. fig. 1. 2.

Habitat in ligno putrescente.

*Larva* subpilosa. incarnata, dorso sanguineo, capite nigro. Romanis in deliciis fuit teste Plinio. vid. *Lyonett. Monogr.*

*Eruca* maxima subterranea, raris pilis obsita, supina parte tota, excepto capite rubra, prona flaua. *Raj. Inf.* 351. 25.

*Puppa* folliculata antice brunnea postice flaua. Forte potius ad *Hepialos* amandanda, quibus et laruae.

*dispar*. 66. B. alis deflexis, masculis griseo fuscoque nebulosis, femineis albidis, lituris nigris. *Syst. Ent.* 570. 49.

*Phalaena dispar*. Linn. *Syst. Nat.* 2. 821. 44.

*Pha-*

*Phalaena pectinicornis* elinguis, alis deflexis albis fascia quadruplici nigra acute undulata. *Geoff. Inf.* 2. 112. 14.

*Phalaena elinguis*, antennis pectinatis, femina alis albis nigro maculatis, mare alis ferrugineis, strigis nigris undatis. *Degeer Inf. Vers. Germ.* 2. 1. 209. 3.

*Merian. Europ.* 1. tab. 18.

*Frisch. Inf.* 1. 14. tab. 3.

*Reaum. Inf.* 2. tab. 1. fig. 11-14.

*Wilk. pap.* 20. tab. 3. a. 7.

*Roes. Inf.* 1. phal. 2. tab. 3.

*Schaeff. Icon.* tab. 28. fig. 3—6.

Habitat in Pomonae arboribus polyphaga, hortorum pestis.

*Larva* pilosa albo lineata, punctis antice coeruleis, posticis rubris. Manus pruritu inficit.

*Puppa* folliculata, punctis quatuor anticis nigris. Tacta in gyrum se voluit.

*Mas* alis fusco griseis nebulosis.

*Femina* duplo maior alis albis, lituris nigris, ova pilis pulverulentis obducit.

67. B. alis deflexis? albidis, strigis nigris, posticis flavis nigro maculatis, abdomine atro, cingulis rubris. *Amasis.*

*Phalaena Amasis. Cram. Inf.* 18. tab. 206. fig. B.

Habitat Surinami.

Caput et thorax alba nigro maculata. Abdomen atrum cingulis quinque sanguineis anoque flavo. Alae anticae albae strigis tribus angulatis nigris.

68. B. alis deflexis cinereis, strigis tribus undatis fuscis. *pudibunda. Syst. Ent.* 570. 50.

*Phalaena pudibunda. Linn. Syst. Nat.* 2. 824. 54. *Fn. Sv.* 1118.

*Phalaena pectinicornis*, elinguis, alis deflexis cinereo undulatis, fasciis transversis obscurioribus, capite inter pedes porrectos. *Geoff. Inf.* 2. 113. 15.

*Phalaena cinerea*, alis oblongis, exterioribus quatuor lineis nigricantibus transversis, distinctis. *Raj. Inf.* 185. 7.



*Goed. Inf.* 3. tab. 5.

*Merian. Europ.* 1. tab. 47.

*Alb. Inf.* tab. 26.

*Rotum. Inf.* 1. tab. 33. fig. 4 — 12.

*Roef. Inf.* 1. phal. 2. tab. 38.

*Degeer Inf.* 1. tab. 16. fig. 11. 12. ]

*Wilk. pap.* 30. tab. 3. c. 2.

*Ammiral.* tab. 18.

*Schaeff. Icon.* tab. 44. fig. 9. 10.

— — — tab. 90. fig. 1. 2. 3.

Habitat in Pomonae arboribus, Corylo polyphaga.

*Larva* fasciculata pilosa, flava, capite luteo, caudæ fasciculo longiori rubro, dorsii quatuor albis.

*Eruca* maior pulcherrima pilosa e viridi flavicans.

*Raj. Inf.* 344. 9.

*Puppa* folliculata, antice nigra, postice brunnea flavo maculata.

*fascelina.* 69. B. alis deflexis cinereis, atomis nigris strigisque duabus fulvis repandis. *Syst. Ent.* 571. 51.

*Phalaena fascelina* elinguis cristata cinerea, alis superioribus antice fasciis duabus angustis fulvo fuscis, scutello bipunctato fulvo. *Linn. Syst. Nat.* 2. 824. 55. *Fn. Sv.* 1119.

*Phalaena* obsoleta cinerea. *Raj. Inf.* 186. 8.

*Goed. Inf.* 1. tab. 36.

*Merian. Europ.* tab. 8.

*Degeer Inf.* 1. tab. 15. fig. 15.

*Roef. Inf.* 1. phal. 2. tab. 37 mal.

*Wilk. pap.* 30. tab. 3. c. 1.

Habitat in Leontodo, Trifolio, Rubo, Populo,

*Larva* pilosa, fasciculata, rubro punctata, fasciculis quinque dorsalibus albis, capitis anique fuscis.

*Puppa* folliculata brunnea, linea dorsali pilosa.

*bucephala.* 70. B. alis deflexis cinereis, strigis duabus ferrugineis maculaque terminali flava. *Syst. Ent.* 571. 52.

*Phalaena bucephala.* *Linn. Syst. Nat.* 2. 816. 31. *Fn. Sv.* 1115.

**CLASS. VI. GLOSSATA. Bombyx. 183.**

*Phalaena pectinicornis* elinguis, alis tubiformibus, superioribus cinereis, fascia duplici ferruginea et extremo circulariter pallescente, subtus omnibus flavescentibus, fascia undulata fusca. *Geoff. Inf.* 2, 123. 28.

*Phalaena media*, alis oblongis ex rufo cinereo fulvo et albicante variis. *Raj. Inf.* 162. 14.

*Goed. Inf.* 1. tab. 34.

*Merian. Europ.* 3. tab. 41.

*Alb. Inf.* tab. 23. fig. 33.

*Frisch. Inf.* 11. tab. 4.

*Degeer Inf.* 1. tab. 13. fig. 14 — 19.

*Roes. Inf.* 1. phal. 2. tab. 14.

*Wilk. pap.* 21. tab. 3. a. 8.

*Schaeff. Icon.* tab. 31. fig. 10. 11.

Habitat in Tilia, Quercu, Pomonae arboribus.

*Larva* pilosa nigra annulis lineisque luteis punctis albis lateralibus.

*Puppa* nuda postice corniculis duobus.

B. alis deflexis nebulosis, posticis brunneis, abdomine *Helops*. brunneo albo annulato.

*Phalaena Helops. Cram. Inf.* 6. tab. 72. fig. C.

Habitat in America.

Magnitudo praecedentis. Alae anticae fuscae ferrugineo nebulosae, posticae brunneae, immaculatae. Corpus brunneum abdomine annulis atris.

B. alis deflexis griseis, fasciis duabus ferrugineis *coeruleo-maculaeque* albida duplicato didyma. *Syst. Ent. cephal.* 572. 53.

*Phalaena coeruleocephala* elinguis cristata, alis deflexis griseis, stigmatibus albidis coadunatis. *Linn. Syst. Nat.* 2. 326. 59. *Fn. Sv.* 1117.

*Phalaena pectinicornis* elinguis, alis deflexis fuscis, macula duplici albo flavescente, geminata. *Geoff. Inf.* 2. 122. 27.

*Phalaena* alis exterioribus pullis duabus tribusque maculis albis e duobus circulis compositis contiguis notatis. *Raj. Inf.* 163. 17.

186 CLASS VI. GLOSSATA. Bombyx.

*Goed. Inf.* 1. tab. 61.

*Merian. Europ.* tab. 9.

*Albin. Inf.* tab. 13. fig. 17.

*Frisch. Inf.* 10. tab. 3. fig. 4.

*Reaum. Inf.* 1. tab. 18. fig. 6—9.

*Roes. Inf.* 1. phal. 2. tab. 16.

*Wilks pap.* 6. tab. 1. a. 12.

Habitat in Pomonae arboribus.

*Larva* nuda coerulescens luteo lineata nigro punctata.

*Eruca* rarius pilosa mali, pyri. *Raj. Inf.* 163. 17.

*Puppa* folliculata, brunnea.

Quies sex hebdomadam.

*argenti-* 73. B. alis deflexis dentatis griseis argenteo maculatis.

*na. Phalaena argentina. Wien. Verz.* tab. 1. a. fig. 2.

— — — tab. 1. b. fig. 2.

Habitat in Germaniae Quercu.

*Larva* nuda grisea, segmentis 4. 11. 12. tuberculatis.

*Puppa* obtusa, brunnea.

*Imago* dorso fasciculis tribus pilorum erectorum.

*decora.* 74. B. alis deflexis, anticis rubro, flauo nigroque variatis, posticis rubris, margine nigro.

*Phalaena decora. Linn. Syst. Nat.* 2. 1068. 14. *Mus.*

*Lud. Vlr.* 174.

Habitat ad Promontorium bon. sp.

Alae anticae maculis quatuor fasciaque postica abbreviata rufis. Margo posticus niger maculis septem flauis.

*Dione.* 75. B. alis deflexis albis nigro striatis punctatisque, subten margine purpureo. *Syst. Ent.* 572. 54. \*

Habitat in America. *Mus. Dom. Monson.*

*Ziczac.* 76. B. alis deflexis dorso dentatis apicibusque macula grisea subocellari, antennis squamatis. *Syst. Ent.* 573. 55.

*Phalaena Ziczac. Linn. Syst. Nat.* 2. 827. 61. *Fa. Sv.* 1116.

*Phalaena pectinicornis* elinguis, alis exterioribus fasciis, venis plurimis, fascia circulari et marginis interioris

CLASS. VI. GLOSSATA. Bombyx. 187.

terioris appendice nigricantibus, inferioribus nigricantibus, limbo lineari fusco. *Geoff. Inf.* 2. 124. 29.

*Goed. Inf.* 3. tab. E.

*Merian. Europ.* tab. 147.

*Albin. Inf.* tab. 14. fig. 20.

*Frisch. Inf.* 3. tab. 1. fig. 2.

*Reaum. Inf.* 2. tab. 22. fig. 9 — 16.

*Roef. Inf.* 1. phal. 2. tab. 20.

*Degeer Inf.* 1. tab. 6. fig. 1 — 10.

*Wilks pap.* 12. tab. 1. d. 1.

*Schaeff. Icon.* tab. 69. fig. 2. 3.

abitat in Salice.

*larva* solitaria nuda, dorso gibberibus duobus, cauda rufa.

*ovipara* folliculata, brunnea.

3. alis deflexis exustis, plaga albida, posticis albis. *distaca*, *Syst. Ent. App.* 831.

*halaena distaca*. *Linn. Syst. Nat.* 2. 826. 60.

abitat in Europae Populo nigra, Betula alba.

3. alis deflexis, anticis nebulosis dorso dentatis, litura Dromedarios anique flavescentibus. *Syst. Ent. Append. darius*. 831.

*halaena dromedarius*. *Linn. Syst. Nat.* 2. 827. 62.

*halaena* elinguis, antennis pectinatis, dorso dentato, alis anticis strigis vndatis fuscis maculaque bascos flava. *Degeer Inf. Vers. Germ.* 2. 1. 223. 2.

abitat in Europae Betula, Alno, Corylo, Populo.

scia postica alarum ferruginea.

*larva* solitaria nuda, brunnea, gibberibus quatuor dorsalibus laruae Ziczac simillima.

3. alis deflexis glaucis, fascia ferruginea, puncto nigro albo annulato, thorace variegato. *Syst. Ent. Coryli*. 573. 56.

*halaena Coryli* elinguis, thorace variegato, alis antice griseo nebulosis, postice coerulescenti glaucis, antennis flavis. *Linn. Syst. Nat.* 2. 823. 50. *Fu. Sv.* 1123.

*Alb. Inf.* tab. 90.

*Wilk.*

*Wilk. pap.* 31. tab. 3. c. 5.

*Degeer Inf.* 1. tab. 18. fig. 4. 5.

*Roef. Inf.* 1. phal. 2. tab. 58.

Habitat in Corylo, Betula.

*Larva* fascicularis pilosa, rubra, fasciculis dorsalibus duobus, caudae vnico, collaribus longioribus.

*Puppa* folliculata, antice nigra, postice brunnea.

*Alphaea.* 80. B. alis deflexis ferrugineis, puncto medio albo strigae punctata fusca. *Syst. Ent.* 573. 57. \*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

*monacha.* 81. B. alis deflexis albis, atro vndatis, abdominis incisuris sanguineis. *Syst. Ent.* 574. 58.

*Phalaena monacha.* Linn. *Syst. Nat.* 2. 821. 43. *Fn. Sv.* 1130.

*Phalaena media* ex cinereo albicante et nigro coloribus varia. *Raj. Inf.* 158. 7.

*Wilk. pap.* tab. 3. a. 4.

*Schaeff. Icon.* tab. 68. fig. 2. 3.

Habitat in Salice, Malo, Larice, Rubo.

*Larua* fusco cinerea, fasciculis dorsi rubris, segmento secundo macula cordata nigra.

*Eruca* Rubum idaeum depascens. *Raj. Inf.* 159. 8.

*Puppa* nigra.

*flaua.* 82. B. alis deflexis flauissimis, apice punctis tribus nigris. *Syst. Ent.* 574. 59. \*

Habitat in India. *D. Koenig.*

*lutea.* 83. B. alis deflexis flauissimis immaculatis. *Syst. Ent.* 574. 60. \*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

*curtula.* 84. B. alis deflexis glaucis, strigis albis maculaque apicis fusca. *Syst. Ent.* 474. 61.

*Phalaena curtula* elinguis, thorace ferruginato, alis deflexis glaucis, striga alba, apicibus macula testacea. Linn. *Syst. Nat.* 2. 823. 52. *Fn. Sv.* 1124.

*Phalaena elinguis*, antennis pectinatis, dorso macula ferruginea, alis glaucis, strigis quatuor albidis maculaque fusca. *Degeer Inf. Versf. Germ.* 2. 1. 234. 11. tab. 5. fig. 1.

*Sepp.*

*Sepp. Inf.* 4. tab. 1. fig. 13 — 18.

*Roef. Inf.* 3. tab. 43.

— — 4. tab. 11. fig. 1 — 6.

*Frisch. Inf.* 5. tab. 6.

Habitat in Salice, Populo.

*Larva* solitaria pilosa fusca, stria dorsali incarnata, verrucis duabus eleuatis punctisque lateralibus rufis.

*Puppa* folliculata, brunnea.

*Imago* quiescit capite inflexo pedibusque anticis porrectis.

Conf. *Phalaena* elinguis, antennis pectinatis, dorso cristata macula ferruginea, alis glaucis, strigis quatuor albis maculaque ferruginea. *Degeer Inf. Vers. Germ.*

2. 1. 231. 9. tab. 4. fig. 24. Vix differre videtur.

85. B. alis deflexis griseis, strigis tribus albidis subanostomomorphis, thorace ferruginato. *Syst. Ent.* 575. 62. *anostomomorphis*.

*Phalaena anostomosis*. Linn. *Syst. Nat.* 2. 824. 53. *Fn. Sv.* 1124.

*Phalaena* elinguis, antennis pectinatis, dorso macula ferruginea, alis griseis, strigis undatis cinereis. *Degeer Inf. Vers. Germ.* 2. 1. 233. 10.

*Goed. Inf.* 1. tab. 33.

*Roef. Inf.* 1. phal. 2. tab. 26.

Habitat in Salice.

*Larva* fusca dorso albo punctato lineaque laterali flaua rubro punctata, tubere humeri anique.

*Puppa* folliculata nigra striis duabus rubris.

*Imago* quiescit capite inflexo pedibusque anticis porrectis.

86. B. alis deflexis flauissimis, strigis duabus obliquis obsulphurea. scurioribus. *Gen. Inf. Mant.* 274. \*

Habitat in Germania. *Dom. de Hattorf.*

87. B. alis deflexis fuscis, maculis tribus viridibus. *Cippus.*

*Phalaena Cippus*. *Cram. Inf.* 5. tab. 53. fig. E.

Habitat Surinami.

Statura praecedentium. Corpus totum brunneum alis maculis tribus viridibus.

88. B. alis deflexis dentatis albidis, nigro venosis, palpis palpina. porrectis pennaceis. *Syst. Ent.* 575. 64.

*Pha-*

190 CLASS. VI. GLOSSATA. Bombyx.

*Phalaena palpina*. Linn. *Syst. Nat.* 2. 328. 64.

*Sepp. Inf.* 4. 17. tab. 4.

*Degeer Inf.* 1. tab. 4. fig. 7.

*Naturf.* 2. tab. 1. fig. 7. mal.

Habitat in Salice.

*Larua* solitaria nuda viridis, linea laterali flava.

*Gnoma*. 89. B. alis deflexis subdentatis cinereo albis, vitta marginali atra, macula alba. *Gen. Inf. Mant.* 274.\*

Habitat Hamburgi. *D. Schulz.*

*capucina*. 90. B. alis deflexis denticulatis ferrugineis, denticulo dorsali reflexo. *Syst. Ent.* 575. 65.

*Phalaena capucina*. Linn. *Syst. Nat.* 2. 332. 79.

Habitat in Europae plantis.

*camelina*. 91. B. alis deflexis denticulatis brunneis, omnibus denticulo dorsali. *Syst. Ent.* 575. 66.

*Phalaena camelina*. Linn. *Syst. Nat.* 2. 332. 80. 1 Sv. 1145.

*Roef. Inf.* 1. phal. 2. tab. 28.

*Reaum. Inf.* 2. tab. 20. fig. 13.

Habitat in Tilia, Alno, Betula.

*Larua* nuda virefcens, corniculis duobus caudalibus stigmatibus purpureis.

*Puppa* folliculata antice brunnea, postice fusca.

*aulica*. 92. B. alis deflexis anticis griseis flavo punctatis, postice fulvis nigro maculatis. *Syst. Ent.* 576. 67.

*Phalaena aulica*. Linn. *Syst. Nat.* 2. 329. 68. *Fa. S.* 1133.

*Clerk. Icon.* tab. 4. fig. 3.

*Merian. Eur.* 2. tab. 8.

*Albin. Inf.* tab. 22.

*Wilk. Inf.* 19. tab. 3. a. 3

*Naturf.* 4. tab. 1. fig. 8.

— — 6. tab. 5. fig. 3.

Habitat in Cynoglossio, Angelica, Vrtica.

*tubricipeda*. 93. B. alis deflexis albidis, punctis nigris, abdomineque quefariam nigro punctato. *Syst. Ent.* 576. 68.

*Phalaena tubricipeda*. Linn. *Syst. Nat.* 2. 329. 69. *A.* Sv. 1138.

*halaena pectinicornis* elinguis, alis deflexis albidis, punctis nigris, abdomine ordinibus quinque punctorum. *Geoff. Inf.* 2. 118. 21.

*Goed. Inf.* 1. tab. 23. 38.

*Merian. Eur.* 1. tab. 46. fig. 65.

*Albin. Inf.* tab. 24. fig. 36.

*Frisch. Inf.* 3. tab. 8.

*Reaum. Inf.* 2. tab. 1. fig. 7 — 9.

*Roes. Inf.* 1. phal. 2. tab. 46. 47.

*Degeer Inf.* 1. tab. 11. fig. 7. 8.

*Schaeff. Icon.* tab. 24. fig. 8. 9.

abitat in Quercu, Atriplice, Oleribus polyphaga. iriat alarum colore et punctorum numero.

*arua* pilosa fusca, punctis coeruleis lineaque dorsali pallida.

*uppa* folliculata coerulelescens, stigmatibus rubris.

B. alis deflexis albis, punctis nigris ramosis, abdomine immaculato. *leporina.*

*halaena leporina.* *Linn. Syst. Nat.* 2. 838. 109. *Fn. Sv.* 1176.

*Degeer Inf.* 1. tab. 12. fig. 10. 11. 17.

*Fuesl. Magaz.* 2. tab. 1. fig. 1.

abitat in Europae Salice, Betula.

finis praecedenti. Alae posticae niueae immaculae, nitidae.

*arua* pilosa, virescens, antice lineis tribus brunneis fasciculisque tribus atris.

*uppa* folliculata, brunnea.

B. alis deflexis cinereis, puncto disci atro posticeque striga purpurascens fracta. *lota.*

*halaena lota.* *Linn. Syst. Nat.* 2. 830. 70. *Fn. Sv.* 1137.

*Clerk. Icon.* tab. 8. fig. 1.

abitat in Europae Salice.

*arua* cinerea, lineis nigris albisque, punctisque albidis.

*uppa* folliculata, brunnea.

B. alis deflexis anticis niueis, fascia lata nigra, antennis filiformibus. *Syst. Ent.* 576. 69. \*

abitat in Sueciae nemoribus.



*compressa*. 97. B. alis compresso adscendentibus niueis, macula communi fusca, centrali grisea, lunula alba.

*Pyrallis Bankiana* alis albis macula communi fusca.  
*Syst. Ent.* 645. 1.

Habitat in Germania.

Parua in hoc genere. Antennae simplices fuscae. Alae valde compressae, postice adscendentes niueae, anticae macula magna communi fusca et in medio macula magna transversalis grisea lunula parua niuea. Versus marginem posticum lunulae fuscae et margo ipse fuscus. Alae posticae niueae. Subtus anticae griseae, posticae niueae.

*Milhauseri*. 98. B. alis deflexis canis, maculis duabus dorsalibus fuscis, antennis apice setaceis. *Syst. Ent.* 577. 70. \*

*Milhauseri Monogr. Dresd.* 1763. tab. 1.

Habitat Dresdae.

*Larua* viridis, spinis dorsalibus furcatis, pedibus subcaudalibus nullis.

*Puppa* folliculata brunnea antice acuminato emarginata.

*Lincus*. 99. B. alis deflexis atris, anticis apice posticis basi fulvis.  
*Phalaena Lincus. Cram. Inf.* 5. tab. 50. fig. 11.

Habitat Surinami.

Antennae filiformes, vix pectinatae. Margo interior alae anticae in acumen definit. Margo posticus alae posticae tenuissime flavus.

*Frigula*. 100. B. alis deflexis anticis griseis, apice albo nigroque striatis. *Syst. Ent.* 577. 7. 2. \*

Habitat Dresdae Saxonum.

*V-nigrum*. 101. B. alis deflexis albis, V nigro notatis. *Syst. Ent.* 577. 73. \*

*Naturf.* 2. tab. 1. fig. 1. mal.

Habitat in Saxoniae Tilia.

*Larua* fasciculata, fasciculis dorsalibus octo, anticis posticisque albis, collari hirsutissimo.

*Naturf.* 12. tab. 1. fig. 8. 9. 10.

*Puppa* viridis, macula thoracis nigra.

B. alis deflexis niueis, ano barbato ferrugineo. *Syst. chryso-*  
*Ent.* 577. 74. *rhoea.*

*halaena chrysoorrhoea.* Linn. *Syst. Nat.* 2. 822. 45.  
*Fn. Sv.* 1128.

*Geoff. Inf.* 2. 117. 20.

*Raj. Inf.* 156. 1.

*Merian. Eur.* 1. 8. tab. 20.

*Frisch. Inf.* 3. 28. tab. 8.

*Reaum. Inf.* 1. tab. 16. fig. 11.

*Roef. Inf.* 1. phal. 2. tab. 22.

*Wilk. pap.* 28. tab. 3. b. 3.

*Edw. Av.* 122. tab. 122.

abitat in Pomonae arboribus polyphaga.

*arma* gregaria, pilosa nigricans, lineis duabus dor-  
salibus rubris, lateribus albo fasciculatis.

*ruca* media pilosa, humeris in gibbum elatis, albo  
nigro rubroque coloribus varia. *Raj. Inf.* 343. 5.

*ypa* folliculata, nigricans.

*ua* lutea lana copiosa fulva tegit.

B. alis deflexis albis, pedibus nigris albo annulatis. *Salicis.*  
*Syst. Ent.* 578. 75.

*halaena Salicis.* Linn. *Syst. Nat.* 2. 822. 46. *Fn. Sv.*  
1129.

*halaena* pectinicornis elinguis, alis deflexis albis pe-  
dum annulis antennisque nigris. *Geoff. Inf.* 2. 116. 19.

*Goed. Inf.* 1. tab. 3.

*Blank. Inf.* tab. 8. fig. A. D.

*Merian. Europ.* 1. tab. 30.

*Albin. Inf.* tab. 84. fig. a — d.

*Frisch. Inf.* 1. tab. 4.

*Reaum. Inf.* 1. tab. 34. fig. 4 — 6.

*Harris Aurel.* 2. tab. 5. fig. k. p.

*Roef. Inf.* 1. phal. 2. tab. 9.

*Degeer Inf.* 1. tab. 11. fig. 13. 14.

*Wilk. pap.* 21. tab. 3. a. 9.

abitat in Salice, Populo.

*irua* pilosa nigra, dorso longitudinaliter albo macu-  
lato rubro punctato.

*ypa* folliculata, pilosa, brunnea.

vies 30 dierum.

*abr. Sp. Inf. T. II.*

N

104. B.

194 CLASS. VI. GLOSSATA. Bombyx.

*Crataegi.* 104. B. alis deflexis rotundatis cinereis, fascia obscurior, ano barbato.

*Phalaena Crataegi.* Linn. *Syst. Nat.* 2. 823. 48. *F. Sv.* 1126.

*Reaum. Inf.* 1. tab. 44. fig. 10.

*Degeer Inf.* 1. tab. 11. fig. 20. 21.

— — *Vers. Germ.* 2. 1. 214. 3.

Habitat in Crataego Oxyacantha.

*Larua* pilosa nigra fasciis albis tuberculisque quatuor ferrugineis.

*Eridanus.* 105. B. alis deflexis niuea, abdomine fuluo annulato.

*Phalaena Eridanus.* *Cram. Inf.* 6. tab. 68. fig. G.

Habitat Surinami.

Antennae filiformes. Huius tamen videtur generis.

*tibialis.* 106. B. alis deflexis niueis, tibiis anticis flavescentibus nigro punctatis. *Syst. Ent.* 578. 76.\*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

*pholeta.* 107. B. alis deflexis albidis, costa palpisque ferrugineis. *Syst. Ent.* 579. 77.\*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

*Corones.* 108. B. alis deflexis, anticis cinereis, posticis niueis. *Syst. Ent.* 579. 78.\*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

*festiua.* 109. B. alis deflexis flavescentibus, basi coeruleo maculatis, apice nigro punctatis. *Syst. Ent.* 579. 79.\*

Habitat in America. *Mus. D. Hunter.*

*Dryas.* 110. B. alis deflexis fuscis, abdomine fuluo, punctis utroque nigris.

*Phalaena Dryas.* *Cram. Inf.* 6. tab. 70. fig. 6.

Habitat Surinami.

Alae posticae basi cyaneae. Abdomen fuluum striis nigris punctorum nigrorum.

*Russula.* 111. B. alis deflexis luteis, margine sanguineo lunulata fusca, posticis subtus immaculatis. *Syst. Ent.* 579. 80.\*

*Phalaena Russula.* Linn. *Syst. Nat.* 2. 830. 71.

*halaena Sannio. Linn. Fn. Sv. 1135.*

*halaena pectinicornis spirilinguis, alis deflexis pallido luteis, limbo roseo, superioribus macula, inferioribus fascia duplici fusca. Geoff. Inf. 2. 129. 39.*

*Robert. Icon. tab. 30. fig. 1.*

*Raj. Inf. 228. 75.*

*Schaeff. Icon. tab. 83. fig. 4. 5.*

*Phalaena Ruffula. Linn. Fn. Sv. 1156.*

*Clerk. Icon. tab. 4. fig. 1.*

*Kleman. Inf. 1. tab. 20.*

habitat in Lactuca, Scabiosa.

B. alis incumbentibus fuscis, linea punctisque duobus *Iacobaeae* rubris, posticis rubris nigro marginatis. *Syst. Ent. 588. 113.*

*halaena Iacobaeae. Linn. Syst. Nat. 2. 839. 111. Fn. Sv. 1155.*

*halaena media, alis exterioribus colore nigro et sanguineo varlis, extimo duntaxat margine nigro. Raj. Inf. 168. 26.*

*Geoff. Inf. 2. 146. 75.*

*Robert. Icon. 1. tab. 20.*

*Blank. Inf. tab. 1. fig. G. K.*

*Petiv. Gazoph. tab. 33. fig. 6.*

*Merian. Eur. 3. tab. 28.*

*Albin. Inf. tab. 34. fig. G. H.*

*Harr. Aurel. tab. 4.*

*Goed. Inf. 1. 56. tab. 9.*

*Raum. Inf. 1. tab. 16. fig. 4 — 7.*

*Roef. Inf. Phal. 2. tab. 49.*

*Wilk. pap. 26. tab. 3. a. 1.*

*Ammir. tab. 3.*

*Edw. Av. tab. 271.*

*Schaeff. Elem. tab. 98. fig. 3.*

— — *Icon. tab. 47. fig. 2. 3.*

abitat in Senecione Iacobaea.

*arua* subpilosa atra, flauo annulata.

*ruca* glabra ex nigro et flauo annulatim variegata.

*Raj. Inf. 169. 26.*

*uppa* tenuissime folliculata, testacea.

*grammica*. 113. B. alis deflexis luteis, anticis flavis nigro striatis, posticis fascia terminali nigra. *Syst. Ent.* 579. 81.  
*Phalaena grammica*. *Linn. Syst. Nat.* 2. 831. 75. *Fn. Sv.* 1134.

*Geoff. Inf.* 2. 115. 17.

*Linn. Amoen. acad.* 5. tab. 3. fig. 31.

*Merian. Eur.* 1. tab. 5.

*Schaeff. Icon.* tab. 92. 2.

*Roef. Inf.* 4. tab. 21. fig. A. D.

Habitat in Plantagine.

*Larva* fusca, linea dorsali alba, pedibus rufis.

*Puppa* nuda, brunnea.

*purpurea*. 114. B. alis deflexis anticis flavis fusco punctatis, posticis rubris nigro maculatis. *Syst. Ent.* 580. 82.

*Phalaena purpurea*. *Linn. Syst. Nat.* 2. 828. 67.

*Geoff. Inf.* 2. 105. 6.

*Merian. Europ.* 1. tab. 6.

*Alb. Inf.* tab. 22.

*Seb. Mus.* 4. tab. 59. fig. 4. 5.

*Roef. Inf.* 1. phal. 2. tab. 10.

Habitat in Europae Ribes.

*Larva* hirsuta, grisea, albo maculata odorem Marti veri spargit.

*Puppa* nuda, brunnea.

*Plantaginis*. 115. B. alis deflexis atris, riualis albis, posticis luteis, margine maculisque nigris. *Syst. Ent.* 580. 83.

*Phalaena Plantaginis* elinguis, alis deflexis atris, riualis flavis, inferioribus rubro maculatis. *Linn. Syst. Nat.* 2. 820. 42. *Fn. Sv.* 1132.

*Phalaena pectinicornis* elinguis, alis deflexis, superioribus fulcis, maculis luteis, inferioribus rubra, maculis quatuor nigris. *Geoff. Inf.* 2. 109. 10.

*Phalaena alpicola*. *Scop. carn.* 507.

*Wilk. pap.* 24. tab. 3. a. 5.

*Roef. Inf.* 4. tab. 24.

*Schaeff. Icon.* tab. 92. fig. 5. 7.

Habitat in Plantagine, Vlmio.

Mas alis posticis luteis, femina coccineis.

*Larva* pilosa atra, dorso brunneo.

*Puppa* atra.

116. B. alis deflexis flauis riuulis punctisque atris, posticis fuscis. *Gen. Inf. Mant.* 280. \* *lugubris.*

*Phalaena sulphuralis.* Linn. *Syst. Nat.* 2. 881. 333.

*Schaeff. Icon.* tab. 9. fig. 14. 15. mal.

Habitat Hamburgi. *D. Schulz.*

117. B. alis deflexis fuscis, anticis cinereo subfasciatis, posticis macula bascos fasciaque rufis. *Syst. Ent.* 580. 84. \* *Vidua.*

*Phalaena Parthenias spirilinguis*, alis deflexis fusco alboque variis, inferioribus luteis punctis duobus nigris. Linn. *Syst. Nat.* 2. 835. 94. *Fn. Sv.* 1160. *It.*

*Westrogoth.* 141. 1.

Habitat in Anglia, Suecia.

118. B. alis deflexis atris, maculis octo albis, posticis flauis villica nigro maculatis. *Syst. Ent.* 2. 581. 85. *villica.*

*Phalaena villica.* Linn. *Syst. Nat.* 2. 820. 41.

*Geoff. Inf.* 2. 106. 1.

*Harris Aurel.* tab. 4.

*Raj. Inf.* 156. 4.

*Alb. Inf.* tab. 21.

*Frisch. Inf.* 10. tab. 2.

*Reaum. Inf.* 1. tab. 31. fig. 4 — 6.

*Roef. Inf.* 4. tab. 28. fig. 2.

— — — tab. 29. fig. 1 — 4.

*Wilk. pap.* tab. 3. a. 2.

Habitat in Oleribus, Vrtica, Millefolio, polyphaga.

*Larua* hirsuta nigricans, fuluo maculata, capite pedibusque rufis.

*Puppa* folliculata, nigra.

119. B. alis deflexis albis nigro fasciatis, posticis sanguineis nigro maculatis. *Syst. Ent.* 581. 86. *Hebe.*

*Phalaena Hebe* elinguis, alis deflexis atris, fasciis albis, inferioribus rubris, maculis nigris. Linn. *Syst. Nat.* 2. 820. 40.

*Phalaena* elinguis pestinicornis, alis deflexis, superioribus albidis riuulis transuersis nigris, inferioribus roseis, macula triplici nigra. *Geoff. Inf.* 2. 109. 9.

*Phalaena festiua.* Berl. *Magaz.* 2. 416. 32.

198 CLASS. VI. GLOSSATA. Bombyx

*Frisch. Inf.* 7. tab. 9.

*Roesl. Inf.* 4. tab. 27. fig. 1. 2.

*Kleman. Inf.* 1. tab. 13. fig. 1 — 4.

*Schaeff. Elem.* tab. 98. fig. 1.

Habitat in Artemisia, Euphorbia.

*Larva* solitaria, hirta, segmentis tribus anterioribus ferrugineis.

*Puppa* nigra.

**Tarquinus.** 120. B. alis deflexis cinereis, macula media atra, lineatum exferente alba. *Gen. Inf. Mant.* 280. \*

*Phalaena Tarquinus.* *Cram. Inf.* 1. tab. 4. fig. B. C.

Habitat Surinami.

**Tarquinia.** 121. B. alis deflexis, anticis nigris, linea hamata interstigas duas albas. *Gen. Inf. Mant.* 280. \*

*Phalaena Tarquinia.* *Cram. Inf.* 1. tab. 4. fig. A.

Habitat Surinami.

Feminam praecedentis credit *Cramer*, an. rite?

**Caia.** 122. B. alis deflexis fuscis, riuulis albis, posticis purpureis nigro punctatis. *Syst. Ent.* 581. 87.

*Phalaena Caia.* *Linn. Syst. Nat.* 2. 819. 38. *Fa. S.*

1131.

*Geoff. Inf.* 2. 108. 8.

*Raj. Inf.* 151. 3.

*Merian. Europ.* tab. 5.

*Albin. Inf.* tab. 20. fig. C. D.

*Frisch. Inf.* 2. tab. 9.

*Sepp. Inf.* 4. 9. tab. 2.

*Roesl. Inf.* 1. phal. 2. tab. 1.

*Goed. Inf.* 1. tab. 17.

*Reaum. Inf.* 1. tab. 36. fig. 6. 7.

*Degeer Inf.* 1. tab. 12. fig. 8. 9.

— — *Vers. Germ.* 2. 1. 215.

*Blanc. Inf.* tab. 9. fig. A. C.

*Wilk. pap.* 18. tab. 3. a. 1.

*Sulz. Inf.* tab. 16. fig. 94.

*Schaeff. Icon.* tab. 29. fig. 7. 8.

Habitat in Oleribus polyphaga.

*Larva* solitaria, hirta nigricans, segmentis vtrinque punctis tribus eleuatis coeruleiscentibus.

*Eruca*

**CLASS. VI. GLOSSATA. Bombyx. 199**

*Eruca densius pilosa, pilis longissimis incanis fulvis et nigris varia. Raj. Inf. 152. 7.*

*Puppa folliculata, nigra, opaca.*

23. B. alis deflexis atris, riuulis rubicundis, posticis rubris nigro punctatis. *Syst. Ent. 582. 88. \** *Virgo.*

*Phalaena Virgo. Linn, Syst. Nat. 2. 820. 39. Mus. Lud. Vir. 381.*

*Clerk. Icon. tab. 42. fig. 5.*

Habitat in America.

24. B. alis deflexis nigris, maculis albis, posticis purpureis, macula centrali margineque nigris. *Menete.*

*Phalaena Menete. Crant. Inf. 6. tab. 70. fig. D.*

Habitat in America.

Minor B. Caia. Alae anticae fuscae macula rotundata alba ad marginem crassiorum baseos, linea in maculas tres diuisa versus marginem interiorem et transversalis abbreviata versus apicem.

25. B. alis deflexis albis nigro maculatis, posticis subtus deflorata. atris, fasciis albis. *Syst. Ent. 582. 89. \**

Habitat in America. *Mus. D. Hunter.*

**\*\*\* Alis incumbentibus.**

6. B. alis incumbentibus, anticis purpurascens, maculis ocellaribus atris, posticis rubris nigro maculatis. *Syst. Ent. 583. 90. \** *Crotalariae.*

*Phalaena Syringa. Cram. Inf. 1. tab. 5. fig. C. D.*

Habitat in Crotalariae Indiae orientalis.

7. B. alis incumbentibus obscuris, maculis numerosis subocellaribus fuscis, posticis rubris nigro maculatis. *Ricini.*

*Syst. Ent. 583. 91. \**

Habitat in Indiae orientalis Ricini. *Mus. Tottianum.*

8. B. alis incumbentibus niueis, anticis costa sanguinea, posticis maculis atris. *sanguinolenta.*

*Cram. Inf. 13. tab. 183. fig. D.*

Habitat in India orientali. *Dom. Lund.*



200 CLASS. VI. GLOSSATA. Bombyx.

Statura praecedentium. Caput niueum palpis antennisque nigris fasciaque postica sanguinea. Thorax niueus fascia media sanguinea. Abdomen fuluum annulis nigris. Alae omnes niueae. Anticae costa sanguinea punctisque aliquot indistinctis nigris minutissimis. Posticae vtrinque maculis aliquot atris.

**Hera.** 129. B. alis incumbentibus virescenti nigris, riulis flavis, posticis rubicundis, maculis tribus nigris. *Syst. Ent.* 583. 92.

*Phalaena Hera.* Linn. *Syst. Nat.* 2. 834. 91.

*Phalaena Plantaginis.* Scop. *carn.* 505.

*Geoff. Inf.* 2. 145. 74.

*Schaeff. Elem.* tab. 10. fig. 1.

— — *Icon.* tab. 29. fig. 1. 2.

*Mouff. Inf.* 91. 7. fig. 4.

*Roej. Inf.* 4. tab. 28. fig. 3.

*Klemau. Inf.* 1. tab. 41.

Habitat in Europae australioris Polygono auiculari, Antirrhino, Plantagine.

*Larva* solitaria, pilosa, nigra, fuluo maculata lineaque dorsali flava, laterali alba.

*Puppa* bis folliculata, nigra.

**Dominula.** 130. B. alis incumbentibus atris, maculis albo flavescentibus, posticis rubris nigro maculatis. *Syst. Ent.* 583. 93.

*Phalaena Dominula.* Linn. *Syst. Nat.* 2. 834. 90.

*Merian. Europ.* 2. tab. 38.

*Alb. Inf.* tab. 22.

*Wilk. pap.* 19. tab. 3. a. 3.

*Schaeff. Icon.* tab. 77. fig. 3. 4.

*Roej. Inf.* 3. tab. 47.

Habitat in Lamio, Salice polyphaga.

*Larva* gregaria pilosa, atra, lincis tribus flavis albo maculatis.

*Puppa* brunnea intra folliculum communem.

**credula.** 131. B. alis incumbentibus corporeque atris, albo punctatis. *Syst. Ent.* 584. 94. \*

*Phalaena Sybaris.* Cram. *Inf.* 6. tab. 71. fig. E.

Habitat in America meridionali.

CLASS. VI. GLOSSATA. Bombyx. 201

12. B. alis incumbentibus nigris, maculis coeruleis flapis *Letrix*.  
albisque, posticis rubro alboque maculatis. *Syst. Ent.*

584. 95.

*Phalaena Letrix*. Linn. *Syst. Nat.* 2. 834. 89. *Mus.*

*Lud. Vlr.* 389.

*Cram. Inf.* 16. tab. 192. fig. 6.

*Edw. Av.* tab. 318.

Habitat in China.

Abdomen rubrum, fasciis nigris.

13. B. thorace variegato, alis griseis, basi apiceque albis *Furcula*.  
nigro punctatis. *Syst. Ent.* 584. 96.

*Phalaena Furcula*. Linn. *Syst. Nat.* 2. 823. 51. *Fn. Sv.*

1122.

*Phalaena* elinguis, antennis pectinatis, thorace cristato variegato, alis griseis, strigis nigris flauo marginatis punctisque nigris. *Degeer Vers. Germ.* 2. 1. 226. 4. tab. 4. fig. 21.

*Harris Inf. angl.* tab. 4. fig. 9.

*Wilk. pap.* 13. tab. 1. fig. 1.

*Sepp. Inf.* 4. 29. tab. 6.

Habitat in Salice.

*Larua* solitaria, nuda, bicaudata, viridis.

*Puppa* folliculata brunnea.

*Oua* nigra.

14. B. alis incumbentibus griseo fuscis, punctis duobus *Colon*.  
nigris distantibus. *Gen. Inf. Mant.* 281. \*

Habitat in Germania. *Dom. de Hattorf.*

15. B. alis incumbentibus griseo nitidis, striga postica *populeti*.  
punctorum nigrorum. *It. Norwag.* d. 6. Aug. \*

Habitat in Populo tremula Norwagiae. *Dom. Stroem.*

*Larua* viridis capite nigro maculato inter folia Populi  
filis iuncta.

16. B. alis incumbentibus, anticis ferrugineis, lunula *antiqua*.  
alba anguli postici, femina aptera. *Syst. Ent.* 584. 98.

*Phalaena antiqua*. Linn. *Syst. Nat.* 2. 825. 56. *Fn. Sv.* 1120.

*Phalaena cinerea ventricola*, corpore breui alarum  
expers. *Raj. Inf.* 173. 24.

*Geoff. Inf.* 2. 119. 23.

*Goed. Inf.* 2. tab. 29. 30.

*Swammerd. Bibl.* tab. 33. fig. 5. 6.

*Merian. Europ.* tab. 84.

*Albin. Inf.* tab. 89.

*Reaum. Inf.* 1. tab. 19. fig. 12. 17.

*Wilk. pap.* 30. tab. 3. c. 3.

*Degeer Inf.* 1. tab. 17. fig. 13. 14.

— — *Vers. Germ.* 2. 1. 238. 2.

*Roes. Inf.* 1. phal. 2. tab. 39.

— — 3. tab. 13.

Habitat in Pruno, Tilia, Alno, polyphaga.

*Larva* fasciculata, fasciculis dorsalibus quatuor albis,  
antennalibus caudalique obscuris.

*Eruca* sublutea pilosa, quatuor in medio dorso agmi-  
nulis pilorum longiorum luteorum et longo pilorum  
nigrorum penicillo in cauda. *Raj. Inf.* 344. 8.

*Puppa* folliculata nigro luteoque variegata.

*Quies* quatuordecim dierum.

*gono-* 137. B. alis incumbentibus fuscis, maculis duabus albis  
*figma.* oppositis, femina aptera. *Syst. Ent.* 585. 99. \*

*Phalaena gonostigma.* *Linn. Syst. Nat.* 2. 826. 57.

*Albin. Inf.* tab. 90.

*Wilk. pap.* 31. fig. 3. c. 4.

*Roes. Inf.* 1. phal. 2. tab. 40.

Habitat in Betula, Quercu.

*Larva* fasciculata, fasciculis octo dorsalibus albis, col-  
laribus duobus analique nigris.

*Puppa* antice flava, postice nigra.

*Quies* octodecim dierum.

*Vmber.* 138. B. alis conuolutis atris, fronte abdomineque fulvis.

*Gen. Inf. Mant.* 282. \*

*Phalaena Vmber.* *Cram. Inf.* 2. tab. 15. fig. F.

Habitat Surinami.

Antennae pectinatae. Alae omnes atrae, posticis basi  
pallidioribus.

39. B. alis conuolutis fulvis, maculis albis numerosis *Histrion*.  
coeruleo cinctis.  
Habitat in Insula Tabago Americae. *Mus. Dom. Teats*.  
Statura B. bellae. Palpi compressi, basi albi, apice nigri.  
Thorax cyaneo niger albo punctatus maculisque  
duabus fulvis. Alae anticae fulvae punctis plurimis  
magnis albis, annulo cyaneo cinctis, apice cyaneae  
punctis albis. Alae posticae, vti omnes subtus, nigrae  
margine postico parum albicante. Abdomen subtus  
album nigro annulatum ano fuluo.
40. B. alis incumbentibus flavis, fasciis sex punctorum *Pylotis*.  
nigrorum, posticis nigro punctatis. *Syst. Ent.*  
585. 100.\*  
*Phalaena Pylotis*. *Drury Inf.* 2. tab. 6. fig. 3.  
*Phalaena cribraria*. *Clerk. Inf.* tab. 54. fig. 4.  
Habitat in noua Hollandia. *Mus. Dom. Banks*.
41. B. alis conuolutis flavis, fasciis sex punctorum nigro- *bella*.  
rum, posticis rubris apice nigris. *Syst. Ent.* 585. 101.  
*Phalaena bella*. *Linn. Syst. Nat.* 2. 884. 348. *Mus.*  
*Lud. Vlr.* 399.  
*Phalaena minor* fulua maculis nigris alba linea pulcre  
adpersis. *Raj. Inf.* 211. 1.  
*Drury Inf.* 1. tab. 24.  
*Cram. Inf.* 10. tab. 109. fig. C. D.  
*Petiv. Gazoph.* 1. tab. 3. fig. 1.  
*Catesb. Carol.* 2. 96. tab. 96.  
Habitat in America septentrionali.
42. B. alis albis, anticis nigro sanguineoque punctatis, *pulchella*.  
posticis apice nigris. *Syst. Ent.* 586. 102.  
*Phalaena pulchella*. *Linn. Syst. Nat.* 2. 884. 349.  
*Phalaena Lotrix*. *Cram. Inf.* 10. tab. 109. fig. E. F.  
*Petiv. Gazoph.* 1. tab. 3. fig. 3.  
Habitat in Solano tomentoso Orientis, Americae in no-  
ua Hollandia. *Mus. Dom. Banks*.
43. B. alis depressis rubris atro punctatis, posticis albo *ornatrix*.  
nigroque variis. *Syst. Ent.* 586. 103.  
*Phalaena Ornatrix*. *Linn. Syst. Nat.* 2. 839. 110. *Mus.*  
*Lud. Vlr.* 384.

*Drury*

204 CLASS. VI. GLOSSATA. Bombyx.

*Drury Inf.* 1. tab. 74. fig. 3.

*Cram. Inf.* 14. tab. 166. fig. C. D. F.

*Seb. Mus.* 4. tab. 39. fig. 19. 20.

Habitat in America meridionali.

An satis distincta a B. bella?

*priuerna.* 144. B. alis incumbentibus, anticis fuscis fascia flaua, posticis fuluis, margine atro.

*Phalaena priuerna.* *Cram. Inf.* 14. tab. 166. fig. E.

Habitat Surinami.

Statura praecedentium. Alae anticae fuscae, basi fuluae striatae, pone medium fascia flaua. Posticae fulvae margine atro. Thorax flauus. Abdomen fuluum.

*Iesuita.* 145. B. alis incumbentibus atris stria fulua. *Syst. Ent.* 586. 104.\*

Habitat in Indiis.

*vestita.* 146. B. alis incumbentibus nigris immaculatis, abdomine subtus villoso albo. *Syst. Ent.* 586. 105.\*

*Phalaena elinguis*, antennis pectinatis, alis fuscis, femina aptera. *Degeer Versf. Germ.* 2. 1. 220. 6. tab. 3. fig. 13.

*Phalaena vicella.* *Wien. Verz.* 288. tab. 1. a. fig. 7.

— — — — — tab. 1. b. fig. 7.

*Tinea inuolucro palearum vnico tecto.* *Geoff. Inf.* 2. 203. 50.

*Reaum. Inf.* 3. tab. 11. fig. 7. 8.

*Frisch. Inf.* 6. 16. tab. 7.

Habitat in Europae Corylo.

*Larua* grisea, nigro punctata, degit intra cylindrum e graminibus aridis filo connexis constructum, quem folii superficiei interiori affigit.

*Imaginis* femina aptera.

*annulata.* 147. B. alis incumbentibus atris, niueo maculatis, tibiis albo annulatis. *Gen. Inf. Mant.* 282.\*

Habitat Hamburgi. *D. Schulz.*

*Graminis.* 148. B. alis depressis griseis, linea trifurca punctoque albidis. *Syst. Ent.* 586. 106.

*Phalaena Graminis.* *Linn. Syst. Nat.* 2. 830. 73. *F.* Sv. 1140.

*Phalaena*

*Phalaena* antennis pectinatis, alis deflexis griseis, stria longitudinali ramosa alba maculisque tribus flavescentibus. *Degeer Inf. Vers. Germ.* 2. L. 247. 5.

*Frisch. Inf.* 10. tab. 21.

*Alt. Holmenf.* 1742. 40. tab. 2.

*Raj. Inf.* 228. 4.?

*Harr. Inf. angl.* tab. 5. fig. 7.

8. *Bombyx popularis* alis deflexis fuscis, albo striatis, stigmatibus albis, pupilla brunnea. *Syst. Ent.* 577. 71.\*

Habitat in Gramine, excepto Alopecuro, prata deval vastans.

*Larva* glabra, obscura, stria laterali dorsali que flauis praedatur a subibus, coruis.

Quies quatuordecim dierum.

149. B. alis incumbentibus griseo fuscoque variegatis, tho. *fulminea*.  
race antice albo, striga nigra. *Gen. Inf. Mant.* 283.\*

Habitat Hamburgi. *D. Schulz.*

150. B. alis incumbentibus atris, rubro flauoque variegatis, *Gloriosas*.  
posticis nigris, margine flauo. *Syst. Ent.* 587. 107.  
Habitat in Indiae orientalis Gloriosa.

151. B. alis incumbentibus atris ante marginem subferrugineis, posticis albis. *Syst. Ent.* 587. 108.

Habitat in Indiae orientalis Crino asiatico. *Museum Tottianum.*

152. B. alis incumbentibus roseis, strigis tribus fuscis, secunda vndata, tertia punctata. *Syst. Ent.* 587. 109. *rosea.*

*Phalaena* pectinicornis elinguis, alis deflexis roseis, superioribus punctorum arcuumque nigrorum ordine duplici. *Geoff. Inf.* 2. 121. 25.

*Phalaena miniata* geometra feticornis, alis rotundatis, omnibus pallide miniatis, anticis linea vndata ad basin, characteribus in medio et punctis versus marginem posticum nigris. *Forst. Cent.* 1. 75. 75.

*Phalaena minor*, alis velut miniaceis, punctis et lineolis nigris in medio notatis. *Raj. Inf.* 227. 86.

*Harr. Inf. angl.* tab. 30. fig. p.

*Natursf.* 14. tab. 1. fig. 18.

Habitat in Europae hortis.

206 CLASS. VI. GLOSSATA. Bombyx.

*rubricollis*. 153. B. alis incumbentibus atra, collari sanguineo, abdomine flavo. *Syst. Ent.* 587. 110.

*Phalaena rubricollis*. *Linn. Syst. Nat.* 2. 840. 113. *Fa. Sv.* 1154.

*Phalaena feticornis spirilinguis*, alis deflexis nigricantibus, collari purpureo, abdomine flavo. *Gress. Inf.* 2. 148. 79. tab. 12. fig. 6.

*Clerk. Icon.* tab. 2. fig. 3.

*Schaeff. Icon.* tab. 59. fig. 8. 9.

Habitat in Europae sylvis haud infrequens.

*fuliginosa*. 154. B. alis incumbentibus rubro fuliginosis, puncto gemino nigro, abdomine sanguineo, dorso nigro. *Syst. Ent.* 588. 111.

*Phalaena fuliginosa spirilinguis laevis*, alis deflexis, rubro fuliginosis, puncto gemino nigro, inferioribus rubro marginatis. *Linn. Syst. Nat.* 2. 836. 95. *Fa. Sv.* 1159.

*Phalaena minor*, alis obscure rufis seu pullis, duabus maculis nigris notatis. *Raj. Inf.* 228. 13.

*Harr. Aurel.* tab. 12.

— — *Inf. Anglic.* tab. 8. fig. 7. vix differe videtur.

*Ammir. Inf.* tab. 30.

*Roes. Inf.* 1. phal. 2. tab. 43.

*Wilk. pap.* tab. 3. a. 14.

Habitat in Rapa, Sinapi, Gramine, Betula.

*Larva* pilosa, ferruginea, capite pedibusque anticis atris. Hyeme in niue Norwagiae maxima in copia obambulans aetatem frigidiorum et annonae caritatem indicat. Stroem.

*Puppa* folliculata atra postice flavo fasciata.

*cribrum*. 155. B. alis incumbentibus, anticis albis transverse nigro punctatis. *Syst. Ent.* 588. 112.

*Phalaena cribrum*. *Linn. Syst. Nat.* 2. 831. 76. *Fa. Sv.* 1136.

Habitat in Europa boreali.

*obscura*. 156. B. alis incumbentibus concoloribus fuscis, anticis punctis tribus albis posticis immaculatis.

Habitat in Italia. *Dr. Allioni.*

Corpus

Corpus paruum, flauum abdomine linea dorsali e punctis atris. Alae anticae obscure fuscae, minime nitidae. Versus apicem puncta tria approximata transversim posita, alba, interioribus duobus maioribus, posticae obscure fuscae omnino immaculatae.

157. B. alis incumbentibus concoloribus, anticis fuscis al- punctata.  
bo punctatis, posticis flauis apice fuscis.

Habitat in Italia. D. Allioni.

Statura praecedentis at paullo minor. Corpus fuscum, abdomine flauo, linea dorsali punctorum nigrorum. Alae anticae fuscae punctis duobus in medio tribusque versus apicem exteriori minori albis. Alae posticae flauae margine late nigro.

## 146. HEPIALUS.

H. alis flauis fuluo striatis, maris niucis. Syst. Ent. Humuli.  
589. 1.

Phalaena Humuli elinguis fulua, antennis thorace brevioribus, maris alis niucis. Linn. Syst. Nat. 2. 833.  
84. Fn. Sv. 1147.

Phalaena media, alis longis, in femina flauis cum lineis rubentibus, in mare albis. Raj. Inf. 157. 3.

Degeer Inf. 1. tab. 7. fig. 5. 6.

— — — Verf. Germ. 2. 1. 266. 4.

Harr. Inf. angl. tab. 4. fig. a — d.

Salz. Hist. Inf. tab. 22. fig. 1.

Habitat in Lupuli radicibus Erysiphen caussans, pellitur stercore suillo.

Larua 16poda subglabra, pallida, capite testaceo.

Puppa folliculata, cylindrica, aculeata, viuacissima.

Oua innumera, minima, nigra.

H. alis cinereis, striga albidior. Syst. Ent. 589. 2. lupulinus.

Phalaena lupulina elinguis lutea, thorace brevioribus, alis macula arcuque cinerascens. Linn. Syst. Nat.  
2. 833. 86. Fn. Sv. 1149.

Clerk. Icon. tab. 9. fig. 4.

Habitat in Europa boreali.



*hectus.* 3. *H. luteus*, alis deflexis, anticis fasciis duabus albidis, obliquis punctato interruptis. *Syst. Ent.* 589. 3.

*Phalaena hecta.* Linn. *Syst. Nat.* 2. 833. 85. *Fn. Sv.* 1148.

*Phalaena clauipes* elinguis, antennis filiformibus brevissimis, alis maiori rufo fulvis, maculis niueis, feminae fuscis, fasciis fulvis. *Degeer Inf.* 7. 596. 4. tab. 44. fig. 14.

*Degeer Inf.* 1. tab. 7. fig. 11.

Habitat in Europae borealis sylvis frequens.

Vespere in aere fluctitat motu in pendulo, at solitarius. Aliam copula vinctam vidi simillimam, at loco fasciarum albidarum fasciae duae oblique ceruinae, in quarum posteriore ad marginem tenuiorem punctum fuscum.

*scalaris.* 4. *H. niueus*, strigis numerosis abbreviatis nigris striaque longitudinali fulva. *Syst. Ent.* 590. 4.\*

Habitat in China. *Mus. Dom. Monson.*

*Aesculi.* 5. *H. niueis*, alis punctis numerosis coerulesco nigris, thorace fenis. *Syst. Ent.* 590. 5.

*Phalaena Aesculi* elinguis laevis niuea, antennis thorace brevioribus, alis punctis numerosis coerulesco nigris, thorace fenis. Linn. *Syst. Nat.* 2. 833. 87. *Fn. Sv.* 1150.

*Phalaena Aesculi.* *Naturf.* 12. tab. 2. fig. 7.

*Wien. Verzeichn.* tab. tit. praef.

*Acta Soc. Berol. phys.* 3. tab. 1. fig. 1. 2.

*Harr. Inf. angl.* tab. 2. fig. 3. 4.

*Pod. Inf.* 88. 16.

*Reaum. Inf.* 2. tab. 38. fig. 1 — 4.

*Roes. Inf.* 3. tab. 48. fig. 5. 6.

*Schaeff. Icon.* tab. 38. fig. 8. 9.

Habitat in Pyro, Aesculo.

*Larva* flava nigro punctata, capite caudaque nigra. Antennae basi pectinatae apice filiformes.

*pyrinus.* 6. *H. niueus*, alis punctis numerosis transuersalibus nigris, thorace duodecim cyaneis. *Syst. Ent.* 590. 6.\*

Habitat in America boreali.

147. N O C T U A.

\* Alis patulis.

N. alis patulis variegatis, subtus ferrugineis nigro vn- *Zenobia*.  
datis.

*Phalaena Zenobia*. *Cram. Inf.* 10. tab. 115. fig. A. B.  
Habitat Surinami.

Magnitudo N. Strigis. Alae omnes supra albo coeruleo ferrugineo nigroque variegatae. Maculae duae albae distinctae in alis anticis, posteriori maiori rubro punctata. Subtus omnes ferrugineae puncto strigisque vndatis nigris.

N. alis patulis concoloribus, albo nigroque reticulatis *Strix*.  
nebulosisque. *Syst. Ent.* 591. 2.

*Phalaena Strix*. *Linn. Syst. Nat.* 2. 833. 82. *Muf.*  
*Lud. Vlr.* 377.

*Phalaena Agrippina*. *Cram. Inf.* 8. tab. 87. fig. A.  
— — — — — tab. 88. fig. A.

*Clerk. Icon.* tab. 51. fig. 1.

*Merian. Surin.* 20. tab. 20.

*Seb. Muf.* 4. tab. 39. fig. 12. 14.

— — — — — tab. 57. fig. 7. 9.

Habitat in America meridionali.

*Larua nuda nigra*, annulis lineaque laterali vire-  
scentibus.

N. alis patulis dentatis fuscis, nigro vndulatis, anticis *Bubo*.  
macula magna ocellari brunnea. *Syst. Ent.* 591. 3.\*

*Phalaena Macrops feticornis spirilinguis*, alis fuscis  
atro vndulatis, superioribus supra ocello ferrugineo,  
subtus albo maculatis. *Linn. Syst. Nat.* 3. 225.

*Cram. Inf.* 15. tab. 171. fig. A. B.

*Satz. Hist. Inf.* tab. 22. fig. 2.

Habitat in China. *Muf. Dom. Monson.*

In quibusdam speciminibus, forte feminis, margo exterior alae posticae reflexus saeculum format, qui magnam copiam serici laeuissimi continet forte ad involuenda oua.

*Fabr. Sp. Inf. T. II.*

Q

4. N.

*Iphianasse*. 4. N. alis patulis dentatis fuscis, ferrugineo vndatis, posticis basi atris, striga coerulescente.

*Phalaena Iphianasse*. *Cram. Inf.* 15. tab. 172. fig. A.  
Habitat Surinami.

Magna. Alae omnes basi obscuriores, anticis maculis duabus subocellaribus ferrugineis, posticis striga coerulescente.

*Aluco*. 5. N. alis patulis dentatis brunneis, nigro vndulatis maculisque tribus marginalibus fuscis. *Syst. Ent.* 591. 4.\*

*Phalaena occidua*. *Cram. Inf.* 15. tab. 173. fig. A. B.  
Habitat in China. *Mus. Dom. Monson*.

*Caprimulgus*. 6. N. alis patulis dentatis fuscis, nigro vndulatis, anticis oculo coerulescente. *Syst. Ent.* 591. 5.\*

Habitat in China. *Mus. Dom. Monson*.

*odora*. 7. N. alis patulis dentatis fuscis, nigro vndatis, anticis oculo atro, lunula coerulea strigaque maculari alba.

*Phalaena odora* pectinicornis spirilinguis, alis crenatis nebulosis, anticis oculo unico, inferioribus sesquialtero. *Linn. Syst. Nat.* 2. 811. 11. *Mus. Ind. Vir.* 374.

*Clerk. Icon.* tab. 50. fig. 1.

*Sloan. Iamaic.* 2. tab. 236. fig. 13. 14.

*Drury Inf.* 1. tab. 3. fig. 1.

*Cram. Inf.* 15. tab. 169. fig. A. B.

Habitat Surinami.

Sexus alter fascia communi e strigis tribus vndatis albis in medio.

*Conf. Phalaena Agarista*. *Cram. Inf.* 15. tab. 170. fig. A. B. vix distincta videtur.

*Mycerina*. 8. N. alis patulis dentatis, anticis vndatis strigaque coerulea, posticis strigis duabus albis coeruleaque.

*Phalaena Mycerina*. *Cram. Inf.* 15. tab. 172. fig. B.  
Habitat Surinami.

Alae anticae obscurae fusco vndatae, ad basin maculae duae ordinariae, posteriori maiori reniformi, in medio striga e lunulis concatenatis coeruleis, quae utrinque lunula alba terminantur. Versus marginem posticum striga vndata atra. Posticae basi nigrae strigis

CLASS. VI. GLOSSATA. Noctua, 211

strigis duabus albis, apice dilutiores striga media  
coerulea duabusque obscurioribus.

N. alis patulis dentatis, nigro undulatis, anticis fascia *Vlula*,  
abbreviata alba maculaque subocellari. *Syst. Ent.*  
592. 6.\*

*Phalaena hieroglyphica. Drury Inf.* 2. tab. 2. fig. 1.

*Phalaena Hermionia. Cram. Inf.* 15. tab. 174. E.

*Phalaena Mygdonia. Cram. Inf.* 15. tab. 174. F.

Habitat in India.

Alae rarius omnino atrae.

Conf. *Phalaena Lathonia. Cram. Inf.* 2. tab. 13. fig. B.

1. N. alis patulis, griseo fuscoque undatis, anticis linea *Spiralis*,  
spirali conuoluta nigra. *Syst. Ent.* 592. 7.\*

Habitat in China. *Mus. Dom. Monson.*

N. alis patulis fuscis, fascia maculaque marginali albis, *crepuscu-*  
anticis ocello. *laris.*

*Phalaena crepuscularis feticornis spirilinguis, alis*  
*griseis fascia maculaque albis, superioribus ocello*  
*fusco. Linn. Syst. Nat.* 2. 811. 13. *Mus. Lud.*  
*Vir.* 378.

*Clerk. Icon.* tab. 53. fig. 1. 2.

*Cram. Inf.* 13. tab. 159. fig. A.

Habitat in America.

Alter sexus pupilla duplici ocelli, alter ocello omni-  
no coeco.

N. alis patulis nigricantibus, anticis lunula vittaque *pandrosa*,  
niueis.

*Phalaena pandrosa. Cram. Inf.* tab. 77. fig. D.

Habitat — — — *Dom. Teats.*

Præcedentibus paullo minor. Antennae pectinatae ra-  
diis ferrugineis. Palpi apice compressi puncto nigro.  
Alae omnes nigricantes margine postico pallidiore,  
anticis lunula media vittaque, quae tamen margi-  
nem posticum haud attingit niueis. Subtus griseae  
striga pallidiore, anticis punctis duobus posticis uni-  
co nigris.

212 CLASS. VI. GLOSSATA. Noctua.

*illibata*. 13. N. alis patulis fuscis, macula apicis nigra. *Syst. Ent.* 592. 8.\*

Habitat in India. *Mus. Dom. Banks.*

\*\* Thorace laeui, alis planis incumbentibus.

*scrobiculata*. 14. N. laevis, alis incumbentibus subdentatis griseis, scrobiculo bascos nigro, posticis vitta marginali nigra. *Syst. Ent.* 592. 9.\*

Habitat in India orientali. *Mus. Tottianum.*

*Dioscoreae*. 15. N. laevis, alis incumbentibus dentatis griseis, posticis luteis, lunula limboque nigris. *Syst. Ent.* 593. 10.\*

*Phalaena fullonica* feticornis spirilinguis, alis flavis, inferioribus concoloribus, disco flavis lunula limboque nigris. *Linn. Syst. Nat.* 2. 812. 16.

*Clerk. Icon.* tab. 48. fig. 1. 2. 3.

*Phalaena Cajeta*. *Cram. Inf.* 3. tab. 30. fig. A. B. C.

*Phalaena Pomona*. *Cram. Inf.* 7. tab. 77. fig. C.

Habitat in Indiae orientalis Dioscoreis. *Dr. Koenig.*

Conf. *Phalaena Ancilla*. *Cram. Inf.* 13. tab. 149. fig. F.

— — — *Procus* — — — 13. tab. 149. fig. G.

— — — *Cocalus* — — — 12. tab. 134. fig. B.

Alae anticae colore nigris variant.

*materna*. 16. N. laevis, alis incumbentibus, posticis fulvis macula margineque albo punctato atris.

*Noctua hybrida* laevis alis incumbentibus, anticis flavis linea niuea, limbo pallidiore fusco maculato. *Syst. Ent.* 593. 11.\*

*Phalaena materna* spirilinguis cristata, alis deflexis griseis, inferioribus luteis, puncto nigro. *Linn. Syst. Nat.* 2. 840. 117.

*Phalaena Salaminca*. *Cram. Inf.* 15. tab. 174. fig. A.

*Phalaena Materna*. *Cram. Inf.* 15. tab. 174. fig. B.

*Drury Inf.* 2. tab. 13. fig. 4.

Habitat in Indiis.

Alarum anticarum color maxime variat.

17. *N. laevis*, alis incumbentibus cinereis, posticis rufescentibus, macula atra. *Syst. Ent.* 593. 12.\* *micror-rhoea.*  
Habitat in noua Hollandia. *Mus. Dom. Banks.*
18. *N. laevis*, alis incumbentibus griseis, punctis strigisque quatuor fuscis, posticis apice nigris albo maculatis. *Syst. Ent.* 593. 13.\* *serua.*  
Habitat in noua Hollandia. *Mus. Dom. Banks.*
19. *N. laevis*, alis incumbentibus, anticis viridibus apice fuscis, posticis flauis, macula atra. *Tirrhaea.*  
*Phalaena Tirrhaea. Cram. Inf.* 15. tab. 172. fig. E.  
Habitat ad Cap. Bon. Spei.  
Corpus fuluum. Alae anticae virides maculis duabus nigris, apice fuscae, posticae flavae macula magna nigra.
20. *N. laevis* alis incumbentibus albis, anticis immaculatis, posticis apice virescenti striatis. *Chione.*  
Habitat in India orientali. *Dom. Lund.*  
Corpus magnum, album, nigro punctatum abdominis, dorso fuluo. Alae anticae supra albae, immaculatae, subtus nigrae linea maculisque aliquot albis, posticae albae apice lineolis maculisque aliquot virescenti nigris nitidis, subtus concolores.
21. *N. laevis*, alis incumbentibus concoloribus fuscis, disco hyalino, thorace niueo nigro punctato. *Astrea.*  
*Syst. Ent.* 594. 14.\*  
*Phalaena Astrea. Drury Inf.* 2. tab. 28. fig. 4.  
Habitat in noua Hollandia. *Mus. Dom. Banks.*
22. *N. laevis*, alis incumbentibus? fuscis, vitta alba, posticis flauis apice coeruleis. *Narcissus.*  
*Phalaena Narcissus. Cram. Inf.* 7. tab. 73. fig. E. F.  
Habitat in China.  
Magna. Thorax fuscus nigro punctatus. Abdomen fuluum nigro punctatum.
23. *N. laevis* alis incumbentibus? anticis fuscis maculis duabus strigaeque albis, posticis flauis nigro maculatis. *bajularia.*  
*Phalaena Bajularia. Cram. Inf.* 15. tab. 172. fig. C.  
Habitat in Amboina.

Corpus fuscum abdomine flavo. Alae anticae fuscacul  
culis duabus posteriori maiori strigaeque versus m  
ginem albis. Posticae flavae fasciis duabus atris  
steriori maculari.

*Quercus.* 24. N. laevis, alis incumbentibus griseis, strigis tril  
fucis. *Syst. Ent.* 594. 15.\*  
Habitat in Europae Quercu.

*puncta-* 25. N. laevis, alis incumbentibus griseis, fusco vnda  
*ta.* punctis quatuor marginis exterioris nigris. *S*  
*Ent.* 594. 16.\*  
Habitat in Anglia. *Mus. Dom. Monson.*

*pallens.* 26. N. laevis, alis incumbentibus pallidis immacula  
marginibus posticis subtus nigro punctatis. *S*  
*Ent.* 594. 17.

*Phalaena pallens.* Linn. *Syst. Nat.* 2. 838. 107.  
*Sv.* 1175.

*Phalaena* antennis setaceis, alis deflexis cinereis, ve  
pallidis. *Degeer Inf. Vers. Germ.* 2. 1. 301. 8. 1  
7. fig. 14.

*Clerk. Icon.* tab. 4. fig. 6.

Habitat in Europae hortis.

*Larva* pilosa nigra, cinereo irrorata, lineisque quatu  
albis.

*Puppa* folliculata, fusca.

*compla-* 27. N. laevis, alis depressis plumbeis, margine exterio  
*na.* pallidis, posticis totis flavis. *Syst. Ent.* 594. 18.

*Phalaena complana.* Linn. *Syst. Nat.* 2. 840. 115. *A*  
*Sv.* 1153.

*Alb. Inf.* tab. 70. fig. G. H.

Habitat in Europae Quercu.

Variat rarius alis anticis totis pallidis.

*quadra.* 28. N. laevis, alis depressis luteis, anticis punctis duob  
cyaneis. *Syst. Ent.* 595. 19.

*Phalaena quadra.* Linn. *Syst. Nat.* 2. 840. 114.

*Wilk. pap.* 24. tab. 3. a. 17.

*Poda Inf.* tab. 2. fig. 8.

*Schaeff. Elem.* tab. 98. fig. 5.

*Roef. Inf.* 1. phal. 2. tab. 17.

Habitat in Pino, Quercu.

*Lar*

*Larva* pilosa, dorso punctis rubris maculoque nigra,  
lateribus lineis nigris.

*Puppa* folliculata, atra.

29. *N. laevis*, alis incumbentibus cinereis, thorace flavo, *deplana*.  
*Syst. Ent.* 595. 20.

*Phalaena deplana* spirilinguis laevis, alis depressis, superioribus liuidis margine bascos nigris. *Linn. Mant.* 1. 539.

Habitat in Europae hortis.

Praecedenti affinis. Pedes antici cyanei.

30. *N. laevis*, alis conuolutis pallidis flavo venosis, ant. conuolutis maculis duabus angulatis atro coeruleis. *ta*.

*Phalaena Entella*. *Cram. Inf.* 18. tab. 208. fig. D.

Habitat ad littora malabarica.

Statura *N. exoletae*. Alae conuolutae omnes pallidae flavo venosae. Anticae maculis duabus angulatis ad costam concatenatis atro coeruleis.

31. *N. laevis*, alis planis, anticis cinereis, strigis duabus *marboneae*.  
flavis posticeque puncto fusco. *Syst. Ent.* 595. 21.\*

*Phalaena marboneae*. *Linn. Syst. Nat.* 2. 837. 103.

Habitat in Europa australi.

32. *N. laevis*, alis depressis planis fusco irroratis, antennis pectinatis. *Boleti*.  
*Gen. Inf. Mant.* 282.\*

Habitat Kiliae in Boletio versicolore.

\* \* \* Thorace laevi, alis deflexis.

33. *N. laevis*, alis deflexis cinereis, basi fulvo albo nigroque maculatis, antennis pectinatis apice setaceis. *Ficus*.  
*Syst. Ent.* 595. 22.\*

Habitat in Ficu racemosa Indiae orientalis. *D. Koenig. Mus. Tottianum*.

34. *N. laevis*, alis deflexis cinereis, strigis punctoque medio albis, basi fulvis, punctis quinque atris. *Caricae*.  
*Syst. Ent.* 596. 23.\*

*Phalaena Alciphron*. *Cram. Inf.* 12. tab. 133. fig. E.  
Habitat in Ficu Indiae orientalis. *Mus. Tottianum*.

35. *N. laevis*, alis deflexis fuscis, strigis tribus obscurioribus stigmatumque postico nigro punctato. *coronata*.  
*Syst. Ent.* 596. 24.\*

Habitat in China.



*Manlia.* 36. N. laevis, alis deflexis brunneis, posticis fascia coerulea margineque nigro.

*Phalaena Manlia.* Cram. Inf. 8. tab. 92. fig. A.

Habitat in Asia.

Magna. Alae anticae brunneae fascia abbreviata in medio obscuriore maculisque ordinariis pallidioribus. Posticae basi brunneae, in medio fascia lata coerulea, apice atrae.

*Geryon.* 37. N. laevis, alis deflexis atris, flauo maculatis, posticis rubris, margine atro.

Habitat in Africa aequinoctiali. *Mus. Dom. Teats.*

Magna. Antennae setaceae, nigrae. Caput et thorax atris, albo punctata. Abdomen fuscum, supra annulis flavis, subtus albis ano fuluo. Alae anticae atrae macula oblonga ad marginem interiorem bascos et quinque in disco flauescentibus. Posticae rubrae, margine late nigro. Apex alarum albicat.

*leonina.* 38. N. laevis, alis deflexis brunneis, stigmatibus griseis, posticis flavis, maculis duabus nigris, abdomine cin-gulis nigris. *Syst. Ent.* 596. 25.\*

Habitat in India. *Mus. Dom. Banks.*

*Dama.* 39. N. laevis, alis deflexis fuscis, albo striatis, posticis al-bis, margine nigro. *Syst. Ent.* 596. 26.\*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

*margina-ta.* 40. N. laevis, alis deflexis variegatis, posticis atris, mar-gine albo. *Syst. Ent.* 597. 27.\*

Habitat in America. *D. Hunter.*

*virescens.* 41. N. laevis, alis deflexis virescentibus, strigis tribus saturationibus. *Gen. Inf. Mant.* 283.\*

Habitat in Americae St. Gruæ.

*Batis.* 42. N. laevis, alis deflexis, anticis fuscis, maculis quinque incarnatis, posticis albidis. *Syst. Ent.* 597. 28.

*Phalaena Batis.* Linn. *Syst. Nat.* 2. 836. 97. *Fn. Sv.* 158.

*Merian. Europ.* 3. tab. 21. fig. 1.

*Reaum. Inf.* 1. tab. 7. fig. 2.

*Roef. Inf.* 4. tab. 26.

Habitat in Ruba.

*Larua* nuda ferruginea, antice posticeque gibba.

*Puppa* postice attenuata, fusca.

43. *N. laevis*, alis deflexis concoloribus fulvis, margine nigro. *Syst. Ent.* 597. 29. *Fulvia*.  
*Phalaena Fulvia*. *Linn. Syst. Nat.* 2. 836. 96. *Muf.*  
*Lud. Vlr.* 385.  
*Clerk. Icon.* tab. 55. fig. 6.  
 Habitat in Indiis.  
 Abdomen nigrum, subtus flavum.
44. *N. laevis* alis deflexis cinereo fuscoque variegatis, sub- *glyphica*.  
 tus luteis fusco fasciatis. *Syst. Ent.* 589. 31.  
*Phalaena glyphica* spirilinguis laevis, alis patulis fuscescentibus, maculis hieroglyphicis nigris, subtus fascia atra. *Linn. Syst. Nat.* 2. 838. 105. *Fn. Sv.* 1161.  
*Phalaena feticornis* spirilinguis, alis patentibus supra fuscis pone subtusque flavescentibus. *Geoff. Inf.* 2. 136. 55.  
*Huffnag. pict.* 4. tab. 11. fig. 3.  
*Reaum. Inf.* 1. tab. 49. fig. 14.  
 Habitat in Europae Verbasco.
45. *N. laevis*, alis deflexis pallidis, fascia late fusca, po- *dipfacea*.  
 sticis albo nigroque variis. *Syst. Ent.* 598. 32.  
*Phalaena dipfacea*. *Linn. Syst. Nat.* 2. 856. 185.  
*Merian. Europ.* 2. tab. 49.  
 Habitat in Sueciae agris secalinis.
46. *N. laevis*, alis deflexis fusco cinereoque variegatis, *Mi*.  
 subtus w. nigro. *Syst. Ent.* 598. 33.  
*Phalaena Mi* spirilinguis laevis, alis deflexis cinereis signo securis pallide marginato, subtus M nigro.  
*Linn. Syst. Nat.* 2. 838. 106. *Fu. Sv.* 1262.  
 Habitat in Europa boreali.  
 Praecedenti nimis affinis.
47. *N. laevis*, alis deflexis cinereis, strigis duabus vnda- *Roboris*.  
 tis albis, macula centrali niuea, lunula nigra. *Gen.*  
*Inf. Mant.* 283. \*  
*Roesf. Inf.* 1. phal. 2. tab. 50.  
 Habitat in Germaniae Quercu.  
*Larua* nuda, viridis, nitens vtrinque lineis duabus flavis.  
*Puppa* nigra.

218 CLASS. VI. GLOSSATA. Noctua.

*albicollis.* 48. N. laevis, alis deflexis, basi albis, apice fuscis, litus duplici alba.

Habitat in Italia. *D. Allioni.*

Media. Caput et thorax alba, immaculata. Antennae setaceae fuscae. Alae anticae basi albae puncto paruo nigro, apice fuscae punctis aliquot atris litumque ad marginem exteriorem et altera fusco puncta ad angulum ani. Subtus fere concolores. Posticae albae, apice fuscae.

*italica.* 49. N. laevis, alis deflexis fuscis, strigis ferrugineis maculaque transuersa marginali alba.

Habitat in Italia. *D. Allioni.*

Statura praecedentis. Caput et thorax fusca, immaculata. Abdomen cinereum segmentorum marginibus albidis. Alae anticae fuscae strigis aliquot videntur fere obsoletis obscurioribus. Macula magna transuersa alba in medio marginis crassioris. Posticae, omnes subtus, atrae basi maculaque albis.

*vulpina.* 50. N. laevis, alis deflexis, anticis obscure cinereis imaculatis, posticis apice nigris, albo maculatis. *Syst.*

*Ent.* 598. 34. \*

Habitat in India orientali. *Mus. Tottianum.*

*destituta.* 51. N. laevis, alis incumbentibus griseis, fusco maculatis striga alba, fuscae innata, submarginali. *Syst. Ent.*

598. 35. \*

Habitat in China. *Mus. Dom. Monson.*

*tigrina.* 52. N. laevis, alis deflexis, anticis variegatis, posticis nigris, fascia maculisque tribus marginalibus albis.

*Syst. Ent.* 599. 36. \*

Habitat in India orientali. *Mus. Tottianum.*

*geometrica.* 53. N. laevis, alis deflexis fuscis, macula magna atra, fasciis duabus, anteriore niuea. *Syst. Ent.* 599. 37. \*

Habitat in India orientali. *D. Koenig.*

*solida.* 54. N. laevis, alis deflexis fuscis, fasciis duabus niueis posteriore videntata. *Syst. Ent.* 599. 38. \*

Habitat in India orientali. *Mus. Tottianum.*

55. *N. laevis*, alis deflexis corporeque albis, antennis linguaeque testaceis. *Syst. Ent.* 599. 39. *nivea.*

*Phalaena nivea. Linn. Syst. Nat.* 2. 838. 208.

Habitat in Europa boreali.

56. *N. laevis*, alis deflexis fuscis, strigis tribus coerulescentibus brunneae adnatis. *Syst. Ent.* 599. 40. \* *inclita.*

Habitat in Brasilia. *Mus. Dom. Banks.*

57. *N. laevis*, alis deflexis albidis, fascia latissima saturatiore, puncto nigro margineque punctato. *Syst. Ent.* 600. 41. *trapezina.*

*Phalaena trapezina. Linn. Syst. Nat.* 2. 836. 99. *Fn. Sv.* 1157.

*Merian. Europ.* tab. 11.

Habitat in Europae Corylo.

*Larua* virefcens, lineis albidis cinereis sulphureisque, saeuit in alias laruas etiam propriae speciei. *Fuesly.*

*Puppa* folliculata, brunnea,

*Quies* 4 hebdomadum.

58. *N. laevis*, alis deflexis griseo ferrugineis, maculis strigaeque posticae flauescentibus, margine nigro punctato. *Syst. Ent.* 600. 42. \* *Cerafa.*

*Raef. Inf.* 1. phal. 2. tab. 53.

Habitat in Ceraso Angliae, Germaniae.

*Larua* flauescens, striis tribus sulphureis, capite coeruleo.

*Puppa* nuda brunnea, postice bidentata.

59. *N. laevis*, alis deflexis brunneis, punctis quatuor approximatim albis, antennis pectinatis. *Gen. Inf. Mant.* 283. \* *monilis.*

Habitat in Angliae hortis.

60. *N. laevis*, alis deflexis griseis, strigis undatis fuscis, posticis cinereis, fascia marginali nigra. *Syst. Ent.* 600. 43. \* *undata.*

Habitat in India orientali. *D. Koenig.*

61. *N. laevis*, alis deflexis dentatis cinereis, strigis fuscis, posticis flauescentibus, vitta marginali atra. *Syst. Ent.* 600. 44. \* *vittata.*

Habitat in India orientali. *D. Koenig.*

320 CLASS. VI. GLOSSATA. Noctua.

- frugalis*. 62. N. laevis, alis deflexis cinereis, fascia obliqua strigae punctorum nigrorum. *Syst. Ent.* 601. 45.\*  
Habitat Tranquebariae. *D. Koenig*.
- flava*. 63. N. laevis, alis deflexis flavis, strigis vndatis ferrugineis. *Syst. Ent.* 601. 46.\*  
Habitat Tranquebariae. *D. Koenig*.
- stigmatisans*. 64. N. laevis, alis deflexis angulatis flavis, strigis obliquis rioribus stigmatibusque niueo. *Syst. Ent.* 601. 47.\*  
Habitat Tranquebariae. *D. Koenig*.
- equestris*. 65. N. laevis, alis deflexis fuscis, fascia lata viridi. *Syst. Ent.* 601. 48.\*  
Habitat Tranquebariae. *D. Koenig*.
- reicta*. 66. N. laevis, alis deflexis albis, macula postica virescente. *Syst. Ent.* 601. 49.\*  
Habitat Tranquebariae. *D. Koenig*.
- elata*. 67. N. laevis, alis deflexis, anticis fusco-nebulosis, litumaculacque cinereis, posticis niueis.  
*Noctua litura*. *Syst. Ent.* 601. 50.\*  
Habitat Tranquebariae. *D. Koenig*.
- lanceolata*. 68. N. laevis, alis deflexis lanceolatis cinereis, anticis fusco irroratis. *Gen. Inf. Mant.* 284.\*  
Habitat in Germania. *D. Schulz*.

\*\*\*\* Thorace cristato, alis incumbentibus

- sponsa*. 69. N. cristata, alis planis cinerascens fuscis vndatis, posticis rubris, fasciis duabus nigris, abdomine vndique cinereo. *Syst. Ent.* 602. 51.  
*Phalaena sponsa*. *Linn. Syst. Nat.* 2. 841. 118.  
*Phalaena feticornis spirilinguis*, alis deflexis, anticis cinereo fuscoque nebulosis, inferioribus rubris fascia duplici transversa nigra. *Geoff. Inf.* 2. 150. 82.  
*Merian. Europ.* 2. tab. 14.  
*Reaum. Inf.* 1. tab. 32. fig. 1 — 7.  
*Roef. Inf.* 4. tab. 19.  
Habitat in Quercu Europae.

*Larva*

*Larva* nodosa, dorso verrucoso, capite coerulescente, corpore variegato.

*Puppa* folliculata, coerulescens.

Conf. *Phalaena lia*. *Cram. Inf.* 3. tab. 33. fig. C. D.

70. *N. cristata*, alis planis cinerascens, posticis rubris, *nupta*. fasciis nigris, abdomine cano subtus albo. *Syst. Ent.* 602. 52.

*Phalaena nupta*. *Linn. Syst. Nat.* 2. 841. 19.

*Raj. Inf.* 152. 4.

*Wilk. pap.* 33. tab. 1. a. 1.

*Vdm. Diff.* 73. tab. 10.

*Roef. Inf. phal.* 2. tab. 15.

Habitat in Europae Salice vitellina.

*Larva* nuda grisea, verruca caudali truncata.

1. *N. cristata*, alis griseis, posticis rubris, fasciis *pastia*. duabus nigris, abdomine supra rubro. *Syst. Ent.* 602. 53.

*Phalaena pastia*. *Linn. Syst. Nat.* 2. 841. 120. *Fn. Sv.* 1166.

*Merian. Inf.* 3. tab. 38. fig. 139.

*Wilk. pap.* 17. tab. 2. b. 1.

*Sepp. Inf.* 4. 33. tab. 7.

Habitat in Salice, Quercu.

Forte praecedentis mera varietas.

2. *N. cristata*, alis dentatis cinereo nebulosis, posticis *Fraxini*. supra nigris, fascia coerulescente. *Syst. Ent.* 602. 54.

*Phalaena Fraxini*. *Linn. Syst. Nat.* 2. 843. 125. *Fn. Sv.* 1165.

*Phalaena feticornis spirilinguis*, alis deflexis, superioribus cinereo fuscoque undulatis, inferioribus nigris, fascia transversa coerulea. *Geoff. Inf.* 2. 151. 83.

*Roef. Inf.* 4. tab. 28. fig. 1.

*Merian. Europ.* tab. 46.

*Ammir. Inf.* tab. 25.

*Wilk. pap.* 45. tab. 1. a. 2.

Habitat in Fraxino, Populo.

3. *N. cristata*, alis incumbens, posticis rubris, fascia *pronuba*. atra submarginali. *Syst. Ent.* 603. 55.

*Pha-*

*Phalaena pronuba*. Linn. *Syst. Nat.* 2. 842. 121. *Fa. Sv.* 1167.

*Phalaena* antennis setaceis, alis brunneis aut cinereis, posticis luteis, fascia marginali nigra. *Degeer Inf.*

*Verf. Germ.* 2. 1. 288. 1.

*Geoff. Inf.* 2. 146. 76.

*Raj. Inf.* 237. 18.

*Goed. Inf.* 1. tab. 14.

*Blanc. Inf.* tab. 6. fig. D. E.

*Merian. Europ.* tab. 49.

*Albin. Inf.* tab. 72. fig. C. D.

*Frisch. Inf.* 10. tab. 15. fig. 4.

*Roef. Inf.* 4. tab. 32.

*Reaum. Inf.* 1. tab. 41. fig. 4.

— — 2. tab. 14. fig. 89.

*Ammiral.* tab. 8.

*Wilk. pap.* 1. tab. 1. a. 1.

*Schaeff. Icon.* tab. 196. fig. 1. 2.

Habitat in Cheirantho, Thlaspi, Senecione.

Color alarum anticarum valde variat.

*Larva* nuda virefcens, lineis duabus dorsalibus nigris interruptis.

*Puppa* nuda, rubra.

*Epione*. 74. N. cristata, alis incumbentibus variegatis, punctis duobus albis, posticis atris, margine albo.

*Phalaena Epione*. *Cram. Inf.* 9. tab. 102. fig. E. F.

Habitat in America boreali.

Statura et magnitudo *P. pronubae*. Alae anticae corulefcentes, inprimis apice rufò nigroque variegatae. In medio puncta duo alba approximata. Subtus disco nigro macula strigaeque albis. Posticae supra atrae margine albo, subtus basi apiceque hepaticae, medio atrae margineque postico albo.

*paranympha*. 75. N. cristata, alis planis, anticis fuscis, strigis angulatis albis nigrisque, posticis flauis, fasciis duabus arcuatis atris. *Syst. Ent.* 603. 56.

*Phalaena paranympha* alis deflexis, cano fuliginosaeque undulatis, inferioribus luteis, fasciis duabus arcuatis atris. Linn. *Syst. Nat.* 2. 842. 122.

*Pha.*

*Phalaena fulminea*. Scop. carn. 510.

Roeß. Inf. 4. tab. 18. fig. 1. 2.

Habitat in Germaniae Pruno.

Larva nuda cinerea, verruca caudali reflexa, acuta.

Puppa laxè folliculata, brunnea.

Quies 4 hebdomadum.

Conf. *Phalaena Grynea*. Cram. Inf. 18. tab. 208. fig. 14.

6. N. cristata, alis planis griseo fasciatis, posticis heluo- *fimbria*.  
lis, macula lineari atra. Syst. Ent. 603. 57.

*Phalaena fimbria*. Linn. Syst. Nat. 2. 842. 123.

Schreb. Inf. fig. 9.

Habitat in Germaniae nemoribus.

7. N. cristata, alis planis fuscis, characteribus atris. Syst. Augur.  
Ent. 604. 58.\*

Habitat in Germania. Dom. de Hattorf.

8. N. cristata, alis incumbentibus ferrugineis, strigis vn- *Segetis*.  
datis obscurioribus, posticis albidis.

*Phalaena Segetum* spirilinguis cristata, alis incumben-  
tibus, superioribus fuscis, stigmatibus ordinariis li-  
neisque transuersis undulatis obscurioribus, inferiori-  
bus lacteo griseis. Wien. Verz. 252. tab. 1. a. 3.

— — — tab. 1. b. 3.

Habitat frequens in Segete Europae australioris, radices  
deuorans.

Larva nuda, liuida, segmentis singulis punctis quatuor  
nigris, capite biftriato.

Mas imaginis antennis pectinatis.

9. N. cristata, alis incumbentibus ferrugineis, strigis fu- *mercato-*  
scis vndatis, posticis apice nigris albo maculatis. Syst. *ria*.  
Ent. 604. 59.\*

*Phalaena Melicerta*. Drury Inf. 1. tab. 23. fig. 1. 3

Cram. Inf. 6. tab. 62. fig. C. D.

Habitat Tranquebariae. D. Koenig.

10. N. cristata, alis incumbentibus subdentatis griseis, *partita*.  
macula bascos atra, stigmatibus duobus brunneis stri-  
gaque media alba. Syst. Ent. 604. 60.\*

Habitat in India. D. Koenig.



*maura.* 81. N. cristata, alis incumbentibus dentatis, cinereo nigroque variis, subtus margine albo. *Syst. Ent.* 604. 61. \*

*Phalaena maura* spirilinguis cristata, alis depresso dentatis, fasciis duabus nigris, inferioribus nigris, fascia alba. *Linn. Syst. Nat.* 2. 843. 124.

*Phalaena Lemur.* *Naturf.* 6. tab. 5. fig. 1. 2

*Schaeff. Icon.* tab. 1. fig. 5. 6.

*Harris Inf. angl.* tab. 1. fig. a. b.

Habitat in Germania, in Anglia frequens.

*libatrix.* 82. N. cristata, alis incumbentibus dentato crosis, rubro griseis, punctis duobus albis. *Syst. Ent.* 604. 62.

*Phalaena libatrix.* *Linn. Syst. Nat.* 2. 831. 78. *Fa. Sv.* 1143.

*Phalaena pectinicornis* elinguis, alis cinereo flavoque rufis margine laceris. *Geoff. Inf.* 2. 121. 26.

*Phalaena elinguis*, antennis pectinatis, dorso cristata, alis crosis rubro griseis, lineis duabus punctisque duobus albis. *Degeer Inf. Verf. Germ.* 2. 1. 240. 1. tab. 5. fig. 5.

*Phalaena media domestica*, alis exterioribus rubro, rubro, cinereo et albicante colore variis cum transversa linea albicante, imo margine laciniato. *Raj. Inf.* 183. 2.

*Phalaena salictaria.* *Pod. Inf.* 92. tab. 2. fig. 9.

*Goed. Inf.* 1. tab. 67.

*Petiv. Gazoph.* tab. 19. fig. 4.

*Albin. Inf.* tab. 32. fig. 50.

*Roef. Inf.* 4. tab. 20.

*Schaeff. Icon.* tab. 124. fig. 1. 2.

*Sulz. Hist. Inf.* tab. 21. fig. 7.

*Harris Inf. angl.* tab. 1. fig. C. D.

Habitat in Glechomate, Rosa, Salice.

*Larva nuda*, viridis, flavo annulata, lineis tribus fuscis stigmatibusque rubris.

*Puppa* folliculata, nigra.

*Quies* 16 dierum.

*plata.* 83. N. cristata, alis incumbentibus fuscis, margine crassiori albo. *Syst. Ent.* 605. 63.

*Pha*

**CLASS. VI. GLOSSATA. Noctua. 225**

*Phalaena plecta* spirilinguis, subcristata, alis brunneis,  
linea nigra margineque crassiori albo. *Linn. Syst.*  
*Nat.* 2. 851. 157. *Fn. Sv.* 1216.

*Kleman. Inf.* 1. tab. 23.

Habitat in Europa.

*Larva* viridis, capite brunneo.

*Puppa* folliculata, atra.

4. *N. cristata*, alis planis fusco cinereoque nebulosis, an- *polymita*.  
gulo ani macula atra. *Syst. Ent.* 605. 64.\*

*Phalaena polymita* spirilinguis cristata, alis cinereis  
fasciatis, macula atra minima marginis tenuioris.

*Linn. Syst. Nat.* 2. 855. 180. *Fn. Sv.* 1217.

Habitat in Europae nemoribus.

1. *N. cristata*, alis incumbentibus cinereo nebulosis, *Brassicae*,  
vncio nigro ad maculam priorem.

*Phalaena Brassicae. Linn. Syst. Nat.* 2. 852. 163. *Fn.*  
*Sv.* 1205.

*Phalaena*, antennis filiformibus cristata, alis deflexis  
griseis, strigis vndatis uigris, macula grisea albaque  
striga albida. *Degeer Inf. Verf. Germ.* 2. 1. 320. 9.

*Frisch. Inf.* 10. tab. 16.

*Roef. Inf.* 1. phal. 2. tab. 29. fig. 4. 5.

*Reaum. Inf.* 1. tab. 40. fig. 16. 17.

— — — tab. 41. fig. 1. 2. 3.

*Albin. Inf.* tab. 28.

*Merian. Europ.* tab. 81.

*Goesd. Inf.* 3. tab. F.

Habitat in Europae Brassica.

5. *N. cristata*, alis incumbentibus fuscis, lineola atra, *exclama-*  
maculaque cordata, posticis albis. *Syst. Ent.* 605. 65.\* *tionis.*

*Phalaena exclamationis. Linn. Syst. Nat.* 2. 850. 155.  
*Fn. Sv.* 1190.

*Phalaena* antennis setaceis, alis incumbentibus griseis,  
maculis quatuor inaequalibus atris. *Degeer Inf. Verf.*  
*Germ.* 2. 293. 3. tab. 6. fig. 22.

*Clerk. Icon.* tab. 1. fig. 4.

*Schaff. Icon.* tab. 112. fig. 1. 2.

Habitat in Europae hortis.

*Fabr. Sp. Inf. T. II.*

P

87. N.

*l. album.* 87. N. cristata grisea, alis incumbentibus l albo notatis.  
*Syst. Ent.* 606. 66. \*

*Phalaena l. album.* Linn. *Syst. Nat.* 2. 850. 154.  
 Habitat in Europa australiori.

*citrago.* 88. N. cristata, alis incumbentibus luteis ferrugineo sub-  
 fasciatis. *Syst. Ent.* 606. 67.

*Phalaena citrago spirilinguis*, cristata, alis depressis  
 luteis, superioribus fasciis tribus ferrugineis obli-  
 quis. Linn. *Syst. Nat.* 2. 857. 189. *Fn. Sv.* 1174.

*Phalaena* antennis filiformibus cristata, alis deflexis lu-  
 teis, strigis quatuor maculisque duabus annularibus  
 ferrugineis. *Degeer Inf. Vers. Germ.* 2. 1. 313. 4. tab.  
 7. fig. 25.

*Albin. Inf.* tab. 33.

*Wilk. pap.* 5. tab. 1. a. 8.

Habitat in Pado, Salice.

*Larva* nuda fusca, lateribus flauis.

*Puppa* folliculata, brunnea.

*viminalis.* 89. N. cristata, alis incumbentibus basi fuscis, strigibus  
 datis fulvis, apice cinereis. *Gen. Inf. Mant.* 234. \*

*Roef. Inf.* 3. tab. 11.

Habitat in Germaniae Salice.

*Larva* virescens, lineis quinque longitudinalibus sin-  
 gisque albis.

*Puppa* vix folliculata, brunnea.

*pagana.* 90. N. cristata, alis incumbentibus griseo fuscis, punctis  
 marginalibus fuscis quinque, albis tribus, postica  
 albis, fascia nigra. *Syst. Ent.* 606. 68. \*

Habitat Tranquebariae. *D. Koenig.*

\*\*\*\*\* Thorace cristato, alis deflexis.

*chrysitis.* 91. N. cristata, alis deflexis orichalceis, margine fascia  
 que griseis. *Syst. Ent.* 606. 69.

*Phalaena Chrysitis.* Linn. *Syst. Nat.* 2. 843. 126. *Fn.*  
*Sv.* 169.

*Phalaena feticornis spirilinguis*, alis deflexis ferrugi-  
 neo fuscis, fascia duplici transversa viridi aurea  
*Geoff. Inf.* 2. 149. 97.

*Pha-*

*Phalaena* antennis filiformibus, dorso cristato, alis deflexis griseis, fasciis duabus aureo viridibus. *Degeer Inf. Versf. Germ.* 2. 1. 311. 2.

*Phalaena* media alis exterioribus duplici arcu transverso viridi aurato, serici instar splendente insignibus. *Raj. Inf.* 182. 45.

*Sepp. Inf.* 5. 4. tab. 1. fig. 7 — 12.

*Merian. Europ.* tab. 39.

*Albin. Inf.* tab. 71. fig. a. b. c. d.

*Natursf.* 6. tab. 3. fig. 5. 6.

*Schaeff. Icon.* tab. 101. fig. 2. 3.

Habitat in Carduis, Vrtica, Mentha.

*Larua* solitaria, gibbosa, viridis albo striata.

*Eruca* geometra maiuscula, coloribus coeruleo viridi et albicante varia. *Raj. Inf.* 183.

*Puppa* folliculata, fusca.

22. *N. cristata*, alis deflexis fuscis, macula magna lunata *orichalceae*. *Syst. Ent.* 607. 70. \*

Habitat Tranquebariae. *D. Koenig.*

23. *N. cristata*, alis deflexis, anticis flauo fuscoque variis, *Festucae*. maculis tribus argenteis. *Syst. Ent.* 607. 71.

*Phalaena Festucae*. *Linn. Syst. Nat.* 2. 845. 131. *Fn. Sv.* 1170.

*Phalaena* antennis filiformibus, dorso cristato, alis deflexis aureis, ferrugineo variis, maculis tribus argenteis. *Degeer Inf. Versf. Germ.* 2. 1. 312. 3.

*Kleman. Inf.* 1. tab. 30. fig. A.

*Albin. Inf.* tab. 84. fig. G. H.

*Petiv. Gazoph.* tab. 7. fig. 7.

*Wilks pap.* 3. tab. 1. a. 17.

*Acta Holm.* 1748. tab. 6. fig. 3. 4.

Habitat in Absinthio, Festuca fluitante.

*Larua* nuda, viridis.

24. *N. cristata*, alis deflexis, anticis fuscis  $\lambda$  aureo in- *Gamma*. scriptis. *Syst. Ent.* 607. 72.

*Phalaena Gamma*. *Linn. Syst. Nat.* 2. 843. 127. *Fn. Sv.* 1171.

*Geoff. Inf.* 2. 156. 94.

*Raj. Inf.* 163. 16.

*Sepp. Inf.* 5. 1. tab. 1 — 6.

*Goed. Inf.* 2. tab. 21.

*Blanc. Inf.* tab. 8. fig. N. P.

*Petiv. Gazoph.* 64. fig. 6.

*Merian. Europ.* 82. fig. 78.

*Albin. Inf.* tab. 79. fig. G. H.

*Frisch. Inf.* 5. tab. 15.

*Reaum. Inf.* 2. tab. 26. fig. 5.

*Wilks pap.* 34. tab. 2. a. 1.

*Schaeff. Icon.* tab. 84. fig. 5.

Habitat in Leguminibus, Oleribus, polyphaga.

*Larva* geometra, dodecapoda, viridis, striis dorsibus albis, laterali flava, capite fusco.

*Puppa* folliculata, brunnea.

*interrogationis.* 95. N. cristata, alis deflexis, anticis fusco cinereoque viridis, signo? albo inscriptis. *Syst. Ent.* 607. 73.

*Phalaena interrogationis.* *Linn. Syst. Nat.* 2. 824. 119.

*Fn. Sv.* 1172.

*Clerk. Icon.* tab. 6. fig. 7.

Habitat in Europa boreali.

*signata.* 96. N. cristata, alis deflexis, obscuris, basi nigro punctatis signo medio? aureo inscriptis. *Syst. Ent.* 608. 74.\*

Habitat Tranquebariae. *D. Koenig.*

*Peponis.* 97. N. cristata, alis deflexis griseis, macula apicis dorsali que brunneo aurea. *Syst. Ent.* 608. 75.\*

Habitat in Peponibus Tranquebariae. *D. Koenig.*

*Iota.* 98. N. cristata, alis deflexis, anticis ferrugineo griseis i refupinato aureo inscriptis. *Syst. Ent.* 608. 76.

*Phalaena Iota.* *Linn. Syst. Nat.* 2. 844. 130.

Habitat in Vrtica, Lamio albo.

*Larva* dodecapoda nuda, viridis, albo maculata.

*pauperta.* 99. N. cristata, alis deflexis, fuscis, cinereo variegatis, posticis albo hyalinis. *Syst. Ent.* 608. 77.\*

Habitat in China. *Mus. Dom. Monson.*

*meticulosa.* 100. N. alis deflexis, eroso dentatis, pallidis, anticis basi incarnata, triangulo fusco. *Syst. Ent.* 608. 78.

*halaena meticuloſa*. Linn. *Syſt. Nat.* 2. 845. 132. *Fn. Sv.* 1164.

*halaena feticornis* ſpirilinguis, alis deflexis margine croſis cinereo fufcis, ſuperioribus triangulo marginali fuſceſcente, incarnatum includente, thorace gibbo. *Geoff. Inf.* 2. 151. 84.

*halaena* antennis filiformibus, dorſo criſtato, alis dentatis deflexis grifeis ferrugineo vndique variis. *Degeer Inf. Verſ. Germ.* 2. 1. 310. 1..

*Goed. Inf.* 1. tab. 56.

*Merian. Europ.* tab. 24.

*Reaum. Inf.* 1. tab. 8. fig. 25. 26.

— — — tab. 14. fig. 12. 13.

*Albin. Inf.* tab. 13.

*Roeſ. Inf.* 4. tab. 9.

*Degeer Inf.* 1. tab. 5. fig. 14.

*Wilks pap.* 3. tab. 1. a. 3.

abitat in Cheirantho, Mercuriali, Vrtica.

*arua* nuda, viridis, linea dorſali interrupta laterali-buſque albis.

*appa* folliculata fuſca poſtice rubra.

N. criſtata, alis deflexis fuſco ferrugineoque variegatis dorſo emarginatis, fronte fulua. *Syſt. Ent.*

609. 79. \*

abitat in China. *Muf. Dom. Monſon.*

N. criſtata, alis deflexis, anticis fuſceſcentibus, arcu *gothica*. punctoque medio atris.

*halaena gothica* ſpirilinguis criſtata, alis deflexis, ſuperioribus fuſceſcentibus, arcu nigro linea alba marginato. Linn. *Syſt. Nat.* 2. 851. 159. *Fn. Sv.* 1192.

*Clark. Icon.* tab. 1. fig. 1.

abitat in Europa boreali. *Dom. Lwud.*

cus niger extrorſum verſus.

N. criſtata, alis deflexis, anticis ſupra decorticatis. *derafa*. *Syſt. Ent.* 609. 80.

*halaena derafa*. Linn. *Syſt. Nat.* 2. 851. 158.

*Naturf.* 2. tab. 1. fig. 7. mal.

abitat in Europa, Anglia.

*Sepp. Inf.* 5. 1. tab. 1 — 8.

*Goed. Inf.* 2. tab. 21.

*Blanc. Inf.* tab. 8. fig. N. P.

*Petiv. Gazoph.* 64. fig. 6.

*Merian. Europ.* 82. fig. 78.

*Albin. Inf.* tab. 79. fig. G. H.

*Frisch. Inf.* 5. tab. 15.

*Reaum. Inf.* 2. tab. 26. fig. 5.

*Wilks pap.* 34. tab. 2. a. 1.

*Schaeff. Icon.* tab. 84. fig. 5.

Habitat in Leguminibus, Oleribus, polyphaga.

*Larva* geometra, dodecapoda, viridis, striis dorsali-  
bus albis, laterali flava, capite fusco.

*Puppa* folliculata, brunnea.

*interrogationis.* 95. N. cristata, alis deflexis, anticis fusco cinereoque va-  
riis, signo? albo inscriptis. *Syst. Ent.* 607. 73.

*Phalaena interrogationis.* *Linn. Syst. Nat.* 2. 824. 129.

*Fn. Sv.* 1172.

*Clerk. Icon.* tab. 6. fig. 7.

Habitat in Europa boreali.

*signata.* 96. N. cristata, alis deflexis, obscuris, basi nigro punctis  
signo medio? aureo inscriptis. *Syst. Ent.* 608. 74.\*

Habitat Tranquebariae. *D. Koenig.*

*Peponis.* 97. N. cristata, alis deflexis griseis, macula apicis dorsali-  
que brunneo aurea. *Syst. Ent.* 608. 75.\*

Habitat in Peponibus Tranquebariae. *D. Koenig.*

*Iota.* 98. N. cristata, alis deflexis, anticis ferrugineo griseis  
i resupinato aureo inscriptis. *Syst. Ent.* 608. 76.

*Phalaena Iota.* *Linn. Syst. Nat.* 2. 844. 130.

Habitat in Vrtica, Lamio albo.

*Larva* dodecapoda nuda, viridis, albo maculata.

*pauperta.* 99. N. cristata, alis deflexis, fuscis, cinereo variegatis, po-  
sticis albo hyalinis. *Syst. Ent.* 608. 77.\*

Habitat in China. *Mus. Dom. Monson.*

*meticulosa.* 100. N. alis deflexis, erosio dentatis, pallidis, anticis  
incarnata, triangulo fusco. *Syst. Ent.* 608. 78.

*Phalaena meticulosa*. Linn. Syst. Nat. 2. 845. 132. Fn. Sv. 1164.

*Phalaena feticornis* spirilinguis, alis deflexis margine crofis cinereo fuscis, superioribus triangulo marginali fuscescente, incarnatum includente, thorace gibbo. Geoff. Inf. 2. 151. 84.

*Phalaena* antennis filiformibus, dorso cristato, alis dentatis deflexis griseis ferrugineo vndique variis. Degeer Inf. Vers. Germ. 2. 1. 310. 1.

Goed. Inf. 1. tab. 56.

Merian. Europ. tab. 24.

Reaum. Inf. 1. tab. 3. fig. 25. 26.

— — — tab. 14. fig. 12. 13.

Albin. Inf. tab. 13.

Roef. Inf. 4. tab. 9.

Degeer Inf. 1. tab. 5. fig. 14.

Wilks pap. 3. tab. 1. a. 3.

Habitat in Cheirantho, Mercuriali, Vrtica.

Larva nuda, viridis, linea dorsali interrupta laterali-  
busque albis.

Puppa folliculata fusca postice rubra.

1. N. cristata, alis deflexis fusco ferrugineoque varie-  
lata. lacera.  
gatis dorso emarginatis, fronte fulva. Syst. Ent.  
609. 79. \*

Habitat in China. Mus. Dom. Monson.

2. N. cristata, alis deflexis, anticis fuscescentibus, arcu gothica.  
punctoque medio atris.

*Phalaena gothica* spirilinguis cristata, alis deflexis, su-  
perioribus fuscescentibus, arcu nigro linea alba mar-  
ginato. Linn. Syst. Nat. 2. 851. 159. Fn. Sv. 1192.

Clerk. Icon. tab. 1. fig. 1.

Habitat in Europa boreali. Dom. Lud.

Arcus niger extrorsum versus.

3. N. cristata, alis deflexis, anticis supra decorticatis. derafa.  
Syst. Ent. 609. 80.

*Phalaena derafa*. Linn. Syst. Nat. 2. 851. 158.

Naturf. 2. tab. 1. fig. 7. mal.

Habitat in Europa, Anglia.



*diffinis-* 105.

*Pha*

Habitat

*affinis-* 106. N. cr.  
ri pun  
609. 83.

*Phalaena*

Habitat in E.

*triptera-*

107. N. cristata,  
libus, rot  
*Syst. Ent.* 6:

*Phalaena tript.*

Habitat in Euro.

*margina-*  
*ta.*

108. N. cristata, ali  
gineis postice fu.  
Habitat in Europa

*Abfynthii.*

109. N. cristata, alis c.  
tetragonum positi

*Phalaena Abfynthii*

scantibus nigricant.

bus albidis. *Lim.*

*Phalaena feticornis* f

nercis, lineis longis

*Clerk. Icon.* tab. 8.

*Frisch. Inf.* 7. tab.

*Roef. Inf.* 1. phal. 2

Habitat in Abfynthio Eur

*arua viridis, maculis flav*

*uppa folliculata brunnea*

is repandis maculaque fusca. *Linn. Syst. Nat.* 181.

feticornis spirilinguis, alis deflexis, superioribus, lineis transuersis undulatis nigris, inferioribus ferrugineis. *Geoff. Inf.* 2. 160. 99.

media, alis longis angustis, exterioribus lineolis nigris, albis atro rubentibus transverse depictis, interioribus obscure rubris. 59. 9.

*Europ.* 2. 23. tab. 9.

*Inf.* 1. tab. 15. fig. 1 — 5.

*Inf.* tab. 11.

phal. 2. tab. 11.

o, Iuglande, Quercu.

is, linea dorsali lateralibusque albis, gibba.

era, lacuis, lineis albis secundum longitudinem varia, angulosa, in extremo dorso

*Raj. Inf.* 159. 9.

deflexis fuscis, stigmatibus marginalibus typica.

idibus reticulatis. *Syst. Ent.* 613. 96.

*Linn. Syst. Nat.* 2. 857. 186. *Fn. Sv.*

phal. 2. tab. 5. 6.

fulice.

interibus albis, linea nigra.

brunnea.

alis cinereo nitidis, fascia media lucipara.

spirilinguis cristata, alis purpureis

linea nigra, stigmate postico flavo.

187. 187. *Fn. Sv.* 1201.

oreali.

ae cinereae, nitidae, in medio

ae tamen marginem tenuiorem

anticis ferrugineis, lunulata. *oleracea.*

postice bidentato. *Syst.*

*satellitica*. 104. N. cristata, alis deflexis, dentatis, brunneis, anticis puncto flavo inter punctula duo alba. *Syst. Ent.* 609. 81.

*Phalaena satellitia*. Linn. *Syst. Nat.* 2. 855. 176.

*Roef. Inf.* 3. tab. 50.

Habitat in Grossularia.

*Larva* nigra, linea laterali alba.

*Puppa* brunnea postice obtuse tridentata.

*diffinis*. 105. N. cristata, alis deflexis ferrugineis, maculis tribus costalibus albis, postice punctis nigris duobus. *Syst. Ent.* 609. 82.

*Phalaena diffinis*. Linn. *Syst. Nat.* 2. 848. 146.

*Harris Inf. angl.* tab. 5. fig. 8.

Habitat in Europae, Angliae Vlmo.

*affinis*. 106. N. cristata, alis deflexis ferrugineis, angulo inferiori puncto nigro gemino, posticis nigris. *Syst. Ent.* 609. 83. \*

*Phalaena affinis*. Linn. *Syst. Nat.* 2. 848. 144.

Habitat in Europae nemoribus.

*triptera*. 107. N. cristata, alis deflexis, maculis tribus longitudinalibus, rotundatis atomisque sulphureis adspersis. *Syst. Ent.* 609. 84. \*

*Phalaena triptera*. Linn. *Syst. Nat.* 2. 848. 145.

Habitat in Europae nemoribus.

*marginata*. 108. N. cristata, alis deflexis flavescentibus, strigis ferrugineis postice fuscis. *Syst. Ent.* 610. 85. \*

Habitat in Europa.

*Absinthii*. 109. N. cristata, alis deflexis canis, fasciis punctisque in tetragonum positis nigris. *Syst. Ent.* 610. 86.

*Phalaena Absinthii* spirilinguis cristata, alis cinerascens nigris fasciatis punctatisque, inferioribus albidis. Linn. *Syst. Nat.* 2. 845. 133. *Fs. Sv.* 112.

*Phalaena seticornis* spirilingula, alis deflexis albidocinereis, lineis longis nigris. *Geoff. Inf.* 2. 158. 95.

*Clerk. Icon.* tab. 8. fig. 19.

*Frisch. Inf.* 7. tab. 12.

*Roef. Inf.* 1. phal. 2. tab. 61.

Habitat in Absinthio Europae.

*Larva* viridis, maculis flavis rufisque lineata.

*Puppa* folliculata brunnea antice viridis.

10. *N. cristata*, alis deflexis fuliginosis, arcis duabus cinescentibus, priore puncto marginali nigro. *Alm.*  
*Phalaena Alm.* Linn. *Syst. Nat.* 2. 845. 134.  
*Degeer Inf.* 1. tab. 2. fig. 25 — 8.  
*Naturf.* 12. tab. 1. fig. 14. 15. 16.  
*Fuesly Magaz.* 2. tab. 1. fig. 5 — 8.  
Habitat in Alno Europae.  
*Larva* nigra maculis dorsalibus flavis pilisque clavatis.  
*Puppa* folliculata, brunnea.
11. *N. cristata*, alis deflexis purpurascentibus, fasciis duabus flavis, posticis obscuris. *Syst. Ent.* 610. 87.  
*Phalaena Delphinii.* Linn. *Syst. Nat.* 2. 857. 188.  
*Phalaena feticornis spirilinguis*, alis deflexis roseis, superioribus fascia duplici limboque albidis. *Geoff. Inf.* 2. 164. 109.  
*Merian. Europ.* 1. tab. 40.  
*Wilks pap.* 3. tab. 1. a. 4.  
*Roef. Inf.* 1. phal. 2. tab. 12.  
Habitat in Europae Delphinio.  
*Larva* laevis, nuda, flauescens, nigro punctata lineisque duabus flavicantibus.  
*Puppa* nuda, brunnea.
12. *N. cristata*, alis deflexis ferrugineis bimaculatis, striga postica vndata pallida. *Syst. Ent.* 610. 88. *Pisi.*  
*Phalaena Pisi.* Linn. *Syst. Nat.* 2. 854. 172. *Sn. Sv.* 1206.  
*Phalaena* antennis filiformibus cristata, alis deflexis ferrugineis, strigis brunneis nigris griseisque strigae marginali vndata alba. *Degeer Inf. Versf. Germ.* 2. 1. 322. 10.  
*Raj. Ins.* 160. 10.  
*Wilks pap.* 4. tab. 1. a. 7.  
*Roef. Inf.* 1. phal. 2. tab. 52.  
*Merian. Europ.* tab. 50.  
Habitat in Piso, Genista.  
*Larva* nuda, ferruginea, lineis quatuor flavis, capite incarnato.  
*Puppa* fusca, incisuris sanguineis.
13. *N. cristata*, alis deflexis pallide incarnatis, macula media fusca, stigmatibus flavis. *Syst. Ent.* 611. 89. \* *mendica.*  
Habitat in Germania, *Dom. de Hattorf.*

232. CLASS. VI. GLOSSATA. Noctua.

*Oxyacanthae*. 114. N. cristata, alis deflexis bimaculatis, margine tenuiore coerulescente. *Syst. Ent.* 611. 90.

*Phalaena Oxyacanthae*. *Linn. Syst. Nat.* 2. 833. 165.  
*Fn. Sv.* 1207.

*Albin. Inf.* tab. 14.

*Wilk. pap.* 12. tab. 1. c. 1.

*Roef. Inf.* 1. phal. 2. tab. 33.

Habitat in Crataego Oxyacantha, Atriplice, Pruno spinosa.

*Larua* albo nigroque varia, cauda gibba.

*Puppa* folliculata, flava.

*Atriplicis*. 115. N. cristata, alis deflexis, anticis fusco nebulosis, liture media flava bifida. *Syst. Ent.* 611. 91.

*Phalaena Atriplicis*. *Linn. Syst. Nat.* 2. 854. 173. *Fn. Sv.* 1196.

*Phalaena feticornis*, spirilinguis, alis deflexis nebulosis fascia vna alteraue aurea. *Geoff. Inf.* 2. 159. 97.

*Wilk. pap.* 3. tab. 1. 2. 2.

*Roef. Inf.* 1. phal. 2. tab. 31.

Habitat in Atriplice, Rumice.

*Larua* nuda rufescens, albo punctata lineaque dorsali fusca.

*Puppa* nuda, brunnea.

*sinuata*. 116. N. cristata, alis deflexis basi cinereis, macula magna sinuata nigra. *Syst. Ent.* 611. 92. \*

Habitat Tranquebariae. *Mus. Tottianum*.

*histrionica*. 117. N. cristata, alis deflexis variegatis, posticis albis, striga marginali nigra. *Syst. Ent.* 612. 93. \*

Habitat Tranquebariae. *D. Koenig*.

*praecox*. 118. N. cristata, alis deflexis cinereis bimaculatis, posticis fascia rufa abbreviata. *Syst. Ent.* 612. 94.

*Phalaena praecox*. *Linn. Syst. Nat.* 2. 854. 174.

*Roef. Inf.* 1. phal. 2. tab. 51.

Habitat in Soncho oleraceo.

*Larua* nuda grisea, linea dorsali alba.

*Puppa* brunnea.

*pyramidea*. 119. N. cristata, alis fuscis, strigis tribus vndatis flavescens, strigibus repandis, posticis ferrugineis. *Syst. Ent.* 612. 95.

*Phalaena pyramidea* spirilinguis cristata, alis cinerescens, superioribus strigis tribus vndatis flavescens.

scitentibus repandis maculaque fusca, *Linn. Syst. Nat.* 2. 856. 181.

*Phalaena seticornis spirilinguis*, alis deflexis, superioribus fuscis, lineis transversis undulatis nigris, inferioribus ferrugineis. *Geoff. Inf.* 2. 160. 99.

*Phalaena media*, alis longis angustis, exterioribus lineolis et areolis nigris, albis atro rubentibus transversis pulcre depictis, interioribus obscure rubris. *Raj. Inf.* 159. 9.

*Merian. Europ.* 2. 23. tab. 9.

*Reaum. Inf.* 1. tab. 15. fig. 1 — 5.

*Ammiral. Inf.* tab. 11.

*Roef. Inf.* 1. phal. 2. tab. 11.

Habitat in Pruno, Iuglande, Quercu.

*Larva* nuda viridis, linea dorsali lateralibusque albis, postice conico gibba.

*Eruca* viridis, glabra, laevis, lineis albis secundum longitudinem ductis varia, angulosa, in extremo dorso eminentia dotata. *Raj. Inf.* 159. 9.

*Puppa* brunnea.

10. *N. cristata*, alis deflexis fuscis, stigmatibus marginalibus strigisque pallidis reticulatis. *Syst. Ent.* 613. 96. *typica.*

*Phalaena typica*. *Linn. Syst. Nat.* 2. 857. 186. *Fn. Sv.* 1195.

*Roef. Inf.* 1. phal. 2. tab. 5. 6.

Habitat in Europae Salice.

*Larva* nuda grisea, lateribus albis, linea nigra.

*Puppa* folliculata, brunnea.

11. *N. cristata*, alis deflexis cinereo nitidis, fascia media lucipara lata fusca.

*Phalaena lucipara spirilinguis cristata*, alis purpureis lucidis, fascia nigra, stigmate postico flavo.

*Linn. Syst. Nat.* 2. 857. 187. *Fn. Sv.* 1201.

Habitat in Suecia, Europa boreali.

*Rustica minor*. Alae totae cinereae, nitidae, in medio fascia late fusca, quae tamen marginem tenuiorem haud attingit.

12. *N. cristata*, alis deflexis, anticis ferrugineis, lunula lutescente strigaeque alba, postice bidentato. *Syst. Ent.* 613. 97. *oleracea.*

334 CLASS. VI. GLOSSATA. Noctua.

*Phalaena oleracea.* Linn. *Syst. Nat.* 2. 853. 171. *Fn. Sv.* 1197.

*Phalaena* alis exterioribus atro rubentibus cum lineola alba prope imum marginem transuersa, duos cuspidēs ad ipsum marginem emittente. *Raj. Inf.* 166. 21.

*Phalaena* antennis setaceis, alis deflexis ferrugineis macula alba flauaque lineaque alba vndata. *Deger Inf. Vers. Germ.* 2. 1. 305. 12.

*Frisch. Inf.* 7. tab. 21.

*Ammiral. Inf.* tab. 7.

*Roef. Inf.* 1. phal. 2. tab. 32.

*Reaum. Inf.* 2. tab. 39. fig. 4.

*Alb. Inf.* tab. 27.

Habitat in oleribus polyphaga.

*Larua* nuda, liuida, nigro punctata, linea dorsali fusca laterali alba.

*Puppa* nuda, nigricans.

*leucomelas.* 123. *N.* cristata, alis deflexis atris, posticis basi niuis. *Syst. Ent.* 613. 98.

*Phalaena leucomelas.* Linn. *Syst. Nat.* 2. 856. 183. *Fn. Sv.* 1184.

*Phalaena* seticornis spirilinguis, alis deflexis vndulato nigris, inferioribus basi albis. *Geoff. Inf.* 2. 149. 10.

*Clerk. Icon.* tab. 1. fig. 2.

Habitat in Europae hortis.

*vinularis.* 124. *N.* cristata, alis deflexis fuscis, fascia grisea, apice bifida flauo marginata. *Syst. Ent.* 613. 99.

Habitat in Saxoniae Coryletis.

*aprilina.* 125. *N.* cristata, alis deflexis, anticis virentibus, maculis variis atris, postice vtrinque punctis trigonis. *Syst. Ent.* 614. 100.

*Phalaena aprilina.* Linn. *Syst. Nat.* 2. 847. 138. *Fn. Sv.* 1178.

*Phalaena* media pulcherrima, alis oblongis, exterioribus dilute coeruleo viridibus maculis et lineolis nigris perbelle depictis. *Raj. Inf.* 158. 4.

*Phalaena* antennis filiformibus cristata, alis albo nigro viridique variis, macula ovali ferruginea annulo albo nigroque. *Deger Inf. Vers. Germ.* 2. 1. 317. 6.

*Deger*

*Degeer Inf.* 1. tab. 5. fig. 22. 23.

*Sulz. Hist. Inf.* tab. 22. fig. 8.

*Roef. Inf.* 3. tab. 39. fig. 4.

Habitat in Quercu.

*Larua* cinereo brunneoque variegata, linea dorsali alba interrupta, segmentoque singulo puncto albo.

*Puppa* brunnea.

16. *N. cristata*, alis deflexis, anticis abdomineque trifariam nigro punctato flavis. *Syst. Ent.* 614. 101.

*Phalaena ludifica. Linn. Syst. Nat.* 2. 848. 143. *Fn. Sv.* 1177.

*Schaeff. Icon.* tab. 92. fig. 3.

*Sepp. Inf.* 4. tab. 9.

Habitat in Europae nemoribus.

Thorax maculatus.

7. *N. cristata*, alis deflexis, anticis viridibus, maculis variis atris, subtus fuscis. *Syst. Ent.* 614. 102.\*

Habitat in Germaniae Lichenibus.

*Triplo minor N. aprilina.*

8. *N. cristata*, alis deflexis, anticis fuscis, fasciis duabus viridibus. *Syst. Ent.* 614. 103.\*

Habitat in Lichene saxatili Germaniae.

9. *N. cristata*, alis deflexis cinereis, anticis lineola bascos caracteribusque nigris. *Syst. Ent.* 614. 104. *Pfi.*

*Phalaena Pfi. Linn. Syst. Nat.* 2. 846. 135. *Fn. Sv.* 1181.

*Gsoff. Inf.* 2. 155. 91.

*Goed. Inf.* 1. tab. 22.

*Merian. Europ.* 3. tab. 42.

*Alb. Inf.* tab. 86.

*Frisch. Inf.* 2. tab. 2.

*Reaum. Inf.* 1. tab. 42. fig. 5. 6.

*Roef. Inf.* 1. phak. 2. tab. 7. 8.

*Anmir. Inf.* tab. 13.

*Wilk. pap.* 28. tab. 3. 6. 4.

Habitat in Pomonae arboribus, Alno, Quercu, polyphaga.

*Larua*



236 CLASS. VI. GLOSSATA. Noctua.

*Larva* pilosa, dorso flavo, lateribus nigris rubro maculatis, thoracis cornu erecto nigro.

*Puppa* folliculata, brunnea.

*Imago* numero et figura characterum nigrorum variat.

*Chi.* 130. *N. cristata*, alis deflexis canis, anticis x nigro notatis. *Syst. Ent.* 615. 105.

*Phalaena Chi. Linn. Syst. Nat.* 2. 846. 136. *Fn. Sv.* 1180.

*Phalaena feticornis spirilinguis*, alis deflexis; superioribus cinereo fuscoque nebulosis lincis vndulatis, et omicro nigris, inferioribus cinereis. *Geoff. Inf.* 2. 156. 93.

*Alb. Inf.* tab. 83. fig. C. D.

*Roef. Inf.* 1. phal. 2. tab. 13.

Habitat in Aquilegia, Soncho.

*Larva* nuda viridis, lincis lateralibus duabus albis.

*Puppa* folliculata, fusca.

*litura.* 131. *N. cristata*, alis deflexis canis, litura media atr puncto albo.

*Phalaena litura spirilinguis cristata*, alis canis, fuscis fusca liturisque tribus marginalibus nigra. *Linn. Syst. Nat.* 2. 858. 192. *Fn. Sv.* 1213.

Habitat in Suecia.

*Media.* Thorax cinereus, fusco varius. Alae canae puncto nigro distincto baseos. In medio macula vel potius litura magna nigra puncto albo. Postice puncta aliquot nigra.

*Persica- riae.* 132. *N. cristata*, alis deflexis fusco nebulosis, stigmate reniformi albo, pupilla lunari flava. *Syst. Ent.* 615. 106.

*Phalaena Persicariae. Linn. Syst. Nat.* 2. 847. 142. *Fn. Sv.* 1268.

*Phalaena feticornis spirilinguis*, alis deflexis, superioribus fuscis lineis vndulatis et omicro nigris, inferioribus cinereis. *Geoff. Inf.* 2. 157. 94.

*Ammir. Inf.* tab. 157.

*Roef. Inf.* 1. phal. 2. tab. 30.

Habitat in Pomonae arboribus polyphaga.

*Larva*

*Larva* nuda viridis, linea dorsali alba, secundo segmento macula quadrata atra, cauda conica.

*Puppa* brunnea.

33. N. cristata, alis deflexis, anticis fuscis, punctis nigris tribus approximatis, posticis liuidis. *Syst. Ent. g. 615. 107.*

*Phalaena Tragopoginis. Linn. Syst. Nat. 2. 855. 177. Fn. Sv. 1189.*

*Phalaena* antennis filiformibus, alis deflexis fuscis nitidis, punctis tribus centralibus nigris, capite flavo.

*Degeer Inf. Verf. Germ. 2. 1. 303. 10. tab. 7. fig. 15.*

*Frisch. Inf. 2. 33. tab. 7.*

Habitat in Tragopogine, Spinaehia, Rumice, Serratula aruensi.

*Larva* viridis, lineis sex albis stigmatibusque albis.

*Puppa* folliculata fusca.

34. N. cristata, alis deflexis obscure griseis, macula fusca punctisque tribus costalibus albis. *hepatica.*

*Phalaena hepatica* spirilinguis cristata, alis glaucescentibus, fascia ferruginea abbreviata terminalique plicata. *Linn. Syst. Nat. 2. 853. 169. Fn. Sv. 1209.*

*Phalaena hepatica. Clerk. Icon. tab. 8. fig. 3.*

Habitat in Suecia.

*Rustica media.* Corpus totum griseum ferrugineo aut fusco mixtum. Macula magna transversalis fusca in medio marginis crassioris et puncta tria parua versus apicem.

5. N. cristata, alis deflexis flavescentibus, strigis duabus fuscis lunulaque lutea. *Syst. Ent. 616. 108.\* turca.*

*Phalaena turca. Linn. Syst. Nat. 2. 847. 140. Fn. Sv. 1221.*

Habitat in Europa boreali.

6. N. cristata, alis deflexis dentatis nebulosis, striga postica dentata alba. *Syst. Ent. 616. 109.\* polyodon.*

*Phalaena polyodon* spirilinguis cristata, cinereo nebulosa, margine postico multidentato. *Linn. Syst. Nat. 2. 853. 170. Fn. Sv. 1219.*

Habitat in Europa boreali.

- Eremita.* 137. N. cristata, alis deflexis ferrugineis, strigis duabus vndatis albis, anteriore obsoleta, posteriore repanda. *Syst. Ent.* 616. 110.\*  
Habitat Lipsiae.
- nigricans.* 138. N. cristata, alis deflexis nigricantibus, stigmatibus ordinariis pallidioribus. *Syst. Ent.* 616. 111.  
*Phalaena nigricans.* Linn. *Syst. Nat.* 2. 855. 178. Fn. Sv. 1220.  
Habitat in Europa boreali.
- oculea.* 139. N. cristata, alis deflexis fusco ferrugineoque varis, stigmate reniformi pallido. *Syst. Ent.* 616. 112.\*  
Habitat in Sueciae nemoribus.
- flavicornis.* 140. N. cristata, alis deflexis, anticis cinereis, strigibus atris, antennis luteis. *Syst. Ent.* 617. 113.  
*Phalaena flavicornis.* Linn. *Syst. Nat.* 2. 856. 179. Fn. Sv. 1204.  
Clerk. *Icon.* tab. 6. fig. 9.  
Habitat in Pomonae arboribus.  
*Larva* nigra, albo punctata, capite brunneo.
- triplicata.* 141. N. cristata, alis deflexis, anticis arcu duplici contrario maculisque tribus glaucis intermediis. *Syst. Ent.* 617. 114.  
*Phalaena triplicata.* Linn. *Syst. Nat.* 2. 854. 175. Fn. Sv. 1202.  
Degeer *Inf.* 1. tab. 6. fig. 20. 21.  
Merian. *Europ.* tab. 97.  
Roef. *Inf.* 1. phal. 2. tab. 34.  
Habitat in Soncho, Humulo, Vertica.  
*Larva* nuda, viridis, gibberibus 2 dorsis, at ani fusca.  
*Puppa* folliculata, brunnea.
- tristis.* 142. N. cristata, alis deflexis cinereis fusco subundatis, margineque exteriori punctato, subtus pallidis, fascia fusca. *Syst. Ent.* 617. 115.\*  
Habitat in Suecia.  
Macula ordinaria lunula obscuriore.
- Rumicis.* 143. N. cristata, alis deflexis cinereo fuscoque varis, latura marginis tenuioris alba.

*Phalaena Rumicis spirilinguis cristata*, alis deflexis cinereo bimaculatis, litura marginis tenuioris alba,  
Linn. Syst. Nat. 2. 852. 164. Fn. Sv. 1200.

Merian. Europ. tab. 82.

Alb. Inf. tab. 32.

Roef. Inf. 1. phal. 2. tab. 27.

Wilk. pap. 26. tab. 3. a. 1

Degeer Inf. 4. tab. 9. fig. 2.

Habitat in Rumice, Soncho.

*Larva* pilosa albo rubroque punctata, linea laterali flava.

*Puppa* brunnea.

4. *N. cristata*, alis lanceolatis conuolutis fusco cinereo-*exoleta*,  
que nebulosis, punctis quatuor marginalibus albis.  
Syst. Ent. 617. 116.

*Phalaena exoleta spirilinguis cristata*, alis lanceolatis conuolutis exoletis, dorso fuscescentibus, collari impresso. Linn. Syst. Nat. 2. 849. 151. Fn. Sv. 1185.

Frisch. Inf. 5. tab. 11. fig. 1.

Wilk. pap. 8. tab. 1. a. 18.

Roef. Inf. 1. phal. 2. tab. 24.

Degeer Inf. 2. tab. 7. fig. 4.

Sulz. Inf. tab. 16. fig. 95.

Schaeff. Icon. tab. 24. fig. 6. 7.

Habitat in Chinopodio, Atriplici, Oleribus.

*Larva* media viridis, punctata, linea laterali albida.

*Puppa* nuda, brunnea.

5. *N. cristata*, alis deflexis dentato eros, margine la-*Verbasci*.  
terali fusco immaculato. Syst. Ent. 618. 117.

*Phalaena verbasci*. Linn. Syst. Nat. 2. 850. 153. Fn. Sv. 1186.

*Phalaena feticornis spirilinguis*, alis deflexis fusco cinereis, superioribus fuscis longitudinaliter striatis.  
Goff. Inf. 2. 158. 96.

*Phalaena media*, alis exterioribus cinereis, lineis et maculis nigris varie depictis. Raj. Inf. 168. 25.

Merian. Europ. 3. tab. 29.

Frisch. Inf. 6. tab. 9.

Roef. Inf. 1. phal. 2. tab. 23.

Reaum. Inf. 1. tab. 43. fig. 9—11.

Wilk.

*Wilks pap.* 7. tab. 1. a. 15.

*Sulz. Hist. Inf.* tab. 22. fig. 7.

*Harr. Aurel.* tab. 8. fig. a. d.

Habitat in Verbasco, Scrophularia.

*Larva* nuda, cinerea, flavo nigroque punctata.

*Eruca* Verbascum depascens cinerea pulcre varieg.

*Raj. Inf.* 168. 25. et 352. 31.

*Puppa* folliculata lutea, stigmatibus nigris.

*umbratica.* 146. N. cristata, alis lanceolatis, striatis, canis, thoracalibus lunatis. *Syst. Ent.* 618. 118.

*Phalaena umbratica.* *Linn. Syst. Nat.* 2. 849. 150.

*Sv.* 1184.

*Roef. Inf.* 1. phal. 2. tab. 25.

Habitat in Soncho.

*Larva* nuda nigricans, trifariam rubro punctata, caudam acuminata.

*Puppa* folliculata, brunnea.

*putris.* 147. N. cristata, alis deflexis, obsoletis, subpunctatis, margine exteriori fusco adiecta macula subocellari. *Syst. Ent.* 618. 119.\*

*Phalaena putris.* *Linn. Syst. Nat.* 2. 850. 152. 1

*Sv.* 1187.

Habitat in Europa.

*rurea.* 148. N. cristata, alis deflexis, griseo fuscoque variis, posticis fuscis, margine crassiori albo. *Syst. Ent.* 618. 120.\*

Habitat in Angliae nemoribus.

*Myrtilli.* 149. N. cristata, alis deflexis, ferrugineis, albo maculatis posticis luteis, fascia lata nigra submarginali. *Syst. Ent.* 619. 121.

*Phalaena Myrtilli* cristata, alis griseis albo variis, inferioribus antice albis postice nigris. *Linn. Syst. Nat.* 2. 853. 167. *Fn. Sv.* 1168.

*Roef. Inf.* 4. tab. 11. fig. A. B. C.

Habitat in Vaccinio viginoso, Myrtilla.

*Larva* nuda viridis, dorso quinquetuberculato, capite coeruleo.

*Puppa* brunnea, stigmatibus albis.

1. *N. cristata*, alis deflexis fuscis, posticis nigris, fascia flava. *Syst. Ent.* 616. 122.\*

*Naturf.* 3. tab. 1. fig. 8.

Habitat in Angliae nemoribus.

Præcedente paullo minor.

## 148. PHALÆNA

\* Antennis pectinatis.

*P. bipectinicornis*, alis angulatis, niueis, immaculatis, *lactearia*.  
antennis apice setaceis. *Syst. Ent.* 619. 1. *Linn. Syst.*  
*Nat.* 2. 856. 194. *Fn. Sv.* 1226.

*Phalaena pectinicornis* spirilinguis, alis patentibus rotundatis niueis, corpore flavo. *Geoff. Inf.* 2. 131. 44.

*Phalaena lactearia*. *Scopol. carn.* 529.

Habitat in Europae nemoribus.

Tenerrima et fere pellucida.

*P. pectinicornis*, alis subangulatis albis, margine postico anticisque subtus fuscis, posticis puncto centrali nigro. *Syst. Ent.* 620. 2.\*

Habitat in Angliae nemoribus.

*P. pectinicornis*, alis angulatis virefcentibus, strigis duabus albis repandis, antennis apice setaceis. *Syst. Ent.* 620. 3. *Linn. Syst. Nat.* 2. 858. 195. *Fn. Sv.* 1227.

*Phalaena* antennis pectinatis apice setaceis, alis horizontalibus viridibus, fasciis duabus albidis, posticis angulatis. *Degeer Inf. Vers. Germ.* 2. 1. 263. 3. tab. 6. fig. 8.

*Rosf. Inf. phal.* 3. tab. 13.

*Reaum. Inf.* 2. tab. 29. fig. 14—19.

*Schaeff. Icon.* tab. 122. fig. 5.

Habitat in Iasmino, Syringa.

*Larva* ferruginea, albo nigroque varia, capite dentato.

*P. pectinicornis*, alis angulatis, dentatis, viridibus, striga undulata obsoleta alba. *Thymia-*

*Phalaena Thymiaria*. *Linn. Syst. Nat.* 2. 859. 199.

*Frisch. Inf.* 10. 20. tab. 17.

Habitat in Germaniae Thymo.

Dentes alarum ferruginei.

*Fabr. Spec. Inf. T. II.* Q

5. P.

242 CLASS. VI. GLOSSATA. Phalaena.

*putatoria*. 5. *P. peëtinicornis*, alis angulatis albis, strigis duabus candidis repandis, antennis apice setaceis. *Syst. Ent.* 620. 4. *Linn. Syst. Nat.* 2. 859. 196. *Fn. Sv.* 1225.  
*Clerk. Icon.* tab. 3. fig. 9.  
 Habitat in Europae nemoribus.

*friatatoria*. 6. *P. peëtinicornis*, alis angulatis albidis, strigis duabus flavescentibus obsoletis. *Syst. Ent.* 620. 5. *Linn. Syst. Nat.* 2. 859. 197.  
*Clerk. Icon.* tab. 55. fig. 4.  
 Habitat in Europae umbrifis.

*punctaria*. 7. *P. peëtinicornis*, alis angulatis cinereis, striga ferruginea ordineque punctorum nigrorum. *Syst. Ent.* 620. 6. *Linn. Syst. Nat.* 2. 859. 200. *Fn. Sv.* 1254.  
*Clerk. Icon.* tab. 5. fig. 11.  
 Habitat in Europa, Anglia.

*Dioraria*. 8. *P. peëtinicornis*, alis dentatis fufcis, strigis purpureis apice pallidis.  
*Phalaena Dioris. Cram. Inf.* 7. tab. 75. fig. F.  
 Habitat Surinami.  
 Strigae, quatuor vndatae et margo pofticus tenuiffime purpureae.

*amatoria*. 9. *P. peëtinicornis*, alis angulatis, omnibus pallidis, puluerulentis, fascia purpurea recta strigaeque fufca repanda. *Syst. Ent.* 621. 7. *Linn. Syst. Nat.* 2. 859. 201. *Fn. Sv.* 1223.  
*Phalaena peëtinicornis* elinguis, alis patentibus angulatis cinereis, fascia duplici transuerfa punctoque obfcuriore atomis cinerascens. *Geoff. Inf.* 2. 128. 7.  
*Wilks pap.* 37. tab. 1. a. 5.  
*Reaum. Inf.* 2. tab. 29. fig. 1—4.  
 Habitat in Europae Quercu.  
*Larva* viridis, cingulis fupra flauis, fubtus rubris.

*falcatoria*. 10. *P. peëtinicornis*, alis falcatis glaucis, anticis undis fasciaeque grifeis, puncto fufco. *Syst. Ent.* 621. 8. *Linn. Syst. Nat.* 2. 859. 202. *Fn. Sv.* 1224.  
*Degeer Inf.* 1. tab. 24. fig. 7.  
 — — — *Versf. Germ.* 2. 1. 257. 7. tab. 6. fig. 1.  
*Naturf.* 9. 97. tab. 1. fig. 6.  
*Schaeff. Icon.* tab. 64. fig. 1.

Habitat

Habitat in Betula alba, Alno.

Larua fusca, striis albis, quiescit capite caudaque eleuatis.

11. *P. pectinicornis*, alis subfalcatis luteis, fascia saturatiora, antennis apice setaceis. *Syst. Ent.* 621. 9.\*

Habitat in Germania.

12. *P. pectinicornis*, alis angulato caudatis albis, strigis fuscis, vndatis fuscis supra duabus, subtus vnica. *Syst. Ent.*

*App.* 831.\*

*Phalaena Geminea. Cram. Inf.* 12. tab. 133. fig. C.

Habitat in India orientali. *Dom. Edler.*

13. *P. pectinicornis*, alis caudato angulatis flauescens, strigis duabus obscurioribus, posticis apice bipunctatis. *Syst. Ent.* 621. 10. *Linn. Syst. Nat.* 2. 860. 203.

*Fn. Sv.* 1222.

*Phalaena feticornis spirilinguis*, alis patentibus sulphureis, linea duplici transuersa obscuriori, inferioribus caudatis. *Geoff. Inf.* 2. 138. 58.

*Phalaena media ochroleucos*, alis amplissimis, exterioribus duabus lineis transuersis, e fuluo virentibus, interioribus vna diuisis. *Raj. Inf.* 177. 1.

*Phalaena antennis filiformibus*, alis latis angulatis luteis, strigis duabus obscurioribus. *Degeer Inf. Vers.*

*Germ.* 2. 1. 327. 3.

*Alb. Inf.* tab. 94.

*Sepp. Inf.* 6. 1. tab. 1.

*Schaeff. Icon.* tab. 93. fig. 8.

*Roef. Inf.* 1. phal. 3. tab. 6.

*Goed. Inf.* 2. tab. 34.

*Petiv. Gazoph.* tab. 51. fig. 6.

*Wilks pap.* 38. tab. 1. 6. 2.

*Clerk. Icon.* tab. 50. fig. 2.

Habitat in Europae Sambuco.

Larua subnodosa, pulla, fusco lineata.

Puppa fusiformis, brunnea, fusco punctata.

Oua costata.

14. *P. pectinicornis*, alis erosius lutescentibus, strigis duabus fuscis. *Syst. Ent.* 622. 11. *lacertinaria.*

*Phalaena lacertinaria pectinicornis*, alis erosius lutescentibus postice saturatioribus, lineis duabus punctoque fuscis. *Linn. Syst. Nat.* 2. 860. 204. *Fn. Sv.* 1229.

Q 2

Pha-



*Phalaena* antennis pectinatis dorso laevi, alis patentibus dentatis lutescentibus, strigis duabus punctoque fuscis. *Degeer Inf.* 2. 1. 242. 3.

*Alb. Inf.* tab. 65.

*Degeer Inf.* 1. tab. 10. fig. 7. 8.

*Reaum. Inf.* 2. tab. 22. fig. 4. 6.

*Wilks pap.* 74. tab. 1. c. 3.

*Sulz. Hist. Inf.* tab. 22. fig. 10.

*Schaeff. Icon.* tab. 66. fig. 2. 3.

Habitat in Quercu, Betula.

*Larva* nuda, rubra, dorso angulato, cauda acuta.

*Puppa* folliculata, conica, brunnea, atomis albis.

*equestraria.* 15. *P. pectinicornis*, alis angulatis, anticis fuscescentibus, fascia lata flava, posticis flavis. *Gen. Inf. Mant.* 285.\*

Habitat Hamburgi. *D. Schulz.*

*alniaria.* 16. *P. pectinicornis*, alis erosis, flavis, fusco pulveratis, strigis duabus fuscis. *Syst. Ent.* 622. 12. *Linn. Syst. Nat.* 2. 860. 205. *Fn. Sv.* 1230.

*Phalaena* elinguis, antennis pectinatis, alis patentibus, dentatis, flavis, fasciis duabus obscuris, collari sulphureo. *Degeer Inf. Vers. Germ.* 2. 1. 219. 4.

*Degeer Inf.* 1. tab. 10. fig. 13. 14.

*Roef. Inf.* 1. phal. 3. tab. 1.

*Schaeff. Icon.* tab. 135. fig. 1. 2.

Habitat in Pomonae arboribus.

*Larva* grisea fusca, luteo punctata, verrucis dorsalibus tribus remotis, caudalibus quatuor approximatis.

*Puppa* coerulea, postice attenuata.

*syringaria.* 17. *P. pectinicornis*, alis suberosis, omnibus griseo lutescentibus, strigis repandis fuscis albisque. *Syst. Ent.* 622. 13. *Linn. Syst. Nat.* 2. 860. 206.

*Phalaena* pectinicornis, alis margine sinuatis, fuscae, nigro, fusco roseoque marmoratis, singulis subtus puncto nigro, superioribus extremo dilatato, recurvis. *Geoff. Inf.* 2. 126. 32.

*Roef. Inf.* 1. phal. 3. tab. 10.

*Wilks pap.* 39. tab. 1. 6. 6.

Habitat in Syringa, laesmino.

*Larva* glauco luteoque varia, dorso aculeis sex, posticis longioribus recurvis.

*Puppa*

CLASS. VI. GLOSSATA. Phalaena. 245

*Pappa folliculata*, farsa, antice grisea, postice brunnea.

3. *P. pectinicornis*, alis angulato dentatis basi rufis, lunula alba, postice cinereis. *Syst. Ent.* 623. 14.\*

*Phalaena lunaria*. *Wien. Verz.* 276. tab. 1. a. fig. 3.  
— — — — — tab. 1. 6. fig. 3.

*Sepp. Inf.* 6. 9. tab. 2.

Habitat in Germaniae Pyro, Salice, Betula.

*Larva* grisea, tuberculis dorsalibus didymis, subtus fasciis albidis nigro punctatis.

*Pappa* folliculata, obtusa brunnea.

4. *P. pectinicornis*, alis angulato dentatis supra pallidis, dentaria. strigis ferrugineis, subtus ferrugineis, lunula obscuriori. *Syst. Ent.* 623. 15.\*

Habitat in Anglia. *Mus. D. Monson.*

5. *P. bipectinicornis*, alis dentatis albis fusco undatis, undaria. antennis apice setaceis. *Syst. Ent.* 623. 16.\*

Habitat Tranquebariae. *D. Koenig.*

6. *P. pectinicornis*, alis angulatis flauis, strigis ferrugineis, angulo ani violaceo. *Syst. Ent.* 623. 17. *dolabraria.*

*Phalaena dolabraria*, pectinicornis alis subflavescentibus, strigis subferrugineis sparsis, lobo postico violaceo. *Linn. Syst. Nat.* 2. 361. 207.

*Sulz. Hist. Inf.* tab. 22. fig. 9.

Habitat in Quercu Germaniae, Angliae.

7. *P. pectinicornis*, alis angulatis viridibus, anticis strigis tribus fuscis. *Gen. Inf. Mant.* 285.\*

*Phalaena Ilyrias*. *Cram. Inf.* 1. tab. 10. fig. E.

Habitat Surinami.

8. *P. pectinicornis*, alis dentatis viridibus, striga punctorum nigrorum, posticis puncto duplici albo. *monilaria.*  
*Gen. Inf. Mant.* 286.\*

Habitat in India. *D. Fothergill.*

9. *P. pectinicornis*, alis anticis griseis, posticis ferrugineis. *Rondelaria.*  
*Syst. Ent.* 623. 18.\*

Habitat in Rondeletia Indiae orientalis.

246 CLASS. VI. GLOSSATA. *Phalaena*:

*vibicaria*. 25. *P. pectinicornis*, alis rotundatis flavescentibus, strigis tribus purpureis, anteriore obsoletiore. *Syst. Ent.* 623. 19.

*Phalaena vibicaria pectinicornis*, alis angulatis, glaucescentibus, puncto, fascia strigaeque purpureis. *Linn. Syst. Nat.* 2. 859. 198. *Fn. Sv.* 1228.

Habitat in Europae nemoribus frequens.

Alae in nostris haud angulatae.

*papilio-naria*. 26. *P. pectinicornis*, alis viridibus subrepandis, striga setualltera repanda. *Syst. Ent.* 624. 20. *Linn. Syst. Nat.* 2. 864. 225. *Fn. Sv.* 1247.

*Frisch. Inf.* 10. tab. 17.

*Roej. Inf.* 4. tab. 12. fig. 3.

Habitat in Europae Betulia.

*Larva* viridis, dorso aculeis decem incurvis rufis.

*Naturf.* 12. tab. 1. fig. 11. 12. 13.

*Puppa* viridi flavoque variegata.

*Quies* 14 dierum.

*macul-aria*. 27. *P. pectinicornis*, alis fulvis apice nigris, fascia maculari alba.

*Papilio Helcita* D. C. alis integerrimis, rotundatis, luteis, concoloribus, limbo nigro, punctis septem albis.

*Syst. Ent.* 475. 141. *Linn. Syst. Nat.* 2. 763. 94.

*Amoen. acad.* 6. 405. 62.

*Phalaena Helcita*. *Cram. Icon.* tab. 39. fig. 4.

*Phalaena fusulis*. *Linn. Mus. Lud. Vr.* 390. 25.

Habitat in India. *Mus. Dom. Teats.*

Magna. Antennae maris pectinatae, feminae setaceae, nigrae. Corpus supra nigrum, albo maculatum triplici serie, subtus fulvum. Alae omnes fulvae apice nigrae, fascia e punctis sex albis.

*catenaria*. 28. *P. pectinicornis*, alis rotundatis albis, lunula media strigaeque postica undata nigra.

*Phalaena catenaria*. *Drury Inf.* 1. tab. 8. fig. 3.

Habitat — — — —

Frons fulva. Punctum fulvum utrinque ante alas.

Alae omnes albae.

Variet interdum alis tantum nigro punctatis.

*pyrrharia*. 29. *P. pectinicornis*, alis subdentatis luteis fusco polverulentis, anticis lunula fusca. *Syst. Ent.* 624. 21.

*Phala-*

*Phalaena prunaria*. Linn. *Syst. Nat.* 2. 861. 208. *Fn.*  
*Sv.* 1232.

*Albin. Inf.* 42. tab. 69.

*Wilks pap.* 40. tab. 1. b. a.

*Schaeff. Icon.* tab. 17. fig. 2. 3.

Habitat in Pruno spinosa.

*Larua* ferruginea, dorso antice posticeque spinis duabus.

30. *P. pectinicornis*, alis griseo nebulosis, omnibus subtus *furcaria*.  
puncto nigro, lunula alba. *Syst. Ent.* 624. 22. \*

Habitat in Germaniae nemoribus.

31. *P. pectinicornis*, alis fuscis, flauo maculatis, subtus *pinaria*.  
nebulosis, fasciis duabus fuscis. *Syst. Ent.* 624. 23.

*Phalaena pinaria* pectinicornis, alis subtus griseis line  
nea alba interrupta fasciisque duabus obscurioribus.

*Linn. Syst. Nat.* 2. 861. 210. *Fn. Sv.* 1233.

*Phalaena* antennis pectinatis, alis horizontalibus, ferru  
gineis, oliuaceo irroratis. *Degeer Inf. Verf. Germ.*

2. 1. 255. 5. tab. 5. fig. 20.

*Clerk. phal.* tab. 1. fig. 10.

*Ammiral. Inf.* tab. 20.

*Sepp. Inf.* 6. 17. tab. 4.

*Schaeff. Icon.* tab. 159. fig. 1. 2.

*β. Clerk. Icon.* tab. 4. fig. 11.

*Reaum. Inf.* 2. tab. 4. fig. 6.

Habitat in Pino, Tilia, Betula.

*Larua* viridis, supra albo, subtus flauo striata.

*Puppa* folliculata, brunnea.

32. *P. pectinicornis*, alis ferrugineis, limbo nigro, posticis *limbaria*.  
subtus nigris albo striatis. *Syst. Ent.* 624. 24. \*

*Harris Inf. angl.* tab. 5. fig. 4.

Habitat in Angliae nemoribus. *Mus. Dom. Banks.*

33. *P. pectinicornis*, alis flauis fusco irroratis, anticis apice *roraria*.  
nigris. *Gen. Inf. Mant.* 285. \*

Habitat in Europa boreali.

*Variat* rarius absque margine nigro alae anticae.

34. *P. pectinicornis*, alis nigro punctatis maculatisque, an- *melana-*  
ticis albidis, posticis lutela. *Syst. Ent.* 625. 25. *Linn. ria.*

*Syst. Nat.* 2. 862. 212. *Fn. Sv.* 1240.

*Clerk. phal.* tab. 4. fig. 2.

Habitat in Europa boreali.

248 CLASS. VI. GLOSSATA. Phalaena.

*atomaria*. 35. *P. pectinicornis*, alis omnibus lutescentibus, strigis atomisque fuscis. *Syst. Ent.* 625. 26. *Linna. Syst. Nat.* 2. 862. 214. *Fn. Sv.* 1245.

*Phalaena* antennis pectinatis, alis horizontalibus dentatis luteis, strigis atomisque fuscis. *Degeer Inf. Verf. Germ.* 2. 1. 249. 2. tab. 5. fig. 14.

*Phalaena isofelata*. *Scop. carn.* 558. ♀

*Phalaena pennata*. *Scop. carn.* 569. ♂

*Geoff. Inf.* 2. 133. 50.

*Frisch. Inf.* 13. tab. 5.

*Albin. Inf.* 42. fig. 69.

*Kleman. Inf.* 1. tab. 34. mal.

Habitat in Europae Tilia.

*Larva* lacuis grisea, lineis numerosis ferrugineis interruptis, postice tuberculis duobus.

*profapia*. 36. *P. pectinicornis*, alis lutescentibus, atomis strigisque tribus fuscis, tertia maculari. *Gen. Inf. Mant.* 285. ♂

*Phalaena profapiaria* pectinicornis, alis lutescentibus, fasciisque vtrinque duabus saturatoribus. *Linna. Syst. Nat.* 2. 864. 222. *Fn. Sv.* 1252.

*Vdm. Dissert.* 66.

*Clerk. phal.* tab. 3. fig. 1.

Habitat in Europae borealis nemoribus.

*pufaria*. 37. *P. pectinicornis*, alis niueis, strigis tribus fuscis obsoletis. *Syst. Ent.* 625. 27. *Linna. Syst. Nat.* 2. 864. 223. *Fn. Sv.* 1251.

*Phalaena* antennis filiformibus, alis latis albis, strigis anticarum tribus, posticarum duabus cinereis. *Degeer Inf. Verf. Germ.* 2. 1. 329. 4. tab. 8. fig. 12.

*Phalaena frigata*. *Scop. carn.* 541.

*Clerk. phal.* tab. 3. fig. 1.

*Schaeff. Icon.* tab. 111. fig. 7.

*Pontop. Atl. Dan.* 1. tab. 30.

Habitat in Europae Betula, Alno.

*Larva* flavescentes, lineis maculisque dorsalibus rubris.

*ocularia*. 38. *P. pectinicornis*, alis rotundatis flauis, omnibus puncto ocellari flauo. *Syst. Ent.* 625. 28. \*

Habitat in India. *Mus. Tottianum*.

Antennae valde pectinatae.

39. *P. pectinicornis*, alis cinereis, striga annulo fasciisque duabus fuscis. *Syst. Ent.* 625. 29.\* *annularia.*  
*Phalaena annulata.* *Naturf.* 6. tab. 4. fig. 3.  
 Habitat in Germaniae hortis.
40. *P. pectinicornis*, alis griseis, atomis fuscis, medio albis, puncto fusco, femina aptera nigro punctata. *defoliaria.*  
*Syst. Ent.* 625. 30. *Linn. Fn. Sv.* 1238.  
*Phalaena defoliaria* alis griseis medio albis, puncto lituraque postica nigris, femina aptera. *Linn. Syst. Nat.* 3. 225.  
*Clerk. phal.* tab. 7. fig. 4.  
*Reaum. Inf.* 2. tab. 30.  
 Habitat in Ceraso, Tilia.  
*Larva* ferruginea, stria laterali sulphurea.  
*Puppa* brunnea.
41. *P. pectinicornis*, alis cinereis, striga fusca vndata albae innata. *Gen. Inf. Mant.* 285.\* *hirtaria.*  
*Phalaena hirtaria* pectinicornis, alis hirtis canis, strigis tribus nigris, posterioribus approximatis, antennis atris. *Linn. Fn. Sv.* 1235.  
*Clerk. Icon.* tab. 7. fig. 1.  
*Degeer Inf.* 1. tab. 22. fig. 6.  
 Habitat in Europae pratis syluaticis.  
*Larva* geometra ferruginea, lineis longitudinalibus palide rufis nigro marginatis.
42. *P. pectinicornis*, alis flavescentibus, strigis duabus, posteriore limbum obscurum determinante. *Syst. Ent.* 626. 31. *vespertina.*  
*Linn. Syst. Nat.* 2. 864. 224.  
 Habitat in Germania, Gallia.
43. *P. pectinicornis*, alis cinereis, anticis fasciis quatuor nigris abbreviatis inaequalibus. *Syst. Ent.* 626. 32. *Wauaria.*  
*Linn. Syst. Nat.* 2. 863. 219. *Fn. Sv.* 1248.  
*Phalaena minor*, alis amplis cinereo albicantibus 4 in exteriorum margine maculis linearibus e rufo nigricantibus. *Raj. Inf.* 179.  
*Petiv. Mus.* 3. n. 4.  
*Albin. Inf.* 47. fig. 78.  
*Merian. Europ.* 1. tab. 25. fig. 151.  
*Frisch. Inf.* 3. tab. 3. fig. 1.

250 CLASS. VI. GLOSSATA. Phalaena.

*Roef. Inf.* 1. phal. 3. tab. 4.

*Wilks pap.* 52. tab. 2. a. 2.

*Ammiral. Inf.* tab. 13. fig. 2. 3.

Habitat in Ribe, Grossularia.

*Larua* subpilosa, viridis, nigro punctata, linea dorsali lateralibusque flavis.

*Puppa* nuda, brunnea, acuminata.

- prafinaria.* 44. *P. pectinicornis*, alis viridibus, strigis tribus, anticis obscurioribus, postica pallida. *Syst. Ent.* 626. 33.\*  
Habitat in Anglia. *Mus. Dom. Banks.*

- ditaria.* 45. *P. pectinicornis*, alis viridibus, maculis marginalibus ferrugineis. *Gen. Inf. Mant.* 285.\*  
Habitat Hamburgi. *D. Schulz.*

- viridaria.* 46. *P. pectinicornis*, alis rotundatis viridibus albo strigis maculisque marginalibus atris. *Syst. Ent.* 626. 34.\*  
Habitat in Angliae nemoribus. *Mus. Dom. Banks.*

- tripunctaria.* 47. *P. pectinicornis*, alis fuscis, anticis fascia maculari, posticis disco albis. *Syst. Ent.* 627. 35.

*Phalaena tripunctaria pectinicornis*, alis fuscescentibus anterioribus fascia interrupta bascosque striga posterioribus disco albis. *Linn. Syst. Nat.* 2. 364. 226. *Mus. Lud. Vlr.* 392.

*Cram. Inf.* 2. tab. 22. fig. E.

*Petiv. Gazoph.* tab. 28. fig. 8.

Habitat in Asia.

Conf. *Phalaena Luficinia*. *Cram. Inf.* 11. tab. 121. fig. E.

*Edw. Av.* 35. tab. 35.

Vix specie differunt.

- Iatropharia.* 48. *P. pectinicornis*, alis concoloribus nigris, anticis maculis duabus, posticis una flavis. *Syst. Ent.* 627. 36.

*Linn. Syst. Nat.* 2. 365. 229. *Mus. Lud. Vlr.* 395.

*Phalaena Osiris*. *Cram. Inf.* 10. tab. 115. fig. E.

*Clerk. Icon.* tab. 55. fig. 3.

*Merian. Surin.* tab. 38. fig. 1.

Habitat in Americae *Iatrophae Gossypifoliae*.

- Catilaria.* 49. *P. subpectinicornis*, alis concoloribus nigris, fasciis duabus virefcentibus strigaeque maculari alba,

*Pha-*

CLASS. VI. GLOSSATA. Phalaena. 251

*Phalaena Catilina. Cram. Inf. 7. tab. 79. fig. E. F.*  
Habitat Surinami.

Affinis praecedenti. Striga alba versus marginem e punctis septem constat. Thorax albo punctatus.

50. *P. pectinicornis*, alis concoloribus albis nigro variis, *Cenaria*.  
corpore flauo nigro maculato.

*Phalaena Cenis. Cram. Inf. 13. tab. 147. fig. E.*  
Habitat in China.

Media. Alae anticae basi albae nigro maculae apice nigrae punctis quinque albis. Posticae albae maculis tribus nigris submarginalibus. Thorax flauus nigro punctatus. Abdomen flauum cingulis sex punctisque tribus apicis nigris.

51. *P. pectinicornis*, alis rotundatis atris coeruleo strigatis maculatisque. *Syst. Ent. 627. 37. \** *fenestria.*  
Habitat in China. *Mus. Dom. Banks.*

52. *P. pectinicornis*, alis luteis, margine venisque dilatatis ramosi nigris. *Syst. Ent. 627. 38. \** *venaria.*

*Phalaena papilionaris. Drury Inf. 2. tab. 2. fig. 4.*

*Phalaena papilionaris. Cram. Inf. 3. tab. 29. fig. A.*

Habitat in India orientali.

53. *P. pectinicornis*, alis omnibus testaceo puluerulentis, *puluerafascia lata ferruginea. Syst. Ent. 627. 39. Linn. Syst. Nat. 2. 862. 215. Fn. Sv. 1243. ria.*

*Clerk. phal. tab. 5. fig. 6.*

Habitat in Europae borealis aruis.

54. *P. pectinicornis*, alis omnibus rufescentibus, *fascia fasciaria. lata ferruginea, margine albo. Syst. Ent. 628. 40.*

*Linn. Syst. Nat. 2. 862. 216. Fn. Sv. 1242.*

*Clerk. Icon. tab. 5. fig. 5.*

Habitat in Europae borealis ericetis.

55. *P. pectinicornis*, alis albidis puluerulentis, 'omnibus *marginastriga marginali punctorum nigrorum. Gen. Inf. Mant. 286. \** *ria.*

Habitat Hamburgi. *D. Schulz.*



*betularia*. 56. *P. pectinicornis*, alis omnibus albis, atomis nigris, thorace fascia nigra, antennis apice setaceis. *Syst. Ent.* 628. 41. *Linn. Syst. Nat.* 2. 862. 217. *Fn. Sv.* 1287.

*Phalaena* antennis pectinatis, alis horizontalibus albis nigro punctatis maculatisque, thorace fasciato.

*Degeer Inf. Vers. Germ.* 2. 1. 250. 1. tab. 5. fig. 11.

*Petiv. Gazoph.* tab. 7. fig. 2.

*Albin. Inf.* tab. 91. 92.

*Wilks pap.* 38. tab. 1. b. 1.

*Degeer Inf.* 1. tab. 17. fig. 22.

*Ammir. Inf.* tab. 21.

*Schaeff. Icon.* tab. 88. fig. 4. 5.

*Kleman. Inf.* 1. tab. 39. fig. 6.

Habitat in Betula, Salice, Rosa, polyphaga.

*Larua* obscura, tuberculis sparsis, capite fisco.

*Puppa* nuda brunnea.

*sacraria*. 57. *P. pectinicornis*, alis flavis, striga obliqua sanguinea.

*Syst. Ent.* 628. 42. \* *Linn. Syst. Nat.* 2. 863. 220.

Habitat in Africa.

*elingua*- 58. *P. pectinicornis*, alis griseis, anticis fascia lata obscuriore, medio puncto nigro. *Syst. Ent.* 628. 43. *Linn.*

*Syst. Nat.* 2. 862. 211. *Fn. Sv.* 1235.

*Roef. Inf.* 1. phal. 3. tab. 9.

Habitat in Europae Pyro.

*Larua* rugosa cinereo brunneopue variegata.

*Puppa* folliculata, antice fusca, postice brunnea.

*Variat* interdum antennis setaceis.

*plumbaria*. 59. *P. pectinicornis*, alis plumbeis, strigis tribus punctoque medio fuscis. *Syst. Ent.* 628. 44. \*

Habitat in Angliae nemoribus.

*purpura*- 60. *P. pectinicornis*, alis luteis, margine anticarumque fasciis duabus purpureis. *Syst. Ent.* 629. 45. *Linn.*

*Syst. Nat.* 2. 864. 221. *Fn. Sv.* 1254.

*Phalaena* pectinicornis, elinguis, alis deflexis luteo rubris, fascia duplici transuersa sanguinea. *Geoff. Inf.* 2. 126. 34.

*Clerk: phal.* tab. 9. fig. 11.

Goed. Inf. 2. tab. 12.

Habitat in Quercu, Pruno spinosa.

61. *P. pectinicornis*, alis atris albo subfasciatis, subtus griseis, fasciis fuscis. *Syst. Ent.* 629. 46. *fuliginaria*.

*Phalaena fuliginaria* pectinicornis, alis nigris albo punctato undulatis, subtus griseiscentibus. *Linn. Fn. Sv.* 1247.

Habitat in Sueciae ericetis.

62. *P. pectinicornis*, alis cinereis, strigis fasciaque postica fuscis. *minutaria*.

Habitat in Italia. *D. Allioni*.

*Minuta*. Corpus nigrum antennis pectinatis. Alae omnes cinereae strigis plurimis fuscis et versus marginem posticam fascia latiori eiusdem coloris.

### Antennis setaceis.

63. *P. feticornis*, alis angulato caudatis flavis, anticis fasciis duabus punctatis rufis. *politata*.

*Phalaena politia*. *Cram. Inf.* 12. tab. 139. fig. E.

Habitat Surinami.

Alae omnes flavissimae, anticis fasciis duabus e punctis numerosis rufis; altera in medio, altera marginali. Inter fascias ad marginem crassiorem lunula fusca punctis rufis.

64. *P. feticornis*, alis angulato caudatis albis fusco fasciatis punctisque tribus caudalibus atris. *caudata*.

*Phalaena fasciata*. *Cram. Inf.* 9. tab. 104. fig. D.

Habitat in Batavia.

Statura omnino *P. strigariae*, at antennae setaceae. Alae omnes albae fasciis numerosis pallide fuscis. Puncta tria atra in apice alarum posticarum.

65. *P. feticornis*, alis angulatis fuscis, fascia niuea. *Syst. fasciata*. *Ent.* 629. 47. \*

Habitat in India orientali.

66. *P. feticornis*, alis angulatis fusco albo nigro fulvoque variis, posticis subtus albis. *Cynata*. *Gen. Inf. Mans.* 236. \*

*Pha-*

254 CLASS. VI. GLOSSATA. Phalaena.

*Phalaena Cynara. Cram. Inf. 2. tab. 14. fig. C. D.*  
Habitat Surinami.

*regulata.* 67. *P. feticornis*, alis angulatis cinereis, anticis limbo fusca. *Syst. Ent.* 629. 48. \*  
Habitat in Brasilia. *Mus. Dom. Banks.*

*falcata.* 68. *P. feticornis*, alis falcatis fulvis, punctis duobus fuscis inter strigas luteas. *Syst. Ent.* 629. 49. \*  
Habitat in Anglia.

*sinuata.* 69. *P. feticornis*, alis emarginatis griseis, punctis duobus fuscis inter strigas albidas. *Gen. Inf. Mant.* 287. \*  
Habitat in Germaniae nemoribus.

*mantata.* 70. *P. feticornis*, alis angulatis fuscis, fascia obscuriorum anticarum puncto ocellari atro. *Gen. Inf. Mant.* 287. \*  
*Phalaena Manto. Cram. Inf. 1. tab. 10. fig. F.*  
Habitat Surinami.

*netrata.* 71. *P. feticornis*, alis angulato dentatis viridibus, punctis macula media alba.  
*Phalaena Netrix. Cram. Inf. 13. tab. 151. fig. E.*  
\* Habitat in America meridionali.

Thorax flavus. Abdomen rufum basi flavum. Alae omnes valde angulatae, virides atomis fuscis marginibus neque postico ferrugineo. Macula magna alba in centro alae posticae.

*phylirata.* 72. *P. feticornis*, alis angulatis viridibus, macula subocellari fulva.  
*Phalaena Phylira. Cram. Inf. 15. tab. 70. fig. D.*  
Habitat Surinami.

Alae omnes virides macula media fulva puncto postico virescente. Margo tenuissime fulvus.

*demandata.* 73. *P. feticornis*, alis subangulatis lutescentibus, strigibus fuscis punctoque medio atro. *Syst. Ent.* 629. 50. \*  
Habitat in Lipsiae hortis.

4. *P. feticornis*, alis dentatis ferrugineis, fascia fusca, *explanata*.  
puncto nigro. *Syst. Ent.* 630. 51.\*  
Habitat in China. *Mus. Dom. Monson.*
5. *P. feticornis*, alis subangulatis cinereis, strigis quatuor fuscis. *cinerata*.  
Habitat in Germaniae nemoribus. *Dom. de Hattorf.*  
Media. Corpus fuscum. Alae omnes obscure cinereae  
strigis quatuor fuscis parum vndatis, quarta obsoleta. Subtus cinereae strigis obsoletis albis.
6. *P. feticornis*, alis dentatis, antice flavis postice fuscis. *dimidiata*.  
*Syst. Ent.* 630. 52.\*  
Habitat in Anglia. *Mus. Dom. Monson.*
7. *P. feticornis*, alis dentatis flavis fusco irroratis, sub-*irrorata*.  
tus macula fasciaque postica fuscis. *Syst. Ent.* 630.  
53.\*  
Habitat in India orientali. *Mus. Tottianum.*
8. *P. feticornis*, alis dentatis flavo fuscoque variis, striga *Carinenta*.  
angulata atra.  
*Phalaena Carinenta. Cram. Inf.* 11. tab. 128. fig. F.  
Habitat in India orientali.  
Maculae duae artae in medio alae anticae.
9. *P. feticornis*, alis angulatis, omnibus viridibus, striga *viridata*.  
pallida. *Syst. Ent.* 630. 54. *Linn. Syst. Nat.* 2.  
865. 230. *Fn. Sv.* 1256.  
*Harris Aurel.* tab. 3. fig. c — i.  
Habitat in Quercu, *Oxyacantha*.
10. *P. feticornis*, alis dentatis griseis, strigis pallidioribus vndatis obsoletis. *obscurata*.  
*Gen. Inf. Mant.* 287.\*  
Habitat in Germania. *D. Schulz.*
11. *P. feticornis*, alis angulatis pallidis, strigis tribus sub-*notata*.  
fuscis, anticis punctis quatuor approximatis. *Syst.*  
*Ent.* 630. 55.  
12. *Phalaena notata feticornis*, alis angulatis pallidis, fasciis tribus subfuscis, anticarum postica nigricante commaculata. *Linn. Syst. Nat.* 2. 866. 232. *Fn. Sv.* 1258.

Clerk.

Clerk. phal. tab. 6. fig. 11.

Habitat in Europae Ericetis.

*nitata.* 82. *P. feticornis*, alis subdentatis obscure griseis, subtus pallidioribus, puncto fasciaque fuscis. *Gen. Inf. Mant.* 288. \*

Habitat in Germania. *D. Schulz.*

*areata.* 83. *P. feticornis*, alis angulatis niueis, anticis macula marginali, posticis strigis quatuor approximatis fuscis. *Phalaena Area. Cram. Inf.* 5. tab. 56. fig. D.

Habitat Surinami.

Statura *P. notatae*. Alae posticae angulatae versus marginem posticum strigis quatuor approximatis fuscis.

*decorata.* 84. *P. feticornis*, alis subangulatis cinereis, anticis fasciis tribus, posticis vnica flauis auro marginatis.

Habitat in Iamaica Insula. *Dom. Teats.*

Media. Alae anticae cinereae, holosericeae fasciis tribus flauis, quarum prima bascos parua, postice auro marginata, secunda approximata, antice auro marginata, tertia maior, communis antice auro marginata et ad marginem crassiolem puncto aureo notata. Margo alae terminatur striga fusco ferruginea. Posticae cinereae holosericeae subangulatae, pone medium fascia lata flaua vtrunque auro marginata et ad angulum ani striga abbreviata atra. Pone fasciam ala pallida macula sesqui altera fulua puncto aureo. Subtus omnes cinereae. Abdomen cinereum segmentorum marginibus tenuissime flauis.

*porata.* 85. *P. feticornis*, alis pallidis rufo puluerulentis, omnibus puncto ocellari albo. *Syst. Ent.* 631. 56. \*

*Phalaena porata* alis subangulatis pallefcentibus, posticis oculo minuto. *Linn. Syst. Nat.* 2. 866, 233.

Habitat in Europa, Anglia.

*emarginata.* 86. *P. feticornis*, alis subemarginatis pallidis, fasciis duabus griseis punctoque fusco. *Syst. Ent.* 631. 57.

*Phalaena emarginata feticornis*, alis omnibus pallidis emarginato angulatis, strigis duabus griseis. *Linn. Syst. Nat.* 2. 866, 236. *Fn. Sv.* 1257.

Clerk.

# **CLASS. VI. GLOSSATA. Phalaena. 257**

*Clerk. phal. tab. 4. fig. 12.*

Habitat in Europae nemoribus.

7. *P. feticornis*, alis coerulescentibus striatis, anticis apice atris, albo maculatis. *bicolora*-*ta*.

*Phalaena bicolora. Cram. Inf. 12. tab. 143. fig. A.*

*Phalaena bicolora. Sulz. Inf. tab. 22. fig. 6.*

Habitat Surinami.

**Magna.** Thorax fuscus albo punctatus. Abdomen fuscum. Alae anticae basi coerulescentes nigro striatae, apice atrae albo unaculatae, posticae coerulescentes albo striatae.

- *P. feticornis*, alis albidis, maculis rotundatis nigris, *grossulari*-*antici* strigis luteis. *Syst. Ent. 631. 58. Linn. Syst. Nat. 2. 867. 242. Fn. Sv. 1261.*

*Phalaena feticornis spirilinguis*, alis patentibus albis maculis inaequalibus nigris plurimis fasciaque transversa lutea. *Geoff. Inf. 2. 136. 57.*

*Phalaena media*, alis amplis albis, maculis crebris nigris et lineis transversis luteis variis. *Raj. Inf. 178. 14.*

*Goed. Inf. 1. tab. 31.*

*Blank. Inf. tab. 7. fig. E. H.*

*Merian. Europ. 1. tab. 29.*

*Albin. Inf. tab. 43. fig. 71.*

*Frisch. Inf. 3. tab. 2.*

*Rosf. Inf. 1. phal. 3. tab. 2.*

*Wlk. pap. 42. tab. 2. a. 1.*

*Edw. Av. tab. 202.*

*Ammiral. Inf. tab. 26.*

Habitat in Ribe, Grossularia frequens.

*Larva* geometra subpilosa, supra cinerea nigro punctata, subtus flava.

*Eruca* geometra, maiuscula, albo nigro et rubro coloribus varia. *Raj. Inf. 179.*

*Pappa* subfolliculata atra, fasciis flavis.

- *P. feticornis*, alis omnibus, strigis confertissimis v. *undulata*-*dulatis* fuscis. *Syst. Ent. 632. 59. Linn. Syst. Nat. 2. 867. 239. Fn. Sv. 1272.*

*Clerk. phal. tab. 6. fig. 2.*

*Vdm. Diff. 71.*

*Fabr. Sp. Inf. T. II.*

R

Habitat

258 CLASS. VI. GLOSSATA. Phalaena.

Habitat in Salice caprea Europae.

Conf. *Harris Inf. angl.* tab. 2. fig. 5. 6. Vix differre videtur, at nostra paullo minor.

*hespara-* 90. *P. feticornis*, alis rotundatis concoloribus coerulescentibus, fascia lata alba.

*Phalaena Hespertia. Cram. Inf.* 5. tab. 56. fig. 6.

Habitat in Indiis.

Alae omnes coeruleae, apice obscuriores, fascia medialba, immaculata.

*flaueola-* 91. *P. feticornis*, alis concoloribus nigris, disco communiflauo crenato. *Linn. Syst. Nat.* 2. 867. 240. *Muf.*

*Lud. Vlr.* 396.

*Phalaena flaueolata. Cram. Inf.* 8. tab. 88. fig. C.

Habitat in Indiis.

*populata.* 92. *P. feticornis*, alis flauo pallidis, anticis subfasciatis, apice subtus fusco contaminatis. *Syst. Ent.* 632. 60.

*Linn. Syst. Nat.* 2. 868. 244. *Fn. Sv.* 1266.

*Clerk. phal.* tab. 5. fig. 14. 15.

Habitat in Populo tremula Europae borealis.

*comitata.* 93. *P. feticornis*, alis anticis flauescentibus, fasciis tribus griseis, puncto apicisque lineola fuscis. *Syst. Ent.*

632. 61. *Linn. Syst. Nat.* 2. 868. 247. *Fn. Sv.* 1264.

Habitat in Europae nemoribus.

*ulmata.* 94. *P. feticornis*, alis albidis, fasciis duabus ferrugineofuscis, postica maculari. *Syst. Ent.* 632. 62. \*

*Sepp. Inf.* 6. 13. tab. 3.

Habitat in Angliae Vlmo, Platano.

*Larva* viridis, nigro lineata, capite caudaque nigra.

*Puppa* folliculata, coerulescens.

*dealbata.* 95. *P. feticornis*, alis albis, subtus nigro venosis. *Syst. Ent.* 632. 63. *Linn. Syst. Nat.* 2. 870. 256.

*Sulz. Hist. Inf.* tab. 23. fig. 3.

Habitat in Germaniae sylvis.

*aurata.* 96. *P. feticornis*, alis rotundatis fulvis, punctis strigisque postica aureis.

*Phalaena Aura. Cram. Inf. 5. tab. 53. fig. F.*

Habitat Surinami.

Striga flexuosa aurea ad marginem posticum. Antennae fuscae.

97. *P. feticornis*, alis griseis, anticis fasciis duabus pallidis repandis, postica semiterminali. *Syst. Ent. 692. prunata.*

64. *Linn. Syst. Nat. 2. 869. 250. Fn. Sv. 1267.*

*Clerk. phal. tab. 7. fig. 3.*

*Frisch. Inf. 5. tab. 14.*

*Ammiral. Inf. tab. 23. fig. 1 — 4.*

Habitat in Pruno, Ribe.

98. *P. feticornis*, alis flauissimis, anticis maculis costalibus *crataegatribus* ferrugineis, media subargentea. *Syst. Ent. 633. ta.*

65. *Linn. Syst. Nat. 2. 868. 243. Fn. Sv. 1283.*

*Phalaena feticornis spirilinguis*, alis patentibus luteis, duplici punctorum cinereorum ordine, superioribus maculis duabus et rachi croceo ferrugineis. *Geoff. Inf. 2. 139. 59.*

*Sepp. Inf. 6. 25. tab. 6.*

*Wilks pap. 39. tab. 1. b. 4.*

*Clerk. phal. tab. 5. fig. 15.*

Habitat in Crataego.

*Larva* grisea, tuberculo dorsali didymo.

*Puppa* fusca, folliculo cinereo ferrugineo maculato.

99. *P. feticornis*, alis omnibus nigris, fascia fulva. *Gen. equestrata. Inf. Mant. 288. \**

Habitat in Germania. *D. Schulz.*

100. *P. feticornis*, alis obscure coeruleis, striga punctorum aureorum. *lucidata.*

*Phalaena lucida. Cram. Inf. 11. tab. 128. fig. D.*

Habitat Surinami.

Alae omnes obscure coerulae strigis obscurioribus.

Versus marginem striga e punctis aureis.

101. *P. feticornis*, alis omnibus albis, margine exteriori *marginatolimbo* fusco interrupto. *Syst. Ent. 633. 66. Linn. ta.*

*Syst. Nat. 2. 870. 257. Fn. Sv. 1279.*

*Phalaena feticornis spirilinguis*, alis patentibus albis

R A

margi-



260 CLASS. VI. GLOSSATA. Phalaena.

marginē vndique interrupto fuscis. *Geoff. Inf. 2.*  
139. 60.

*Clerk. phal. tab. 2. fig. 5.*

*Sulz. Inf. tab. 16. fig. 96.*

Habitat in Europae Corylo.

*miata.* 102. *P. feticornis*, alis griseis, fasciis tribus viridibus, intermedia latiore fusco vndata. *Syst. Ent. 633. 87.*

*Phalaena miata feticornis*, alis virescenti vndulatis, fascia media lata triplici onotata. *Linn. Syst. Nat. 1.*  
869. 249. *Fn. Sv. 1268.*

*Clerk. phal. tab. 8. fig. 2.*

*Reaum. Inf. 2. tab. 29. fig. 11 — 13.*

*Amnstrl. Inf. tab. 32. fig. 2.*

Habitat in Sueciae Pruno spinosa, Fraxino.

*coerulata.* 103. *P. feticornis*, alis obscuris viridibus fusco fasciatis fascisque duabus coeruleis. *Gen. Inf. Mant. 288.\**

Habitat Hamburgi. *D. Schulz.*

*rufata.* 104. *P. feticornis*, alis cinereis, postice ferrugineis, stigma alba punctisque marginalibus nigris. *Syst. Ent. 633.*  
68.\*

Habitat in Angliae nemoribus.

*albicillata.* 105. *P. feticornis*, alis omnibus nigricantibus, fascia lata alba immaculata. *Syst. Ent. 634. 69. Linn. Syst. Nat. 2.*  
870. 255. *Fn. Sv. 1278.*

*Clerk. phal. tab. 1. fig. 12.*

Habitat in Europa boreali.

*hastata.* 106. *P. feticornis*, alis omnibus nigris albo maculatis, fasciis duabus albis nigro punctatis hastato dentatis. *Syst. Ent. 634. 70. Linn. Syst. Nat. 2. 870. 254. Fn.*  
*Sv. 1276.*

*Phalaena* antennis filiformibus, alis latis albis fasciis vndatis maculisque hastatis nigris. *Degeer Inf. Vef.*  
*Germ. 2. 1. 334. 7. tab. 8. fig. 20.*

*Clerk. phal. tab. 1. fig. 9.*

*Kleman. Inf. 1. tab. 44.*

Habitat in Europae Betula.

*Larva* fusco brunnea, maculis lateralibus flexuosis  
*Puppa* folliculata, brunnea.

107. *P. feticornis*, alis omnibus nigris albo subundatis, *tristata*. fasciis duabus albis immaculatis. *Syst. Ent.* 634. 71. *Linn. Syst. Nat.* 2. 869. 252. *Fn. Sv.* 1277.

*Clerk. phal.* tab. 1. fig. 13.

Habitat in Betula Europae borealis.

108. *P. feticornis*, alis omnibus flavescentibus, lineis ni- *clathrata*. gris decussatis. *Syst. Ent.* 634. 72. *Linn. Syst. Nat.* 2. 867. 238. *Fn. Sv.* 1275.

*Phalaena feticornis spirilinguis*, alis patentibus fuscis, utrinque maculis albis quadrangulis tessulatis, *Geoff.* *Inf.* 2. 135. 53.

*Clerk. phal.* tab. 2. fig. 11.

*Phalaena retialis. Scop. carn.* 536.

Habitat in Europae Ericetis.

109. *P. feticornis*, alis griseis, anticis striga, fascia puncto- *gristata*. que fuscis. *Gen. Inf. Mant.* 288. \*

Habitat Hamburgi. *D. Schulz.*

110. *P. feticornis*, alis rotundatis nigris, striga ferruginea, *Syrata*. anticis margine crassiori cinereo.

*Phalaena Syero. Cram. Inf.* 2. tab. 97. fig. F.

Habitat Surinami.

Parua. Corpus cinereum abdomine nigro. Alae concolores nigrae striga postica ferruginea. Margo crassior alarum anticarum late cinereus.

111. *P. feticornis atra*, alis erectis, anticis apice albis. *chaero-* *Syst. Ent.* 634. 73. *Linn. Syst. Nat.* 2. 866. 237. *Fn. phyllata*. *Sv.* 1274.

*Alt. Soc. Berol. nat.* 3. tab. 1. fig. 3.

*Petiv. Gazoph.* tab. 69. fig. 8.

Habitat in Chaerophyllo sylvestri.

112. *P. feticornis*, alis cinerascens, anticis fasciis tri- *flavata*. bus abbreviatis fuscis. *Syst. Ent.* 634. 74. *Linn. Syst.*

*Nat.* 2. 871. 260. *Fn. Sv.* 859.

*Goed. Inf.* 3. tab. k.

*Merian. Europ.* tab. 120.

*Frisch. Inf.* 7. tab. 19.

Habitat in Pomonae arboribus.

262 CLASS. VI. GLOSSATA. Phalaena.

*bilineata*. 113. *P. feticornis*, alis luteis testaceo vndatis, fascia repanda, margine fusco alboque. *Syst. Ent.* 635. 75. *Lin.*  
*Syst. Nat.* 2. 868. 245. *Fn. Sv.* 1284.

*Clerk. phal.* tab. 6. fig. 13.

Habitat in Europae nemoribus frequens.

*bimacula*. 114. *P. feticornis*, alis albis, anticis maculis duabus costalibus fuscis. *Syst. Ent.* 635. 76. \*

Habitat in Lipsiae hortis.

*volutata*. 115. *P. feticornis*, alis omnibus viridibus, strigis duabus albis. *Syst. Ent.* 635. 77. \*

*Harris Inf. angl.* tab. 8. fig. 3.?

Habitat in Germania.

*fimbriata*. 116. *P. feticornis*, alis plumbeis, fascia lata fusca, flauo marginata. *Gen. Inf. Mant.* 289. \*

*Phalaena XIII. Schaeff. Icon.* tab. 80. fig. 6: 7.

Habitat in Germania. *D. Schulz.*

*quadrisfciata*. 117. *P. feticornis*, alis griseo fuscis, fascia lata nigra, puncto subocellari atro. *Gen. Inf. Mant.* 289. \* *Lin.*  
*Fn. Sv.* 1253.

Habitat in Germania. *D. Schulz.*

*ocellata*. 118. *P. feticornis*, alis albidis, fascia lata abbreviata fusca, puncto subocellari atro. *Syst. Ent.* 635. 78. \*

*Phalaena ocellata feticornis*, alis albidis, anticis fasciis duabus fuscescentibus maculaque apicis ocellari didyma. *Lin. Syst. Nat.* 2. 870. 258. *Fn. Sv.* 1280.

*Phalaena Alstroemeriana. Sutz. Hist. Inf.* tab. 23. fig. 5.

Habitat in Europa boreali.]

Inuenitur et *pectinicornis*.

*lyneata*. 119. *P. feticornis*, alis rotundatis albis, fasciis duabus punctoque apicis fuscis. *Syst. Ent.* 635. 79. \*

Habitat in Angliae nemoribus.

*vinulata*. 120. *P. feticornis*, alis albidis, fasciis duabus nigris, posteriori terminali. *Gen. Inf. Mant.* 289. \*

Habitat in Germania. *Dom. de Hattorf.*

121. *P. feticornis*, alis cinereis, posticis supra fuscis, fascia *Ianata*. punctisque tribus albis. *Syst. Ent.* 635. 80.

*Phalaena Ianata* feticornis, alis anterioribus subtus cinereis, posticis subtus fascia, punctisque tribus albis. *Linn. Syst. Nat.* 2. 870. 259. *Mus. Lud. Vlr.* 398.

Habitat in Indiis.

122. *P. feticornis*, alis flavescentibus, striga nigra postice pallidioribus, femina aptera nigro punctata. *Syst. Ent.* 636. 81. *Linn. Syst. Nat.* 2. 874. 281. *Fn. Sv.* 1293.

*Phalaena* alis latis exterioribus albicantibus cum tinctura viridi aut coerulea. *Raj. Inf.* 181. 4.

*Phalaena* antennis filiformibus, alis griseis, strigis obscurioribus, femina aptera. *Degeer Inf. Verf. Germ.* 2. 1. 331. 6.

*Reaum. Inf.* 2. tab. 27. fig. 8. 9.

*Degeer Inf.* 1. tab. 24. fig. 11. 19.

*Frisch. Inf.* 3. 8. tab. 2.

*Roef. Inf.* 3. tab. 14. ♀

*Roef. Inf.* 3. tab. 40. fig. 6. ♂

Habitat in Pomonae arboribus.

*Larva* ferruginea, linea laterali flava.

123. *P. feticornis*, alis rotundatis viridibus, margine ferrugineo. *Syst. Ent.* 636. 82.\* *expectata.*

Habitat in Asia. *Dom. Drury.*

124. *P. feticornis*, alis anticis testaceis, fasciis tribus griseis, puncto prominulo apicisque lineola supra fusca. *Syst. Ent.* 636. 83. *Linn. Syst. Nat.* 2. 868. 246. *chenopodiata.*

*Fn. Sv.* 1263.

*Alb. Inf.* tab. 47. fig. 79.

*Kleman. Inf.* tab. 37.

Habitat in Europae Chenopodio.

*Larva* nuda, virescens aut fusca, lineata, segmentis angulatis.

*Puppa* folliculata, flavescens.

*Imago* mare subpectinicorni.

- Centum-notata.* 125. *P. feticornis*, alis ferrugineo fuscoque variis, fascia media grisea C atro notata. *Gen. Inf. Mant.* 289.\*  
*Phalaena centumnotata.* *Naturf.* 6. tab. 4. fig. 4.  
 Habitat in Germania. *D. Schulz.*
- duplicata.* 126. *P. feticornis*, alis griseis, strigis duabus fuscis duplicatis. *Syst. Ent.* 636. 84.\*  
 Habitat in Angliae nemoribus.
- lunulata.* 127. *P. feticornis*, alis fuscis postice strigis vndatis flauis. *Gen. Inf. Mant.* 289.\*  
 Habitat in Germaniae nemoribus.
- Alchemillata.* 128. *P. feticornis*, alis fuscescentibus, anticis vndatis, fascia niuea cinereo vndata lineolaque intra apicem. *Linn. Syst. Nat.* 2. 869. 253. *Fn. Sv.* 1282.  
*Degeer Inf.* 1. tab. 22. fig. 16.  
 — — *Versf. Germ.* 2. 1. 259. 10.  
 Habitat in Europae borealis Alchemilla.  
 Duplo minor *P. rectangulata.*  
*Larua* glabra viridis, flauo fasciata lineaque laterali alba.
- rectangulata.* 129. *P. feticornis*, alis viridibus fusco subfasciatis. *Syst. Ent.* 636. 85. *Linn. Syst. Nat.* 2. 872. 270. *Fn. Sv.* 1295.  
*Clerk. phal.* tab. 8. fig. 6.  
 Habitat in Europae borealis sylvis.
- succenturiata.* 130. *P. feticornis*, alis albidis, puncto nigro limboque obscuriore. *Syst. Ent.* 637. 86. *Linn. Syst. Nat.* 2. 872. 267. *Fn. Sv.* 1294.  
*Clerk. phal.* tab. 5. fig. 2.  
 Habitat in Europae borealis Ericetis.
- ferrugata.* 131. *P. feticornis*, alis purpurascensibus cinereo vndulatis, postice macula didyma fusca. *Syst. Ent.* 637. 87. *Linn. Fn. Sv.* 1292.  
 Habitat in Europa boreali.
- maculata.* 132. *P. feticornis*, alis flauis nigro maculatis. *Syst. Ent.* 637. 88.  
*Phalaena macularia.* *Linn. Syst. Nat.* 2. 862. 213.  
*Scop. carn.* 557.  
 Habitat in Angliae Ericetis frequens.

133. *P. feticornis*, alis flavis nigro punctatis, punctis *cribrata*.  
baseos suboccellatis.

*Phalaena cribraria*. *Cram. Inf.* 18. tab. 208. fig. C. G.

*Phalaena cribraria*. *Clerk. Icon.* tab. 54. fig. 7. 8.

*Drury Inf.* 2. tab. 6. fig. 3.

Habitat in India orientali.

Affinis videtur praecedenti, at puncta nigra baseos an-  
nulo albo cincta.

134. *P. feticornis*, alis rotundatis niuceis, anticis pun- *punctata*.  
cto medio fusco margineque nigro punctato. *Syst.*

*Ent.* 637. 89.\*

Habitat in Angliae nemoribus.

135. *P. feticornis*, alis omnibus griseo vndatis, puncto *mediata*.  
centrali fusco margineque nigro punctato. *Gen. Inf.*

*Mant.* 290.\*

Habitat in Germania. *Dom. de Hattorf.*

136. *P. feticornis*, alis flauissimis, fascia lata fusca retror- *sociata*.  
sum dentem exferente. *Syst. Ent.* 637. 90.\*

*Harris Aurel.* tab. 35. fig. 1.

Habitat in Angliae nemoribus.

137. *P. feticornis*, alis flavescentibus fusco vndatis, anti- *contrata*.  
cis puncto centrali nigro. *Gen. Inf. Mant.* 290.\*

Habitat Hamburgi. *D. Schulz.*

138. *P. feticornis*, alis cinereis, puncto centrali margi- *cinerata*.  
neque postico vndato fuscis.

Habitat in Germania. *Dom. de Hattorf.*

Alae omnes cinereae vel pallidae, basi immaculae, pun-  
cto centrali fusco: margo posticus late fuscus cine-  
reo vndatus: subtus concolores.

139. *P. feticornis*, alis virescentibus, anticis fasciis dua- *purpura*-  
bus purpureis. *Syst. Ent.* 637. 91. *Lin. Fn. Sv.* *ta.*

1302.

Habitat in Europae borealis ericetis.

140. *P. feticornis*, alis fulvis, strigis vndatis fusco ferru- *corruga*-  
gineis. *ta.*

Habitat in Italia. *Dr. Allioni.*

266 CLASS. VI. GLOSSATA. *Phalaena*.

Parua. Alae omnes dilute fulvae, strigis plurimis vndatis obscurioribus. Subtus concolores puncto centrali nigro.

*immuta*-141. *P. feticornis*, alis niueis, strigis vndatis obscurioribus, margine postico nigro punctato. *Syst. Ent.* 637. 92.\*

*Phalaena immutata feticornis*, alis albis concoloribus, strigis pallidis, puncto margineque postico nigro punctato. *Linn. Syst. Nat.* 2. 871. 264. *Fn. Su.* 128. *Clerk. phal.* tab 6. fig. 12.

*Phalaena feticornis spirilinguis*, alis patentibus albidis atomis cinerascens et fascia vndata ferruginea. *Geoff. Inf.* 2. 140. 63.

Habitat in Angliae nemoribus frequens.

*variegata*-142. *P. feticornis*, alis omnibus purpureis, maculis margineque flauis. *Gen. Inf. Mant.* 291.\*

Habitat Hamburgi. *D. Schulz.*

*diaphana*-143. *P. feticornis*, alis hyalino albis nigro punctatis, apice strigis fulvis abbreviatis.

*Phalaena diaphana*. *Cram. Inf.* 10. tab. 113. fig. G. Habitat in Sierra Leon Africae.

Parua. Alae omnes albae nigro punctatae et versus apicem strigae variae, abbreviatae fulvae.

*cingulata*-144. *P. feticornis*, alis omnibus fuscis, striga niuea. *Syst. Ent.* 638. 93. *Linn. Syst. Nat.* 2. 874. 280. *Fn. Su.* 1303.

*Clerk. phal.* tab. 2. fig. 10.

Habitat in Europae borealis cricetis.

*lineata*-145. *P. feticornis*, alis striatis niueis immaculatis. *Syst. Ent.* 638. 94.\*

Habitat in Americae meridionalis insulis.

*costata*-145. *P. feticornis*, alis niueis, anticis costa ferruginea. *Syst. Ent.* 638. 95.\*

Habitat in Americae meridionalis insulis.

CLASS. VI. GLOSSATA. Phalaena. 267

147. *P. feticornis*, alis concoloribus rufis, margine nifiſſicata.  
gro albo punctato. *Syſt. Ent.* 638. 96.\*  
Habitat in Africa aequinoctiali. *Dom. Drury.*
148. *P. feticornis*, cinerea, capite thoracis alarumque capitata.  
margine baſi ferrugineis. *Syſt. Ent.* 638. 97.\*  
Habitat in India. *D. Koenig.*  
Statura *P. Vorticatae*.
149. *P. feticornis*, cyaneo nigra albo maculata. *Syſt. Ent. aeſtuata.*  
638. 98.  
*Phalaena aeſtuata feticornis*, alis fulcis concoloribus,  
maculis albis oblongis, coadunatis. *Linn. Syſt. Nat.*  
2. 867. 241. *Muf. Lud. Vlr.* 397.  
Habitat in Indiis.
150. *P. feticornis*, alis apice flauis, anticis ſtrigis duabus *argenta-*  
*argenteis*, poſticis macula quadruplici nigra. *Syſt. ta.*  
*Ent.* 618. 99.\*  
Habitat in China. *Muf. Dom. Monſon.*
151. *P. feticornis*, alis atris albo maculatis, ano rufo. *melata.*  
*Gen. Inf. Mant.* 291.\*  
*Phalaena Melite. Cram. Inf.* 2. tab. 18. fig. D.  
Habitat in America boreali.
152. *P. feticornis*, alis flauis, maculis margineque poſtico *ſulphura-*  
fulco ferrugineis. *Syſt. Ent.* 639. 100. *ta.*  
Habitat in Indiis. *Muf. Dom. Monſon.*
153. *P. feticornis*, alis flauſcentibus, margine exteriori *ſinuata.*  
poſticoque purpureis ſinuatis.  
Habitat in Africa aequinoctiali. *Muf. Dom. Banks.*  
Media. Caput et thorax purpurea antennis ferrugineis.  
Abdomen album ano purpureo. Corpus ſubtus ni-  
ueum. Alae anticae flauſcentes margine exteriori et  
poſtico late purpureis varie ſinuatis, poſticae flau-  
ſcentes margine poſtico maculis duabus purpureis.  
Subtus omnes concolores, at margo niger.
154. *P. feticornis*, alis albis fulco faſciato maculatis tho- *vorticata.*  
race anoque flauis. *Syſt. Ent.* 639. 101. *Linn. Syſt.*  
*Nat.* 2. 873. 272. *Fv. Sv.* 1297.

*Phalae-*



*Phalaena feticornis spirilinguis*, alis patentibus albo fuscoque nebulosis, ano flavo. *Geoff. Inf.* 2. 135.

*Phalaena* antennis filiformibus, alis horizontalibus albis, maculis ovalibus nigris, thorace caudaque nigris. *Degeer Inf. Vers. Germ.* 2. 1. 326.

*Phalaena* minor, alis oblongis ex albo et coeruleo nigricante variis ad exortum flavis. *Raf. Inf.* 222. 73. *Goed. Inf.* 2. tab. 13.

*Petiv. Gazoph.* tab. 32. fig. 8.

*Alb. Inf.* tab. 37. fig. 60.

*Reaum. Inf.* 1. tab. 49. fig. 17. 18.

*Geoff. Inf.* 1. tab. 28. fig. 18. 19.

*Roef. Inf.* 1. phal. 4. tab. 14.

*Harris Aurel.* 13. tab. 6. fig. a — p.

*Schaeff. Elem.* tab. 98. fig. 4.

— — *Icon.* tab. 119. fig. 1. 2.

Habitat in Vrticæ, Pomona.

*Larva* 16poda, nuda, albida, dorso fusco, collari bipunctato.

Latet intra folium Vrticæ, quod contorquet filisque connectit. Coniungit Geometras Linnaei cum Pyralidibus.

*Puppa* folliculata, brunnea.

*funerata.* 155. *P. feticornis*, alis rotundatis atris, anticis maculis duabus, posticis vnica hyalinis.

*Phalaena Tages. Cram. Inf.* 9. tab. 97. fig. D.

*Phalaena Vfeus. Cram. Inf.* 9. tab. 97. fig. E.

Habitat Surinami.

Affinis videtur *P. atrali*. Abdomen fuscum.

*limbata.* 156. *P. feticornis*, alis rotundatis luteis, puncto ocellari margineque postico fuscis. *Syst. Ent.* 639. 102. *Lin. Syst. Nat.* 2. 873. 273.

Habitat in Europæ borealis nemoribus.

*metallata.* 157. *P. feticornis*, alis niveis, costa maculaque media aureis.

Habitat in Africa æquinoctiali. *Mus. Dom. Banks.*

Media. Corpus album ano aureo. Alae omnes niveae purpureo aliquot micantes. Anticæ costa late maculaque media magna aureis. Posticæ macula media

**CLASS. VI. GLOSSATA. Phalaena. 259**

dia strigaeque postica fere obsoleta aurea. Subtus colorata.

158. *P. feticornis*, alis hyalino albis, margine nigro cin- *hyalina-*  
 ctis, ano lutescente. *Syst. Ent.* 640. 103. *Linn. Syst.* *td.*  
*Nat.* 2. 874. 279.

Habitat in America, in Asia. *D. Koenig.*

159. *P. feticornis*, alis cinereis albo maculatis, anticis ob- *potamo-*  
 soletis reticulatis. *Syst. Ent.* 640. 104. *Linn. Syst.* *gata.*  
*Nat.* 2. 873. 275. *Fn. Sv.* 1299.

*Reaum. Inf.* 2. tab. 32. fig. 11.

Habitat in Potamogetonte natante.

160. *P. feticornis*, alis pallide subfasciatis, anticis punctis *stratiota-*  
 tribus nigris. *Linn. Syst. Nat.* 2. 873. 276. *Fn.* *ta.*  
*Sv.* 1200.

*Geoff. Inf.* 2. 142. 66.

*Degeer Inf.* 1. tab. 37. fig. 16. 18.

Habitat in Stratiote Aloide.

*Larva* aquatica, hexapoda, viridis, spiraculis latera-  
 libus fasciculatis.

*Puppa* folliculata, vtrisque attenuata ferruginea sub-  
 aquatica.

161. *P. feticornis*, alis cinereis, omnibus conformiter al- *nympha-*  
 bo maculato-reticulatis. *Syst. Ent.* 640. 105. *Linn.* *ta.*  
*Syst. Nat.* 2. 873. 274. *Fn. Sv.* 1298.

*Phalaena minor*, alis vtrisque albo et fusco colore va-  
 riis. *Raj. Inf.* 227. 87.

Habitat in Nymphaea, Lemna, frequens.

162. *P. feticornis*, alis niueis, posticis fascia terminali ni- *lemnata.*  
 gra, punctis quatuor albis. *Syst. Ent.* 640. 106.  
*Linn. Syst. Nat.* 2. 874. 278. *Fn. Sv.* 1301.

*Phalaena aquatica minima alba sive argentea. Raj. Inf.*  
 205. 102.

*Reaum. Inf.* 2. tab. 12. fig. 14. 15.

*Harris Auct.* 15. tab. 7. fig. 1—p.

Habitat in Lemna aliisque plantis aquaticis ad paludes  
 frequentissima.

270 CLASS. VI. GLOSSATA. Phalaena.

- perspecta*. 163. *P. feticornis*, alis anticis griseis hyalino maculatis, posticis hyalinis, margine flauo hyalino maculato. *Syst. Ent.* 640. 107.\*  
Habitat in India orientali. *D. Koenig.*
- elevata*. 164. *P. feticornis*, alis flauissimis fusco punctatis, subtus immaculatis. *Gen. Inf. Mant.* 291.\*  
Habitat in Americae meridionalis insulis.
- indicata*. 165. *P. feticornis*, alis flauis, anticis strigis tribus, posticis duabus vndatis fuscis. *Syst. Ent.* 640. 108.\*  
Habitat in Americae meridionalis insulis. *Mus. Tottenhamum.*
- derogata*. 166. *P. feticornis*, alis albis, strigis plurimis fuscis. *Syst. Ent.* 641. 109.\*  
Habitat in India orientali. *D. Koenig.*
- cieclata*. 167. *P. feticornis*, alis fusco flauoque variis, maculis hyalinis sparsis. *Syst. Ent.* 641. 110.  
Habitat in India orientali. *D. Koenig.*
- fenestrata*. 168. *P. feticornis*, alis albohyalinis, strigis flauis numerosis flauescentibus. *Syst. Ent.* 641. 111.\*  
Habitat in Americae meridionalis insulis.

\* \* \* Alis forficatis.

- farinalis*. 169. *P. palpis recuruatis*, alis politis lutescentibus, strigis repandis albis, basi apiceque glaucis. *Syst. Ent.* 641. 112.

*Phalaena farinalis* palpis recuruatis, alis politis fuscescentibus, strigis repandis albidis, arca interiecta glauca. *Linn. Syst. Nat.* 2. 881. 327. *Fn. Sv.* 1348.

*Clerk. Icon.* tab. 2. fig. 14.

Habitat in Europae farina.

Sedens caudam erigit.

- glaucinalis*. 170. *P. palpis recuruatis*, alis glaucis, strigis duabus fuscis remotis. *Syst. Ent.* 641. 113. *Linn. Syst. Nat.* 2. 881. 328. *Fn. Sv.* 1349.

*Clerk.*

*Clerk. phal. tab. 3. fig. 4.*

Habitat in Europa boreali.

171. *P. palpis* antennis pectinatis brevioribus, femoribus *barbalis*.  
anticis barba porrecta. *Syst. Ent.* 641. 114. *Linn.*  
*Syst. Nat.* 2. 881. 329. *Fn. Sv.* 1345.

*Clerk. phal. tab. 5. fig. 3.*

Habitat in Europa boreali.

172. *P. palpis* porrectis longitudine antennarum pectinata- *tentac-*  
tarum, alis cinerascens, strigis tribus fuscis. *Syst. laris.*  
*Ent.* 641. 115. *Linn. Syst. Nat.* 2. 381. 330. *Fn. Sv.* 1344.

*Phalaena* antennis pectinatis, alis horizontalibus griseis, strigis duabus fuscis, palpis longitudine antennarum. *Degeer Inf. Vers. Germ.* 2. 1. 260. 12.

*Degeer Inf.* 1. tab. 5. fig. 1.

Habitat in Europa boreali.

173. *P. palpis* porrectis approximatis thorace longioribus, *proboscis-*  
alis griseis, strigis ferrugineis. *Syst. Ent.* 642. 116. *alis.*  
*Linn. Syst. Nat.* 2. 881. 331. *Fn. Sv.* 1346.

*Clerk. phal. tab. 5. fig. 4.*

*Sulz. Hist. Inf.* tab. 23. fig. 6.

Habitat in Europa.

174. *P. palpis* porrectis thorace longioribus, alis subgriseis, punctis duobus muricatis lineaque apicis nigris. *rostralis.*  
*Syst. Ent.* 642. 117. *Linn. Syst. Nat.* 2. 881. 332.  
*Fn. Sv.* 1347.

*Roes. Inf.* 1. phal. 4. tab. 6.

Habitat in Europae Carpino, Humulo.

*Larva* nuda, 14poda, viridis, strigis obliquis lineisque lateralibus albis.

*Puppa* folliculata, brunnea, obtusa, antice carinata.

175. *P. palpis* recurvatis, alis griseis, strigis tribus fuscis, *nemor-*  
intermedia flexuosa. *Syst. Ent.* 642. 118. *alis.*

*Harris Inf. angl.* tab. 6. fig. 2.?

Habitat in Angliae nemoribus.

176. *P. palpis* porrectis thorace longioribus, alis griseis, *palpalis.*  
posticis margine crassiori albo. *Syst. Ent.* 642. 119.\*

Habitat in Angliae nemoribus.

177. *P.*

372 CLASS. VI. GLOSSATA. Phalaena.

*forficatis.* 177. P. alis glabris pallidis, strigis obliquis ferrugineis.  
*Syst. Ent.* 642. 120. *Linn. Syst. Nat.* 2. 882. 334.  
*Fn. Sv.* 1352.

*Phalaena feticornis spirilinguis, triangularis, alis deflexis flavis, lineis duabus obliquis ferrugineis. Geoff. Inf.* 2. 166. 111.

*Reaum. Inf.* 1. tab. 16. fig. 13. 14.

Habitat in Europae Brassica.

*vndatis.* 178. P. alis obscure cinereis, strigis tribus obliquis vndatis niueis.

Habitat in Italia. *D. Allioni.*

Magnitudo praecedentis. Corpus fuscum. Alae obscure cinereae strigis tribus obliquis vndatis niueis. Margo posticus itidem niueus sex parium punctorum fuscorum.

*sagittalis.* 179. P. palpis recurvis, alis cinereis, macula costali atra posticis flavis, margine fusco.

*Noctua Sagitta. Syst. Ent.* 597. 90.

Habitat in India orientali.

Caput acuminatum.

*verticalis.* 180. P. alis glabris pallidis subfasciatis, subtus fusco vndatis. *Syst. Ent.* 643. 121. *Linn. Syst. Nat.* 2. 882. 335.  
*Fn. Sv.* 1353.

*Phalaena feticornis spirilinguis triangularis, alis deflexis albo virescentibus, subtus lineis duabus transversis nigris. Geoff. Inf.* 2. 166. 112.

*Phalaena parva ex albido sublutea, alis exterioribus duplici linea transversa nigricante oblique ducta notatis. Raj. Inf.* 257. 48.

*Reaum. Inf.* 2. tab. 19. fig. 1 — 5.

*Roef. Inf.* 1. phal. 4. tab. 4.

*Wilks pap.* 24. tab. 3. a. 16.

Habitat in Europae Vrtica.

*Larva* 16poda, pilosa, capite pedibusque flavis.

*Puppa* brunnea, segmentis tribus ultimis denticulo notatis.

*salicis.* 181. P. alis cinereis, anticis strigis tribus obliquis fulvis, antennis pubescentibus.

Pha

Fm

CLASS VI. GLOSSATA. Phalaena. 273

*Phalaena faticalis.* *Wien. Verz.* 285. tab. 1. a. fig. 5.

— — — — — tab. 1. b. fig. 5.

Habitat in Salice.

*Larua* 14poda, nuda, viridis, linea dorsali saturatiore.

*Puppa* nigra, nitida.

*Imago* statura obtusa fere Pyralidis, Strigae alarum e flavo fulvoque compositae.

182. *P. palpis recurvatis, alis fuscis nitidis, anticis puncto fuscalis.*  
subocellari niueo. *Syst. Ent.* 643. 122. \*

Habitat in India orientali. *Mus. Tottianum.*

183. *P. palpis recurvatis, alis cinereis, linea dorsali fasciata dorsalis.*  
que postica fuscis. *Syst. Ent.* 643. 123. \*

Habitat in Brasilia. *Mus. Dom. Banks.*

184. *P. palpis recurvatis, alis cinereis, margine crassiori pinguina-*  
nigro subfasciato. *Syst. Ent.* 643. 124. *Linn. Syst. his.*

*Nat.* 2. 882. 336. *Fn. Sv.* 1350.

*Phalaena* antennis filiformibus elinguis, palpis porrectis, alis horizontalibus achatinis, maculis strigisque nigris et ferrugineis. *Degeer Inf. Vers. Germ.* 2. 1.

271. 2. tab. 6. fig. 11.

*Reaum. Inf.* 3. tab. 20. fig. 5. 6. 7. 9. 10. 11.

*Roland. At. Holm.* 1755. 51. tab. 2.

*Schaeff. Icon.* tab. 6. fig. 8. 9.

Habitat in pinguibus, butyro, lardo aliisque. Frequens intra domos et culinas, rarius in ventriculo humano, inter vermes pessima.

*Larua* 16poda, laevis, fusca, nitida.

*Puppa* nuda, brunnea, obscura.

185. *P. alis flavis, strigis tribus vndatis ferrugineis.* *ferrugalis.*  
Habitat in Africa aequinoctiali. *Dom. Banks.*

Statura omnino praecedentium. Os album palpis apice ferrugineis. Alae omnes flavescentes anticis strigis tribus, posticis duabus ferrugineis. Subtus omnes palidae.

186. *P. alis griseo fuscis striatis, macula reniformi A latino inscripta.* *Syst. Ent.* 644. *Linn. Syst. Nat.* 2. 882. 338. *fecalis.*

*Fabr. Sp. Inf. T. II.*

S

Ro-

*Rotund. Alt. Holm. 1752. 62.*

Habitat intra culmos fecalinos, eos intra vaginam erigens, ex vno in alterum migrans; vnde spicæ albae cassæ. *Roland.*

*vinetis. 187. P. alis griseo fuscis, maculis duabus costalibus albis, corpore niueo.*

Habitat in Germania. *Dom. de Hattorf.*

Magnitudo præcedentis. Lingua supra alba. Alae antice planæ, griseo fuscae postice striga vndata punctisque marginalibus nigris. Maculae duae albidæ ad costam. Posticæ supra fuscae, subtus pallidæ punctis duobus, striga punctisque marginalibus nigris. Corpus subtus cum pedibus niueum.

*sanguinea-188. P. alis flavis, fasciis duabus sanguineis, priore exte-*  
*lis. rius dilatata, perforata. Syst. Ent. 644. 126. Linn.*  
*Syst. Nat. 2. 882. 339.*

Habitat in Europa boreali.

*sticticalis. 189. P. alis griseis, striga marginali flava margineque ni-*  
*gro ciliari. Syst. Ent. 644. 127. Linn. Syst. Nat. 2.*  
*883. 340. Fn. Sv. 1354.*

Habitat in Europa boreali.

*fascialis. 190. P. palpis recurvatis, alis fuscis, fasciis duabus flavis.*  
*Syst. Ent. 644. 128. \**

Habitat in nova Hollandia. *Mus. Dom. Banks.*

*strigalis. 191. P. alis griseis, strigis duabus flavis costaque purpu-*  
*rascente.*

Habitat in Germaniæ nemoribus. *Dom. de Hattorf.*

Magnitudo *P. forficalis*. Caput et thorax subferruginea. Alae antice obscure griseae strigis duabus distinctis flavis costaque purpurascente. Posticæ griseae strigis obsoletioribus.

*recurva-192. P. palpis recurvatis, alis fuscis, anticis fasciis duabus*  
*lis. fuscis, posteriore abbreviata. Syst. Ent. 644. 129. \**

Habitat in India orientali. *Mus. Tottianum.*

CLASS. VI. GLOSSATA. Phalaena. 275

1293. P. alis subferrugineis, striga flava, margine crassiori *punctalis*. albo punctato. *Syst. Ent.* 644. 130. \*

Habitat in Europae borealis nemoribus.

1294. P. alis purpurascens, omnibus luteo maculatis. *purpuralis*. *Syst. Ent.* 644. 131. *Linn. Syst. Nat.* 2. 883. 342. *Fn.* *lis.* *Sv.* 1356.

Habitat in Europa frequens.

*Variat rarius alis fuscis flavo maculatis.*

1295. P. alis cinereo fuscis, strigis anticarum tribus, posticarum duabus vndatis argenteis. *argentalis*.

Habitat in agro Pedemontano. *D. Allioni.*

Magnitudo P. purpuralis. Alae omnes cinereo fuscoque variae. Anticae striga bascos parva, abbreviata, puncto parvo nigro, striga altera ante medium angulata, tertia pone medium vndata. Posticae strigis duabus, posteriore vndata. Margo posticus omnium ciliaris argenteo fuscoque varius.

1296. P. alis purpurascens, maculis duabus costalibus *costalis*. margineque postico flavis. *Syst. Ent.* 644. 132. \*

Habitat in Angliae nemoribus.

1297. P. alis atris, omnibus maculis duabus niueis. *Syst. atralis*. *Ent.* 645. 133. \*

*Phalaena gmaculata feticornis*, alis nigris, omnibus maculis duabus albis rotundatis. *Linn. Mant.* 540.

*Phalaena Funeraria*. *Mull. Zool. Dan.* 132. 1524.

*Phalaena Funeraria*. *At. microsc.* 4. tab. 16. fig. 17.

Habitat in Angliae, Europae borealis nemoribus.

1298. P. alis flavis, fascia purpurea. *Syst. Ent.* 645. 134. \* *dentalis*.

Habitat in India orientali. *D. Koenig.*

149. PYRALIS.

- a. P. alis fuscis, fasciis duabus niueis, posteriore vndata. *Bankiana*.

Habitat in Angliae sylvis. *Dom. Teats.*

Magna. Alae anticae fuscae, nitidae litura bascos fasciisque duabus obliquis niueis, posteriore maiore,

S 2 quae



276 CLASS. VI. GLOSSATA. *Pyralis*.

quae versus anteriora dentem emittit. Alae posticae cinereae.

*Leeana*. 2. P. alis pallidis, macula centrali fusca. *Gen. Inf. Mant.* 292. \*

Habitat in Europae nemoribus, frequens in Anglia.

*mundana*. 3. P. alis denudatis albidis, strigis duabus punctoque medio obsoleto fuscis. *Syst. Ent.* 645. 2.

*Phalaena mundana*. *Linn. Syst. Nat.* 2. 812. 17. *Fa. Sv.* 1343.

*Phalaena* antennis filiformibus, alis albedo hyalinis, strigis duabus vndatis fuscis. *Degeer Inf. Vers. Germ.*

2. 1. 353. 9.

*Schaeff. Icon.* tab. 159. fig. 6. 7.

*Hollar. Inf.* tab. 4. fig. 2.

Habitat in Europae Lichenibus.

*Larua* pilosa, brunnea.

*prasinana*. 4. P. alis anticis viridibus, strigis duabus obliquis flauicantibus, posticis albis. *Syst. Ent.* 645. 3.

*Phalaena prasinana* antennis filiformibus, alis desicis viridibus, strigis duabus albis obliquis, posticis albis.

*Degeer Inf. Vers. Germ.* 2. 1. 296. 1.

*Phalaena* spirilinguis, humeris latis, alis viridibus, linea duplici transuersa albida. *Geoff. Inf.* 2. 172. 124.

*Roef. Inf.* 4. tab. 10.

*Degeer Inf.* 1. tab. 3.

*Reaum. Inf.* 1. tab. 39. fig. 13. 14.

*Pod. Inf.* 93. tab. 2. fig. 8. ?

Habitat in Alno, Quercu Europae.

*Larua* viridis, linea laterali flauicante, segmento secundo dorsi tuberculato.

*Puppa* viridis, linea dorsali atra.

*fagana*. 5. P. alis viridibus, strigis tribus obliquis albis, antennarum pedibusque fulvis.

*Phalaena prasinana* alis superioribus viridibus, strigis duabus obliquis flauis, margine postico, antennarum pedibusque fulvis. *Linn. Syst. Nat.* 2. 875. 285. *Fa. Sv.* 1306.

*Ammiral. Inf.* tab. 2. fig. 1.

*Harris Aurel.* tab. 10. fig. 1.

*Petiv.*

*Petiv. Gazoph.* tab. 7. fig. 11.

*Albin. Inf.* tab. 31.

*Roef. Inf.* 4. tab. 22.

*Wilks pap.* 6. tab. 1. a. 13.

Habitat in Fago syluatica Europae.

Affinis praecedenti, attamen distincta.

*Variat* alis margine sanguineo, vti in figura *Roefelii*.  
et *Wilks*, attamen vix distincta.

*Larua* viridis, linea laterali flaua, cauda bifida.

*Puppa* intra folliculum, gibbosum, ferrugineum.

6. *P. alis rhombeis, anticis viridibus immaculatis. Syft. viridana.*

*Ent.* 656. 4.

*Phalaena viridana. Linn. Syft. Nat.* 2. 875. 266.

*Phalaena feticornis spirilinguis, humeris latls, antennis flauescens, alis dilute fuscis. Geoff. Inf.* 2.  
171. 123.

*Frisch. Inf.* 3. tab. 8.

*Reaum. Inf.* 2. tab. 18. fig. 6. 7.

*Roef. Inf.* 1. phal. 4. tab. 1.

*Wilks pap.* 4. tab. 1. n. 5.

*Harris Aurel.* tab. 10. fig. 1. 2.

Habitat in Europae Quercu.

*Larua* nuda, 16poda, viridis, nigro punctata, pedibus  
posticis flauis.

*Puppa* brunnea, postice stylo bifido.

7. *P. alis rhombeis, anticis viridibus, margine albo. Syft. clorana.*

*Ent.* 646. 5.

*Phalaena clorana. Linn. Syft. Nat.* 2. 876. 287. *Fn.*

*Sv.* 1308.

*Degeer Inf. Verf. Germ.* 2. 1. tab. 10. fig. 8. 9.

*Roef. Inf.* 1. phal. 4. tab. 3.

*Reaum. Inf.* 2. tab. 18. fig. 6. 7.

*Frisch. Inf.* 3. 18. tab. 2.

Habitat in Salicibus.

*Larua* virescens, albo punctata, lateribus fusco macu-  
latis.

8. *P. alis fuscis, margine exteriori albido, medio ramum vncana. recuruum album exferente. Syft. Ent.* 646. 7.

*Phalaena vncana. Linn. Syft. Nat.* 2. 875. 284. *Fn. Sv.*  
1305.

*Clerk. phal.* tab. 3. fig. 7.

Habitat in Europa boreali.

Statura et magnitudo fere *Noctuae*.

*plumbana.* 9. *P. alis plumbeis, lineolis tribus maculaque magna apice ferrugineis. Gen. Inf. Mant.* 292. \*

Habitat Hamburgi. *D. Schulz.*

*Hattorfiana.* 10. *P. alis fuscis, strigis cinereis sesquitertiis. Syst. Ent.* 647. 8. \*

Habitat in Germania. *Dom. de Hattorf Hannoneranus.*

*Rosana.* 11. *P. alis anticis testaceis, fascia grisea obliqua. Syst. Ent.* 647. 9.

*Phalaena Rosana. Linn. Syst. Nat.* 2. 876. 293. *Fa. Sv.* 1314.

*Phalaena antennae filiformibus, alis testaceis, strigis maculisque obscurioribus, posticis fuscis. Degener Inf. Vers. Germ.* 2. 1. 344. 1.

*Merian. Europ.* 1. 7. tab. 19.

*Reaum. Inf.* 2. tab. 15. fig. 5 — 9.

*Blanc. Inf.* tab. 8. fig. E — H.

Habitat in Rosa, Berberide.

*pallidana.* 12. *P. alis pallide flavis, strigis duabus obliquis ferrugineis. Gen. Inf. Mant.* 292. \*

Habitat Hamburgi. *D. Schulz.*

*Smeathmanniana.* 13. *P. alis exalbidis, fasciis duabus obliquis fuscis, anteriore abbreviata, posteriore interrupta. Habitat in Anglia. Mus. Dom. Teats.*

Statura *P. humanae*. Alae anticae albae, in medio fascia obliqua fusca, quae tamen marginem exteriorem haud attingit. Pone hunc fascia altera in medio interrupta. Alae posticae obscurae.

*decussana.* 14. *P. alis cinereo nigro argenteoque tenuissime fasciatis, apice macula subocellari cinerea. Syst. Ent.* 647. 10. \*

Habitat in Lipsiae hortis.

*ameriana.* 15. *P. alis anticis testaceis, litura communi ferruginea. Syst. Ent.* 647. 11.

*Phalaena Americana*. Linn. *Syst. Nat.* 2. 877. 298. *Fn.*  
*Sv.* 1310.

*Reaum. Inf.* 2. tab. 18. fig. 6. 7.

Habitat in Salice.

16. *P.* alis anticis fuscis, maculis quatuor dorsalibus atro ferrugineis. *Syst. Ent.* 647. 12. \* *maculana.*

Habitat in Sueciae nemoribus.

17. *P.* alis anticis fuscis immaculatis. *Syst. Ent.* 648. 13. *fuscana.*  
*Phalaena fuscana*. Linn. *Fn. Sv.* 1311.

Habitat in Europae borealis nemoribus.

18. *P.* alis anticis obscure cinereis, maculis duabus marginalibus fuscis. *Forstera-*

Habitat in Angliae nemoribus. *Dom. Teats.*

Caput et thorax fusca, immaculata. Alae anticae obscure cinereae atomis vix conspicuis fuscis subundatae maculisque duabus magnis marginalibus, fuscis.

19. *P.* alis anticis testaceis, fascia obliqua fusca. *Syst. Ent.* *Xylostea-*  
648. 14. *na.*

*Phalaena Xylosteana* alis superioribus testaceis, cruce communi ferruginea. Linn. *Syst. Nat.* 2. 876. 295.  
*Fn. Sv.* 1313.

*Phalaena* antennis filiformibus, alis testaceis, fascia obliqua fusca. *Degeer Inf. Vers. Germ.* 2. 1. 345. 2.  
*Degeer Inf.* 1. tab. 27. fig. 9. 10.

Habitat in Europa boreali.

20. *P.* alis anticis testaceis, margine postico rufo, medio ministrana.

litura ferruginea, lineola alba. *Syst. Ent.* 648. 15.  
*Phalaena ministrana*. Linn. *Syst. Nat.* 2. 877. 300. *Fn.*  
*Sv.* 1131.

*Clerk. phal.* tab. 2. fig. 12.

Habitat in Europae borealis hortis.

1. *P.* alis anticis fuscescentibus, macula centrali fulva, servana.

basi flavis. *Gen. Inf. Mant.* 292. \*

Habitat in Germania. *Dom. de Hattorf.*

2. *P.* alis cinereis, macula brunnea communi transversa. *Vdman-*  
*Syst. Ent.* 648. 16. *niana.*

*Phalaena Vdmanniana*. Linn. *Syst. Nat.* 2. 880. 32.

*Fn. Sv.* 1332.

Habitat in Europa boreali.

*Solan-  
driana*.

23. *P. alis pallidis albo subfasciatis, macula communi dor-  
sali ferruginea.* *Syst. Ent.* 648. 17.

*Phalaena Solandriana* alis superioribus pallidis, ma-  
cula communi dorsali ferruginea. Linn. *Syst. Nat.*

2. 878. 310. *Fn. Sv.* 1317.

*Phalaena* antennis filiformibus, alis pallide testaceis,  
feminae macula triangulari alba, maris macula fer-  
ruginea marginis tenuioris. *Degeer Inf. Vers. Germ.*  
2. 1. 358. 3.

*Degeer Inf.* 1. tab. 28. fig. 29. 30.

Habitat in Europae borealis Betula.

*Larva* cinerea, nigro irrorata, capite primoque corpo-  
ris segmento fuscis.

*Stroemia*-24. *P. alis fuscis, maculis duabus dorsalibus cinereis.*

*na.*

Habitat in Germania. *Dom. de Hattorf.*

Statura praecedentium. Alae anticae testaceo fuscae ma-  
culis duabus magnis cinereis ad marginem tenuio-  
rem. Margo exterior albo striatus. Alae posticae fuscae.

*Zoëgana*. 25. *P. alis flavis, puncto medio lituraque postica ferrugi-  
neis.* *Syst. Ent.* 648. 18.

*Phalaena Zoëgana* alis flavis puncto medio ferrugi-  
neo, postice ferrugineis macula flava: Linn. *Syst.*  
*Nat.* 2. 876. 289.

*Atk. Soc. Berol. nat.* 3. tab. 1. fig. 9.

*Clerk. phal.* tab. 4. fig. 4.

β. *Phalaena hamana* alis superioribus flavis puncto  
lituraque postica hamata ferrugineis. Linn. *Syst. Nat.*

2. 876. 290. *Fn. Sv.* 1309.

*Harris Inf. angl.* tab. 1. fig. 1. 2.

*Clerk. phal.* tab. 4. fig. 4. 6.

Habitat in Europa boreali.

*cruciana*. 26. *P. alis ex albido cinereis, postice macula arcuata fer-  
ruginea.* *Syst. Ent.* 649. 19.

*Phalaena cruciana*. Linn. *Syst. Nat.* 2. 880. 323. *Fn. Sv.*  
1333.

Habitat in Europae borealis nemoribus.

17. P. alis exalbido cinereis immaculatis, capite fuluo. *cinerana*.  
Habitat in Germania. *Dom. de Hattorf*.  
Nimis praecedenti affinis, at alae immaculatae.
18. P. alis cinereis, fasciis obliquis fulvis, albo marginatis. *obliquana*.  
Habitat in Anglia. *Mus. Dom. Teats*.  
Magna. Antennae subpectinatae, fuscae. Caput et thorax ferruginea. Alae anticae cinereae fasciis tribus obliquis fulvis margine omni albo. Fascia prima bascos, secunda latissima a medio marginis crassioris ad angulum ani inaequalis, tertia apicis minor obliqua.
19. P. alis anticis fusciscentibus, margine exteriori bascos niueo nigro punctato. *lactana*.  
*Syst. Ent. 649. 20.\**  
Habitat in Sueciae nemoribus.
30. P. alis fuscis, macula communi dorsali rhombea alba. *Brunnichana*.  
*Syst. Ent. 649. 21.*  
*Phalaena Brunnichana. Linn. Syst. Nat. 2. 380. 319.*  
Habitat in Europae borealis hortis.
31. P. alis luteis, fasciis tribus arcuatis maculaque disci atra, punctis tribus argenteis. *arcuana*.  
*Syst. Ent. 649. 22.*  
*Phalaena arcuana* alis luteis, fasciis tribus arcuatis, macula nigra tripunctata lineolisque duabus argenteis. *Linn. Syst. Nat. 2. 877. 296. Fn. Sv. 1317.*  
*Clerk. phal. tab. 10. fig. 8.*  
Habitat in Europae borealis Coryletis.
32. P. alis anticis luteis, characteribus I. L. thoraceque argenteis. *Lecheana*.  
*Syst. Ent. 649. 23.*  
*Phalaena Lecheana. Linn. Syst. Nat. 2. 877. 301. Fn. Sv. 1318.*  
*Clerk. phal. tab. 10. fig. 2.*  
Habitat in Europae borealis Pado.
33. P. alis anticis flauissimis, venis sanguineis reticulatis. *Christierna*.  
*Syst. Ent. 649. 24.*  
*Phalaena Christierna. Linn. Syst. Nat. 2. 877. 303.*  
*Schaeff. Icon. tab. 2. fig. 12.*  
Habitat in Europa boreali, in Anglia.
34. P. alis anticis testaceis, fasciis fuscis sesquitertiis. *auellana*.  
*Ent. 650. 25.*

283 CLASS. VI. GLOSSATA. Pyralis.

*Phalaena anellana*. Linn. *Syst. Nat.* 2. 877. 297. *Fa. Sv.* 1313.

Habitat in Europae borealis Corylo, Pyro.

*fasciana*. 35. P. alis anticis obscure cinereis, fascia media lituraque apicis fuscis.

Habitat in Angliae hortis. *Dom. Testis.*

Corpus statura et magnitudine P. anellanae. Alae anticae obscure cinerae atomis plurimis fuscis. In medio fascia lata sinuata fusca et in margine postico litura fusca. Alae posticae cinerae.

*Hastiana*. 36. P. alis anticis fuscis, fascia antrorum obliqua alba. *Syst. Ent.* 650. 26.

*Phalaena Hastiana*. Linn. *Syst. Nat.* 2. 878. 311. *Fa. Sv.* 1328.

*Clerk. phal.* tab. 2. fig. 7.

Habitat in Europae borealis Salice.

*Lediana*. 37. P. alis fuscis, fasciis duabus rubro argenteis, posteriore puncto intruso. *Syst. Ent.* 650. 27.

*Phalaena Lediana*. Linn. *Syst. Nat.* 2. 879. 313. *Fa. Sv.* 1330.

Habitat in Europae borealis Ledo.

*Schalleriana*. 38. P. alis canis, macula marginali ferruginea. *Syst. Ent.* 650. 28.

*Phalaena Schalleriana* alis canis, macula laterali ferruginea, apicem alae attingente. Linn. *Syst. Nat.* 2. 879. 315.

Habitat in Europa.

*Modaeriana*. 39. P. alis anticis flavescentibus, macula marginali posteriorius fusca. *Syst. Ent.* 650. 29.

*Phalaena Modaeriana*: Linn. *Syst. Nat.* 2. 880. 31. *Fa. Sv.* 1331.

Habitat in Europa boreali.

*Schreberiana*. 40. P. alis anticis griseis, macula marginali trigona alba. *Syst. Ent.* 650. 30.

*Phalaena Schreberiana*. Linn. *Syst. Nat.* 2. 976. 316. *Fa. Sv.* 1332.

Habitat in Europae borealis nemoribus.

1. P. alis anticis luteo ferrugineis, macula marginali tri- *Holmia-*  
gona alba. *Syst. Ent.* 650. 31. *na.*  
*Phalaena Holmiana.* Linn. *Syst. Nat.* 2. 878. 308. *Fn.*  
*Sv.* 1305.  
*Clerk. phal.* tab. 10. fig. 7.  
Habitat in Pomonae arboribus.
2. P. alis anticis flauis, fascia obliqua testacea maculaque *gnomana.*  
postica marginali ferruginea. *Syst. Ent.* 651. 32.  
*Phalaena gnomana.* Linn. *Syst. Nat.* 2. 876. 294. *Fn.*  
*Sv.* 1337.  
*Clerk. phal.* tab. 4. fig. 13.  
Habitat in Europae borealis nemoribus.
3. P. alis anticis ferrugineis fusco maculatis reticulatis- *oporana.*  
que. *Syst. Ent.* 651. 33.  
*Phalaena oporana.* Linn. *Syst. Nat.* 2. 876. 292. *Fn.*  
*Sv.* 1315.  
*Wilk. pap.* 17. tab. 2. a. 4.  
Habitat in Pomonae arboribus.
14. P. alis anticis fusco cinereis, punctis fuscis, centrali *Ilucana.*  
solitario atro.  
Habitat in Angliae hortis. *Mus. Dom. Testis.*  
Magna in hoc genere. Alae anticae latae fusco cinereae  
fere vndatae. Ante medium puncta duo approxima-  
ta fusca, tunc punctum centrale solitarium atrum.  
Versus marginem posticum puncta duo siue tria ap-  
proximata fusca. Margo ipse striga punctorum fu-  
scorum. Alae posticae cinereae.
45. P. alis anticis canis, atomis nigris sparsis, posticis ex- *Logiana.*  
albidis. *Syst. Ent.* 651. 34.  
*Phalaena Logiana.* Linn. *Syst. Nat.* 2. 879. 317. *Fn.*  
*Sv.* 1336.  
*Clerk. phal.* tab. 10. fig. 3.  
Habitat in Europa boreali.
16. P. alis anticis niueis, subtus fuscis. *Syst. Ent.* 651. 35. *Gonana.*  
*Phalaena Gonana.* Linn. *Syst. Nat.* 2. 879. 318. *Fn.*  
*Sv.* 1342.  
Habitat in Europae borealis hortis.



- Turnbergiana.* 47. P. alis anticis cinereis, punctis quatuor eleuato fasciculatis atris.  
Habitat in Germania. *Dom. de Hattorf.*  
Statura et magnitudo P. asperanae. Corpus totum fuscum, immaculatum. Alae obscurae, minime nitidae punctis quatuor per paria approximatis eleuato fasciculatis nigris.
- asperana.* 48. P. alis anticis basi albis, apice fuscis scabris. *Gen. Inf. Mant.* 292\*.  
Habitat Hamburgi. *Dr. Schulz.* Kiliae, *Dom. de Schefstedt.*
- duplana.* 49. P. alis anticis pallidis, punctis duobus approximatis bascos nigris. *Gen. Inf. Mant.* 293\*.  
Habitat Kiliae. *Dom. de Schefstedt.*
- squamana.* 50. P. alis virescentibus scabris. *Syst. Ent.* 651. 36\*.  
Habitat in Anglia, *Mus. Monson*, in Quercu Kiloniae, *Dom. de Schefstedt.*  
Puncta duo eleuata atra, alterum versus basin, alterum in centro.
- literana.* 51. P. alis rhombeis, anticis viridibus, characteribus atris *Syst. Ent.* 646. 6.  
*Phalaena literana.* *Linn. Syst. Nat.* 2. 876. 288.  
Habitat in Germaniae nemoribus.  
Alae anticae scabrae.
- emargana.* 52. P. alis subcaudatis flauis fusco reticulatis fasciaque lata fusca, margine crassiori late emarginato. *Syst. Ent.* 651. 37\*.  
Habitat in Angliae hortis. *Dom. Lee.*  
Aliam vidi simillimam, at absque emarginatura alae.
- caudana.* 53. P. alis subcaudatis griseis, margine interiori fuluo exteriori emarginato. *Syst. Ent.* 38\*.  
Habitat in Angliae hortis.
- frabrana.* 54. P. alis anticis cinereis fusco subfasciatis scabris, margine exteriori postice exciso.  
Habitat in Angliae Quercu. *Mus. Dom. Teats.*

Statura et magnitudo praecedentium. Corpus cinereum.  
Alae anticae cinereae punctis eleuatis scabrae, posticae strigis vndatis fuscis subfasciatae. Margo exterior a medio ad apicem contrahitur, vt quasi excisus.  
Alae posticae cinereae.

5. P. alis anticis flauis, maculis duabus costalibus ful-phureis. *Syst. Ent. 652. 39.\**

*Phalaena fagana. Wien. Verz. 28. 7. tab. 1. a. b.*  
— — — — — 1. b. b.

Habitat in Saxoniae Quercu.

6. P. alis cinereis, litura media sanguinea, puncto al-bido. *Syst. Ent. 652. 40.\**

Habitat in Anglia. *Mus. Dom. Banks.*

7. P. alis anticis luteis ferrugineo reticulatis, medio li-tura fuscescente. *Syst. Ent. 652. 41.* *Forskah-liana.*

*Phalaena Forskahliana. Linn. Syst. Nat. 2. 878. 304.*  
*Fn. Sv. 1319.*

*Clerk. phal. tab. 10. fig. 2. 6.*

Habitat in Europae Rosa.

Alae subtus laete flavae.

8. P. alis anticis flauis luteo reticulatis duplici xx nota-tis. *Syst. Ent. 652. 42.* *Loeflingiana.*

*Phalaena Loeflingiana. Linn. Syst. Nat. 2. 878. 305.*  
*Fn. Sv. 1323.*

*Clerk. phal. tab. 10. fig. 6.*

Habitat in Europae hortis.

Praecedenti affinis, at maior et alae subtus fuscae.

9. P. alis anticis luteis flauo punctatis, fasciis quatuor ar-genteis, tertia bifida. *Syst. Ent. 652. 43.* *Bergmanniana.*

*Phalaena Bergmanniana. Linn. Syst. Nat. 2. 878. 307.*  
*Fn. Sv. 1314.*

*Phalaena* antennis filiformibus, alis luteis nitidis, strigis quatuor argenteis. *Degeer Inf. Vers. Germ. 2. 1. 346. 4.*

*Clerk. phal. tab. 10. fig. 5.*

Habitat in Europae borealis hortis.

*Testiana*. 60. *P.* alis anticis griseis, macula punctisque duobus centralibus nigris.

Habitat in Angliæ nemoribus.

Paullo maior *P. Alstroemianæ*. Corpus cinereum antennis nigris. Alæ antice griseae macula centrali punctisque duobus oblique positis approximatis nigris. Puncta quatuor s. quinque nigra ad marginem posticum.

*Alstroemiana*. 61. *P.* alis albidis, anticis macula media laterali fusco ferruginea. *Syst. Ent.* 653. 44.

*Phalaena Alstroemiana*. *Linn. Syst. Nat.* 2. 879. 314. *Fn. Sv.* 1335.

*Clerk. phal.* tab. 10. fig. 1.

Habitat in Europæ borealis nemoribus.

*Fabriciana*. 62. *P.* alis anticis nigricantibus, anticis pari punctorum albidorum. *Syst. Ent.* 553. 45.

*Phalaena Fabriciana*. *Linn. Syst. Nat.* 2. 880. 324.

Habitat in Europæ borealis Pruno spinosa in sepibus frequens.

*Conwangana*. 63. *P.* alis ferrugineo fuscis, argenteo punctatis, maculae dorsali flauicante. *Syst. Ent.* 653. 46.\*

Habitat in Angliæ nemoribus capta d. XXVI. Iun. prope Oxoniam.

*Scheffeldiana*. 64. *P.* alis atris, macula magna communi, strigis atomisque coeruleiscenti argenteis. *Gen. Ins. Mant.* 193.\*

Habitat Kiliae in Alnetis. *Dona. de Scheffeldt.*

*Koenigiana*. 65. *P.* alis fulvis, apice nigris fulvo maculatis. *Syst. Ent.* 653. 47.\*

Habitat in India orientali. *D. Koenig.*

*aurana*. 66. *P.* alis fuscis, maculis duabus aureis. *Syst. Ent.* 653. 48.\*

Habitat in Angliæ nemoribus. *Mus. Dona. Banks.*

*pomonæ*. 67. *P.* alis nebulosis, postice macula rubra aurea. *Syst. Ent.* 653. 49.

*Phalaena Pomonæ*. *Linn. Syst. Nat.* 2. 892. 401. *Fn. Sv.* 1412.

*Frijth*

*Frisch. Inf.* 7. tab. 13.

*Roef. Inf.* 1. phal. 4. tab. 10.

*Reaum. Inf.* 2. tab. 40. fig. 9. 10.

*Wilks pap.* 5. tab. 1. a. 9.

Habitat inter Europae Poma.

*Larva* 16poda, nuda, rufa, capite nigro.

*Puppa* folliculata, brunnea.

68. P. alis testaceis, fasciis punctisque argenteis, medio *Schulzia-*  
litura atra. *Gen. Inf. Mant.* 293.\* *na.*

Habitat in Germania. *D. Schulz.*

69. P. alis fuscis, fasciis tribus cinereis, secunda intror-  
sum bifida. *Syst. Ent.* 654. 50.\*

Habitat in Sueciae nemoribus.

70. P. alis cinereis subfasciatis, apice acuminato recurvatis, *unguica-*  
margine exteriori postico transverse albo striatis. *na.*  
*Syst. Ent.* 654. 51.\*

*Phalaena unguicella.* *Linn. Syst. Nat.* 2. 887. 365. *Fn.*  
*Sv.* 1382.

*Clerk. phal.* tab. 12. fig. 7.

Habitat in Europae borealis nemoribus.

71. P. alis fuscis, arcu dorsali albo apiceque strigis argen-  
teis. *Syst. Ent.* 654. 52.\*

*Tinea* alis nigris, superioribus macula quadrata alba,  
ex quatuor lineolis albis transversis composita. *Geoff.*  
*Inf.* 2. 189. 16.

Habitat in Europae hortis.

72. P. alis flavescentibus, apice strigis ferrugineis argen-  
teisque, puncto triplici atro. *Syst. Ent.* 654. 53.\*

Habitat in Sueciae nemoribus.

73. P. alis basi fuscis linea semicirculari pallida, apice aura-  
tis, margine crassiori nigro pallidoque striato. *Gen.*  
*Inf. Mant.* 294.\*

Habitat in Kiliae Quercu. *Dom. Lund.*

74. P. alis anticis fuscis, apice albis. *Syst. Ent.* 654. 54. *Cynosta-*  
*Pha- na.*

288 CLASS. VI. GLOSSATA. *Pyralis*.

*Phalaena Cynosbatella*. Linn. *Syst. Nat.* 2. 887. 368.  
Fn. Sv. 1379.

*Phalaena*, antennis filiformibus laevis, alis deflexis,  
anticis fuscis, postice albis fusco maculatis. *Deger*  
*Inf. Vers. Germ.* 2. 1. 359. 5.

*Degeer Inf.* 1. tab. 34. fig. 4. 5.

*Merian. Europ.* tab. 28.

Habitat in Rosae Gemmis.

*Larva* 16poda vtrinque attenuata, ferruginea, capite  
nigro.

*Puppa* folliculata, brunnea.

Ala terminatur puncto atro.

*resinana*. 75. P. alis fuscis argenteo subfasciatis. *Syst. Ent.* 654. 55.

*Phalaena resinella*. Linn. *Syst. Nat.* 2. 894. 206. Fn.  
Sv. 1423.

*Phalaena* antennis filiformibus, alis deflexis griseis,  
fasciis maculisque griseis. *Deger Inf. Vers. Germ.*  
2. 1. 303. 10.

*Frisch. Inf.* 10. tab. 9.

*Deger Inf.* 1. tab. 33. fig. 12. 13.

*Roef. Inf.* 1. phal. 4. tab. 16.

Habitat intra Resinae glomerem exsudantem e vulnere  
ramorum Pini.

*Larva* 16poda, nuda, lutea, capite brunneo.

*Puppa* folliculata, fusca.

*applanata*. 76. P. alis depressis fuscis, punctis tribus centralibus al-  
bis. *Gen. Inf. Mant.* 294. \*

Habitat Kiloniae prope Grez. *Dom. de Schestedt*.

*Heraclea*. 77. P. alis depressis griseis, anticis subtus fuscis. *Syst. Ent.*  
na. 655. 56.

*Phalaena Heracleana* clinguis recurvicornis, alis gri-  
seis, puncto fusco, thorace flavo. Linn. *Syst. Nat.* 2.  
880. 326. Fn. Sv. 1354.

*Reaum. Inf.* 2. tab. 6. fig. 1 — 4.

*Deger Inf.* 1. tab. 29. fig. 6. 7.

*Schaeff. Icon.* tab. 2. fig. 3. 4.

Habitat in umbellis plantarum umbelliferarum in pri-  
mis Heraclei, contortis destructisque floribus caules  
ex alis foliorum penetrans. Linn.

*Larva*

*Larva viridis, nigro punctata; lineis dorsalibus tribus obscurioribus.*

71. P. alis depressis fuscis, capite thoraceque niuis. *Syst. depressa-  
Ent. 655. 57.\**  
Habitat in Sueclae floribus. *na.*

# 150. TINEA.

1. T. alis griseis, postice emarginatis, dorso canaliculato *cerealis*.  
fusco. *Syst. Ent. 655. 1.\**

*Phalaena cereana* alis griseis, arcu dorsali complanato exasperato carinatis, scutello apice albo. *Linn. Syst. Nat. 2. 874. 282.*

*Reaum. Inf. 3. tab. 19. fig. 14. 15.*

*Blom. Act. Holm. 1764. tab. 2.*

Habitat in Apum alucaribus, ceram depascens.

2. T. alis oblongis cinereis, punctis duobus atris ante *colonella*.  
strigam curuam vndulatam obsoletam. *Syst. Ent. 655. 2.*

*Phalaena colonella. Linn. Syst. Nat. 2. 883. 346. An. Su. 1358.*

*Phalaena* antennis filiformibus, alis conuolutis achatinis, strigis duabus vndatis punctoque centrali nigris. *Deger Inf. Verf. Germ. 2. 1. 343. 5.*

*Clerk. phal. tab. 3. fig. 8.*

Habitat in Europa boreali.

3. T. alis oblongis cinereis antice albidis, postice pallidis, striga vndata nigra. *Syst. Ent. 656. 3.*

*Phalaena sociella* alis oblongis cinereis antice albidis, postice striga pallida. *Linn. Syst. Nat. 2. 883. 345. Fn. Su. 1349.*

*Phalaena* antennis filiformibus, alis conuolutis cinereis, strigis obliquis maculisque ferrugineis. *Deger Inf. Verf. Germ. 2. 1. 343. 4.*

*Clerk. phal. tab. 3. fig. 11.*

Habitat in Apis lapidariae fauis.

4. T. alis oblongis griseis, antennarum articulis duobus inferioribus incrassatis. *Gen. Inf. Mant. 294.\**

Habitat Kiliae. *Dom. Land.*

*Fabr. Sp. Inf. T. II.*

T

5. T.

*gelatella*. 5. T. alis fulco cinereis, anticis vitta alba.

*Phalaena gelatella* alis cinereis, superioribus linea alida punctisque duobus nigris remotis, femina aptera. *Linn. Syst. Nat.* 2. 883. 344. *Fn. Sv.* 1450.

*Clerk. phal.* tab. 8. fig. 3.

Habitat in Europae borealis pomariis.

Femina rudimentis tantum alarum.

*euonymel*. 6. T. alis anticis niueis, punctis 50 nigris. *Syst. Ent.* 656. 4.

*Phalaena euonymella*. *Linn. Syst. Nat.* 2. 885. 350. *Fn. Sv.* 1363.

*Tinea* alis superioribus albis, punctis nigris, inferioribus fuscis. *Geoff. Inf.* 2. 183. 4.

*Frisch. Inf.* 3. tab. 16.

*Merian. Europ.* 2. tab. 2.

*Alb. Inf.* tab. 60. fig. A. C.

*Reaum. Inf.* 2. tab. 13. fig. 12. 13.

*Roef. Inf.* 2. phal. 4. tab. 8.

*Wilks pap.* 4. tab. 1. a. c.

*Sulz. Inf.* tab. 16. fig. 99.

Habitat in Euonymo, Pado, Sorbo.

*Larva* gregaria, nuda, flaua, nigro punctata, sub tentorio communi.

*Puppa* folliculata, brunnea.

*Varietas* maior Americana punctis paucioribus maioribus in *Mus. Dom. Teats*.

*padella*. 7. T. alis anticis plumbeis, punctis 20 nigris, *Syst. Ent.* 656. 5.

*Phalaena padella*. *Linn. Syst. Nat.* 2. 885. 351. *Fn. Sv.* 1364.

*Phalaena* antennis filiformibus, alis cinereis, punctis 20 nigris. *Degeer Inf. Vers. Germ.* 2. 1. 354. 1. tab. 10. fig. 14. 15. 16.

*Frisch. Inf.* 5. 38. tab. 16. fig. 1.

*Reaum. Inf.* 2. tab. 12. fig. 5 — 9.

*Roef. Inf.* 1. phal. 4. tab. 7.

*Harris Aurel.* 7. tab. 3. fig. k. m.

*Schaeff. Icon.* tab. 145. fig. 2. 3.

Habitat in Pomonae arboribus.

*Larus*

*Larva* gregaria, nuda, grisea, segmentis vtrinque puncto atro, sub tentorio communi.

*Puppa* folliculata, flavicans.

- l. T. alis anticis albis, punctis lineaque communi dentato atra, posticis nigris. *Syst. Ent.* 656. 6. *pascella.*

*Phalaena pascella.* Linn. *Syst. Nat.* 2. 334. 347.

*Sulz. Hist. Inf.* tab. 23. fig. 9.

Habitat in Germaniae arvis.

- p. T. alis anticis albis, linea communi sinuato repanda maculisque lateralibus albis. *Syst. Ent.* 657. 7. *sequella.*

*Phalaena sequella.* Linn. *Syst. Nat.* 2. 335. 352. *Fn.*

*Sv.* 1365.

*Clerk. phal.* tab. 10. fig. 14.

Habitat in Europae boreali.

10. T. alis niueis, fascia punctisque numerosis fuscis. *cornella.*  
*Syst. Ent.* 657. 8.<sup>o</sup>

Habitat in Saxoniae Corno.

11. T. alis luteis, anticis punctis nigris trifariis. *Syst. irrorella.*  
*Ent.* 657. 9.

*Phalaena irrorella.* Linn. *Syst. Nat.* 2. 335. 354. *Fn.*

*Sv.* 1361.

*Phalaena* antennis pectinatis, alis deflexis rotundatis luteis nigro punctatis, corpore nigro, ano luteo.

*Deger Inf. Vers. Germ.* 2. 1. 248. 6. tab. 5. fig. 12.

*Clerk. phal.* tab. 4. fig. 5.

*Sulz. Hist. Inf.* tab. 23. fig. 10.

Habitat in Europae borealis sylvis.

12. T. alis albidis fusco lineatis, apice strigis duabus *linetella.*  
fuscis.

Habitat in agro Pedemontano. *D. Allioni.*

Statura praecedentium. Palpi valde porrecti. Margo posticus, ciliaris.

13. T. alis supra luteis immaculatis, subtus fuscis. *Syst. lutarella.*  
*Ent.* 657. 10.

*Phalaena lutarella* alis corporeque luteis immaculatis.

*Linn. Syst. Nat.* 2. 336. 355. *Fn. Sv.* 1360.



*Clerk. phal. tab. 4. fig. 9.*

Habitat in Europae borealis nemoribus.

*mesomella.* 14. T. alis anticis pallidis, margine flavo punctisque duobus fuscis.

*Tinea mesomella* alis supra albis, subtus luteis, internis fuscis. *Syst. Ent.* 657. 11.

*Phalaena mesomella.* *Linn. Syst. Nat.* 2. 886. 356. *Fa. Sv.* 1362.

*Clerk. phal. tab. 4. fig. 14.*

Habitat in Europae gramine.

Punctum paruum ad marginem exteriorem et alterum ad marginem interiorem oppositum.

*Oxyacanthella.* 15. T. alis fusco nebulosis, friga albida margineque postico albo. *Syst. Ent.* 657. 12.

*Phalaena Oxyacanthella.* *Linn. Syst. Nat.* 2. 886. 357.

Habitat in Crataego Oxyacantha.

*Pinetella.* 16. T. alis anticis flavis, maculis duabus albissimis, anteriore oblonga, posteriore ovata. *Syst. Ent.* 657. 13.

*Phalaena Pinetella.* *Linn. Syst. Nat.* 2. 886. 358. *Fa. Sv.* 1368.

*Clerk. phal. tab. 4. fig. 15.*

Habitat in Europae Pinetis.

*pratella.* 17. T. alis anticis cinereis, linea albissima, postice ramosa, apice striis obliquis. *Syst. Ent.* 658. 14.

*Phalaena pratella.* *Linn. Syst. Nat.* 2. 886. 360. *Fa. Sv.* 1369.

*Phalaena* antennis filiformibus, alis convolutis argenteis, subtus griseis. *Degeer Inf. Vers. Germ.* 2. 1. 341. 1.

*Clerk. phal. tab. 3. fig. 14.*

Habitat in Europae borealis pratis graminosis frequens.

*pascuella.* 18. T. alis cinereis, linea albissima, margine postico nigro punctato. *Syst. Ent.* 658. 15.

*Phalaena pascuella*, alis superioribus pallidis, linea maculisque duabus oblongis albissimis. *Linn. Syst. Nat.* 2. 886. 359. *Fa. Sv.* 1367.

*Phalar-*

*Phalaena antennis filiformibus, alis tenuioribus albidis, lineis longitudinalibus argenteis, palpis oblongis.*

*Degeer Inf. Verf. Germ. 2. 1. 343. 2.*

Habitat in Europae borealis pascuis frequens.

19. T. alis cinereis, linea unica albiſſima abbreviata. *Syst. culmella.*  
*Ent. 658. 16.*

*Phalaena culmella* alis superioribus albidis, excavato striatis, punctis nigris septem alisque nitentibus.

*Linn. Syst. Nat. 2. 886. 361. Fn. Sv. 1370.*

*Phalaena minor* pascuorum. *Raj. Inf. 229. 82.*

*Reaum. Inf. 1. tab. 17. fig. 13. 14.*

Habitat in Europa graminosis frequens.

20. T. argentea, antennis fusco annulatis. *Syst. Ent. argentella.*  
*658. 17.*

*Phalaena argentella.* *Linn. Syst. Nat. 2. 895. 421.*

*Fn. Sv. 1432.*

*Phalaena antennis filiformibus, alis argenteis, cauda lutea.* *Degeer Inf. Verf. Germ. 2. 1. 357. 2. tab. 10. fig. 19.*

Habitat in Rubo, Vrtica.

*Larva* gregaria, glabra, fusca.

*Puppa* brunnea, folliculata, folliculo siueo.

21. P. alis anticis flavis, lateribus sanguineis. *Syst. Ent. carnella.*  
*658. 18.*

*Phalaena carnella.* *Linn. Syst. Nat. 2. 887. 363.*

*Sulz. Hist. Inf. tab. 23. fig. 12.*

Habitat in Germaniae pratis.

22. T. alis anticis cinereo flavescentibus, striga marginali argentea. *strigella.*

Habitat in Germania.

Statura praecedentium. Palpi porrecti, compressi.

Alae anticae flavescentes, immaculatae sola striga marginali argentea. Posticae fusco cinereae striga eadem marginali argentea.

23. T. alis anticis fuscis, margine crassiori albo, palpis marginali porrectis incrassatis. *la.*

Habitat in Germania. *Dom. de Hattorf.*

294 CLASS. VI. GLOSSATA. Tinea.

Minor *T. carnella*. Alae anticae fuscae, margine triflori niveo, postico ciliato pallido. Puncta duo minutissima in medio disci. Alae posticae cinercae.

*sigrella*. 24. *T. nigra*, alis fuscis, abdomine rufo. *Syst. Ent.* 658. 19.\*  
Habitat in Daniae nemoribus.

*cryptella*. 25. *T. alis* griseis, linea dimidiata albissima strigaeque postica vndata argentea. *Syst. Ent.* 658. 20.\*  
Habitat in India orientali. *D. Koenig.*

*gramella*. 26. *T. alis* cinereis, strigis duabus angulatis fuscis argenteaeque marginali.

Habitat in Anglia. *Mus. Dom. Teats.*

Paullo minor *P. culmella*. Alae anticae griseae strigis duabus fuscis, quae in medio angulum formant. Margo posticus terminatur striga argentea.

*salicella*. 27. *T. alis* coerulescenti cinereis, latere dorsali punctis albis, thorace cristato. *Syst. Ent.* 659. 21.

*Phalaena Salicella*. *Linn. Syst. Nat.* 2. 887. 367. *Fs. Sv.* 1378.

*Roef. Inf.* 1. phal. 4. tab. 9.

*Wilks pap.* 5. tab. 1. a. 10.

Habitat in Europae Salice.

*Larva* nuda, 16poda, fascia albo punctata.

*Puppa* antice nigra, postice brunnea.

*scalacella*. 28. *T. alis* coerulescenti cinereis, punctis vitta media maculisque rhombeis contiguis atris.

*Phalaena scalacella*. *Acta Soc. Berol. nat. curios.* 3. tab. 1. fig. 6.

Habitat in Germaniae Salice.

Maiuscula in hoc genere. Alae anticae cinereae vitta longitudinali media atra et maculae septem rhombae vittae contiguae. Margo atro punctatus.

*foenella*. 29. *T. alis* fuscis, maculis albidissimis, anteriore falcata, posteriori reniformi. *Syst. Ent.* 659. 22.

*Phalaena foenella*. *Linn. Syst. Nat.* 2. 888. 369. *Fs. Sv.* 1375.

*Clerk. phal.* tab. 2. fig. 13.

Habitat in Europae plantis.

30. T. alis nigris, postice albis, capite niueo. *Syst. Ent. tapezella*.  
659. 23.  
a. *Tinea nigra*, capite niueo, alis pone albis. *Geoff.*  
*Inf.* 2. 187. 13.  
*Phalaena tapezella*. *Linn. Syst. Nat.* 2. 888. 371. *Fn.*  
*Sv.* 1414.  
*Tinea vestiura* maior. *Raj. Inf.* 204. 98.  
*Reaum. Inf.* 3. tab. 20. fig. 2. 4.  
*Clerk. phal.* tab. 11. fig. 12.  
ß. *Tinea fusca*, capite fusco, alis pone albis. *Geoff. Inf.*  
2. 188. 14.  
Habitat in Europae tapetibus, pellibus, intra cylindros  
meatus rodens.
31. T. alis canis, medio puncto nigro, capite griseo. *pellionel-*  
*Syst. Ent.* 659. 24. *la.*  
*Phalaena pellionella*. *Linn. Syst. Nat.* 2. 888. 372. *Fn.*  
*Sv.* 1415.  
*Phalaena plumbea*, nitida puncto nigro in medio ala-  
rum. *Geoff. Inf.* 2. 184. 6.  
*Roef. Inf.* 1. phal. 4. tab. 17.  
*Reaum. Inf.* 3. tab. 6. fig. 12. 16.  
Habitat in Europae vestimentis pilosis.  
*Larua* albicans linea dorsali rubra intra cucullum la-  
titans.  
*Puppa* flauescens.
32. T. alis cinereis, costa albis, apicibus adscendentibus *vestianel-*  
plumosis. *Syst. Ent.* 660. 25. *la.*  
*Phalaena vestianella*. *Linn. Syst. Nat.* 2. 888. 370. *Fn.*  
*Sv.* 1417.  
*Scop. Carn.* 669.  
Habitat in vestimentis, quae misere destruit.
33. T. alis cinereis, thorace vtrinque puncto albo. *Syst. farcitella*.  
*Ent.* 660. 26.  
*Phalaena farcitella*. *Linn. Syst. Nat.* 2. 888. 373. *Fn.*  
*Sv.* 1416.  
*Roef. Inf.* 1. phal. 4. tab. 15.  
*Reaum. Inf.* 3. tab. 6. fig. 9. 10.  
Habitat in vestimentis, pellibus, intra cucullum latens.
34. T. alis sublinearibus ferrugineis, macula costali alba. *stigmatel-*  
la.  
Habitat in Anglia. *Mus. Dom. Teats.*

Parua, compressa, linearis. Antennae albicantes. Alae anticae adscendentes, ferrugineae, nitidae macula magna alba in medio costae, quae denticulum marginem anteriorem fere attingentem exserit. Pedes ferruginei apice albi.

*limbella*. 35. T. alis ciliatis exalbidis, limbo maculisque fuscis. *Syst. Ent.* 660. 27.\*  
Habitat in Germaniae nemoribus.

*mellonella*. 36. T. alis canis postice purpurascens, striga alba, scutello nigro, apice candido. *Syst. Ent.* 660. 28.  
*Phalaena mellonella*. Linn. *Syst. Nat.* 2. 888. 375.  
*Fn. Sv.* 1383.

*Swammerd. Bibl.* 1. tab. 26. fig. 2.

*Roef. Inf.* 3. tab. 41.

*Reaum. Inf.* 3. tab. 19. fig. 7 — 9.

Habitat in Apum alucaribus, angiportus fauos penetrantes fruens.

*Larva* nuda, 16poda, grisea capite brunneo.

*Puppa* folliculata, brunnea.

*cucullatella*. 37. T. alis argenteis, arcu dorsali fusco. *Syst. Ent.* 660. 29.  
*Phalaena cucullatella* alis albido cinereis, antice striga nigra recurvata. Linn. *Syst. Nat.* 2. 889. 376. *Fn. Sv.* 1384.  
*Roef. Inf.* 1. phal. 4. tab. 11.

Habitat in Europae Sorbo intra folliculum conicum pilosum.

*Larva* pilosa, 14poda, fusca, dorso albo maculato.

*Puppa* folliculata, obtusa, fusca.

*arcella*. 38. T. alis niueis, arcu communi maculisque duabus marginalibus fuscis. *Gen. Inf. Mant.* 294.\*  
Habitat in Germaniae hortis. *D. Schulz.*

*arbutella*. 39. T. alis rufis, strigis argenteis, intermediis bidis. *Syst. Ent.* 661. 30.

*Phalaena arbutella* alis sanguineis, fasciis argenteis tenuissimis. Linn. *Syst. Nat.* 2. 889. 380. *Fn. Sv.* 1385.  
Habitat in Europae borealis Vna Vrsl.

*tessella*. 40. T. alis nigris, fasciis duabus albis remotissimis tessellatis. *Syst. Ent.* 661. 31.

*Phalaena tessella*. Linn. *Syst. Nat.* 2. 889. 381. *Fn. Sv.* 1430.

Habitat in Europa.

41. T.

41. T. alis anticis niueis, lineola bascos, fascia media *Clematella*.  
punctoque apicis nigris.  
Habitat in Angliae Clematide. *Mus. Dom. Tests.*  
Parua. Caput fuscum. Thorax niueus. Alae niueae  
nitidae lineola marginali bascos nigra. In medio fa-  
scia obliqua angulata nigra et in apice punctum vnum  
alterumque nigrum. Alae posticae cinercae, imma-  
culatae.
42. T. alis cinereis, vitta fusca, palpis porrectis, anten- *bicoßella*.  
nis pilosis. *Syst. Ent.* 661. 32.\*  
*Phalaena bicoßella* alis lanceolatis, cinereo albidis, co-  
sta longitudinali fusca, margine niueo. *Linn. Syst.*  
*Nat.* 2. 890. 387. *Fn. Sv.* 1438.  
*Clerk. phal.* tab. 3. fig. 15.  
*Scop. carn.* 642.  
Habitat in Europae hortis.
43. T. alis anticis testaceis, linea longitudinali alba. *parenthe-*  
*Phalaena parenthesella*. *Linn. Syst. Nat.* 2. 890. 384. *sella*.  
*Fn. Sv.* 1337.  
Habitat in Europa.
44. T. alis linearibus testaceis, antennis mediocribus. *elongella*.  
*Gen. Inf. Mant.* 295.\*  
*Phalaena elongella*. *Linn. Syst. Nat.* 2. 890. 385. *Fn.*  
*Sv.* 1436.  
Habitat in Europa boreali.
45. T. alis anticis linearibus fuscis, maculis tribus flauis. *cylindrel-*  
*Gen. Inf. Mant.* 295.\* *la*.  
Habitat Hamburgi. *D. Schulz.*
46. T. alis anticis linearibus testaceis, punctis 4 fuscis. *punctella*.  
*Syst. Ent.* 661. 34.  
*Phalaena punctella*. *Linn. Syst. Nat.* 2. 890. 386. *Fn.*  
*Sv.* 1337.  
Habitat in Europae pomariis, sedens apicibus alarum  
innititur.
47. T. alis fusco cinereis, lineola media fasciaque posti- *undella*.  
ca undatis albis. *Syst. Ent.* 661. 35.\*  
Habitat in Germaniae nemoribus.

*cinclælla*. 48. T. alis nigris, anticis striga argentea. *Syst. Ent.* 662. 36.  
*Phalaena cinclælla* alis nigris, superioribus fascia lineari argentea transuersa. *Linn. Syst. Nat.* 2. 891. 390.  
*Fn. Sv.* 1380.

*Phalaena verticella*. *Scop. carn.* 651.

Habitat in Europae fruticibus.

Sedens sese circumgyrat saepius circa idem punctum.

*leucatella*. 49. T. alis nigricantibus, fasciis duabus, posteriore oblietioe capiteque albis. *Syst. Ent.* 662. 37.

*Phalaena leucatella*. *Linn. Syst. Nat.* 2. 891. 391. *Fn. Sv.* 1444.

Habitat in Europa.

*strobilella*. 50. T. alis fusco argenteoque vndatis, posticis fuscis, margine albo. *Syst. Ent.* 662. 38.

*Phalaena strobilella*. *Linn. Syst. Nat.* 2. 892. 402. *Fn. Sv.* 1419.

*Phalaena* antennis filiformibus, alis deflexis fuscis, strigis achatinis. *Degeer Inf. Vers. Germ.* 2. 1. 362. 9. tab. 9. fig. 5.

*Clerk. phal.* tab. 12. fig. 10.

Habitat in Abietis strobilis.

*spunclælla*. 51. T. alis fusco cinereis, puncto marginali albo. *Gen. Inf. Mant.* 295. \*

Habitat Hamburgi.

*4spunclælla*. 52. T. alis fuscis, punctis duobus albidis.

Habitat in Germania.

Alae anticae omnino fuscae, nitidae punctis duobus in medio approximatis albis, altero ad marginem tenuiorem maiori transuerso.

*dodecella*. 53. T. grisea nigro punctata, alis pallido fasciatis, punctis trium parium fuscis.

*Phalaena dodecella* alis griseis pallido fasciatis, punctis trium parium fuscis. *Linn. Syst. Nat.* 2. 902. 404. *Fn. Sv.* 1421.

*Phalaena* antennis filiformibus, alis cinereis, fasciis quatuor vndatis fuscis, tuberculisque sex vndatis nigris. *Degeer Inf. Vers. Germ.* 2. 1. 365. 3.

*Degeer Inf.* 1. tab. 22. fig. 22. 23.

Habitat intra Pini Turiones.

*Larva* 16poda, laevis, ferruginea.

*Puppa* folliculata, conica, ferruginea.

*Imago* palpis reflexis corniformibus.

54. *T. alis fusco cinereis, punctis duobus dorsalibus nigro-fuscella.*  
gris. *Syst. Ent.* 662. 39. \*

*Phalaena fuscella.* *Linn. Syst. Nat.* 2. 893. 409. *Fn. Sv.*  
1349.

Habitat in Europae plantis.

55. *T. alis fusco cinereis nitidulis immaculatis.* *Syst. Ent.* cinerella.  
662. 40.

*Phalaena cinerella* alis fusco cinereis immaculatis nitidulis, palpis corniformibus. *Linn. Syst. Nat.* 2.  
891. 392. *Fn. Sv.* 1445.

Habitat in Europae plantis.

56. *T. alis nigris, macula bascos fasciaque media interrupta sulphureis.* *Syst. Ent.* 662. 41.

*Phalaena bractella* alis nigris, macula marginali anticeque luteo aureis. *Linn. Syst. Nat.* 2. 894. 415. *Fn. Sv.* 1426.

*Clerk. phal.* tab. 12. fig. 4.

Habitat in Europae plantis.

Antennae nigrae ante apicem albae.

57. *T. alis fuscis, macula communi dorsali duplici alba trigonella.* *Syst. Ent.* 663. 42.

*Phalaena trigonella.* *Linn. Syst. Nat.* 2. 891. 393. *Fn. Sv.* 1373.

Habitat in Europae plantis.

58. *T. alis fuscis, macula rhombea nigra.* *Syst. Ent.* 663. 43. rhomboidella.  
*Phalaena rhomboidella.* *Linn. Syst. Nat.* 2. 891. 394. *Fn. Sv.* 1372.

Habitat in Europae borealis sylvis.

59. *T. alis nigro, albo argenteoque nebulosis, capite albo.* gemmella.  
*Syst. Ent.* 663. 44.

Pha-



*Phalaena gemmella*. Linn. *Syst. Nat.* 2. 894. 414. *Fa. Sv.* 1388.

*Clerk. phal.* tab. 10. fig. 10.

Habitat intra quercus folia, subcutanea.

*capitella*. 60. T. alis fuscis, maculis tribus flavescentibus, capite ferrugineo. *Syst. Ent.* 663. 45.

*Phalaena capitella*. Linn. *Syst. Nat.* 2. 894. 418. *Fa. Sv.* 1374.

*Phalaena nasuta*, alis aureo fuscis, fasciis argenteis, media interrupta. *Vdm. Diff.* 79.?

Habitat in Angliae hortis, capt. d. VI. Mai.

*sulphurella*. 61. T. fusca, capite, thorace alarumque basi maculaque costali sulphureis. *Gen. Inf. Mant.* 295.\*

Habitat in Germaniae nemoribus. *D. Schulz.*

*compositella*. 62. T. alis fuscis, macula communi dorsali argentea et strigis quatuor composita. *Syst. Ent.* 663. 46.\*

Habitat in Angliae nemoribus.

*minutella*. 63. T. alis atris, maculis duabus ferrugineis suboppositis. *Phalaena minutella* alis nigricantibus maculis transversis ferrugineis. Linn. *Syst. Nat.* 2. 895. 410. *Fa. Sv.* 1440.

*Phalaena nigra*, maculis 4 transversis ferrugineis. *It. Scanic.* 216.

Habitat in Europae hortis.

Maculae duae marginales fere oppositae flavae in medio alae. Caput hirtum, ferrugineum.

*Calthella*. 64. T. atra, alis anticis totis aureis, capite ferrugineo. *Syst. Ent.* 663. 47.

*Phalaena Calthella*. Linn. *Syst. Nat.* 2. 894. 422. *Fa. Sv.* 1432.

Habitat in Calthae palustris floribus frequens.

*Petiverella*. 65. T. alis subgriseis, litura dorsali flava apiceque punctis tribus nigris. *Syst. Ent.* 664. 49.

*Phalaena Petiverella*. Linn. *Syst. Nat.* 2. 895. 423. *Fa. Sv.* 1381.

*Clerk. phal.* tab. 12. fig. 11.

Habitat in Europae floribus.

66. T. alis auratis argenteo subfasciatis, angulo ani macula atra, punctis tribus argenteis. *Fuscella*.

Habitat in Germaniae nemoribus. *Dom. de Hattorf*.

Magnitudo T. Geoffroyellae. Alae anticae obscure aureae, postice strigis quinque argenteis, quarum tres posteriores abbreviatae marginem tenuiorem haud attingunt. Ad angulum ani macula transuersa atra punctis tribus argenteis viuacissimis et ad apicem alia macula atra, immaculata. Posticae cinereae, immaculatae.

67. T. alis flavescentibus argenteo striatis, maculis duabus Geoffroy-sulphureis marginalibus, antennis mediocribus. *Syft. ella*.  
*Ent. 664. 50.*

*Phalaena Geoffroyella. Linn. Syst. Nat. 2. 896. 439.*

Habitat in Europa australi.

68. T. antennis breuibus albo annulatis, alis liuidis, atomis oblongis fuscis. *Syft. Ent. 664. 51. Mouffetella*.

*Phalaena Mouffetella. Linn. Syst. Nat. 2. 896. 431.*

*Fn. Sv. 1397.*

Habitat in Europa boreali.

69. T. alis fuscis, linea vnica aurea antrorsum arcuata. *Albinella. Syft. Ent. 664. 52.*

*Phalaena Albinella. Linn. Syst. Nat. 2. 897. 435. Fn.*

*Sv. 1398.*

Habitat in Europae borealis arboretis.

70. T. alis auratis, strigis duabus argenteis. *Gen. Inf. Seppella. Mant. 296. \**

Habitat in Angliae arboretis. *Dom. Teats.*

71. T. alis cinereis, strigis tribus abbreviatis lineolisque Sulzerella. duabus apicis argenteis fusco marginatis.

Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*

Antennae breues, filiformes, nigrae. Corpus griseo hirtum. Alae cinereae quasi granulatae. Ante medium macula nigra, in qua strigae duae valde abbreviatae argenteae. Pone medium macula alia nigra striga longiori argentea. Ad apicem par macularum fuscarum, in quibus lineola argentea.

**Goedartella.** 72. T. alis auratis, fasciis duabus punctisque tribus argenteis. *Syst. Ent.* 664. 53.

*Phalaena Goedartella* alis auratis fasciis duabus argenteis, priore antrorsum, posteriore retrorsum arcuata. *Linn. Syst. Nat.* 2. 897. 436. *Fn. Sv.* 1401.  
*Clerk. phal.* tab. 12. fig. 14.

Habitat in Alnetis, in Betulae gemmis.

**Merianella.** 73. T. alis nigris, fasciis tribus argenteis transverse diuaticatis. *Syst. Ent.* 664. 54.

*Phalaena Merianella.* *Linn. Syst. Nat.* 2. 897. 438.  
*Fn. Sv.* 1402.

*Tinea* nigro aurata lineis argenteis transversis tribus, antennis extremo albis. *Geoff. Ins.* 2. 198. 43.

*Reaum. Inf.* 1. tab. 17. fig. 12.

Habitat in Europae Pruno, Pado.

**Wilshella.** 74. T. alis nigris, fasciis quatuor argenteis arcuatis, secunda recta. *Linn. Syst. Nat.* 2. 897. 439. *Fn. Sv.* 1403.

*Tinea* nigro aurata, lineis argenteis transversis quatuor, antennis nigris. *Geoff. Inf.* 2. 199. 43. *β.*

*Reaum. Inf.* 3. tab. 4. fig. 8.

Habitat in Europae Vlmo.

**Schieffermüllerella.** 75. T. alis auratis, strigis tribus nigro argenteis, tertia externe bifida.

Habitat in Hamburgensibus arboretis. *D. Schulz.*

Affinis T. Lyonnetellae. Caput et thorax atra, auro marginata. Antennae et pedes nigrae, albo annulatae. Alae auratae nitidae strigis tribus nigro argenteis tertia, ad marginem crassiore late bifida.

**Bergstraeßerella.** 76. T. alis planis auratis, limbo atro punctisque argenteis numerosis.

Habitat in Germaniae Pomariis. *Dom. de Hattorf.*

Antennae atrae. Caput et thorax fusca. Alae auratae nitidae limbo omni atro et in hoc limbo puncta albidiora, in disco aureo magis argentea.

**Lyonnetella.** 77. T. alis flauis, fasciis quatuor argenteis, tertia superne bifida. *Syst. Ent.* 665. 56.

*Pha-*

*Phalaena Lyonetella.* Linn. *Syst. Nat.* 2. 897. 440.

*Fn. Sv.* 1404.

Habitat in Europae nemoribus.

Fascia vltima marginalis.

78. T. alis albis, lineolis duabus argenteis posticeque vn. *Bonnetella.*  
*Syst. Ent.* 665. 57.

*Phalaena Bonnetella.* Linn. *Syst. Nat.* 2. 897. 441.

*Fn. Sv.* 1399.

Habitat in Europae Populetis.

79. T. alis nigris, disco flauo, strigis lineis duabus pun- *Schaeffere-*  
 ctisque tribus argenteis. *Syst. Ent.* 665. 58. *rella.*

*Phalaena Schaefferella.* Linn. *Syst. Nat.* 2. 898. 443.

*Fn. Sv.* 1409.

*Phalaena Fagi.* Linn. *It. Scanic.* 112.

Habitat in Fagi foliis subcutanea.

80. T. alis atris, punctis quatuor argenteis maculisque *Stroemel-*  
 quatuor marginalibus flauis. *It. Norwag.* d. 8. Iul. *la.*

Habitat in Norwagiae nemoribus, capt. d. 8. Iul.

81. T. alis planis atris, fascia media maculisque duabus *Gleichenella.*  
 apicis oppositis argenteis.

Habitat in Angliae nemoribus. *Mus. Dom. Teats.*

Parua. Antennae breues, atrae. Corpus atrum argento  
 nitidulum. Alae atrae nitidae fascia lata in medio,  
 maculisque duabus oppositis versus apicem argenteis.  
 Alae posticae cinerae.

82. T. alis nigris apice fulvis, strigis interruptis argen- *Rhediella.*  
 teis. *Syst. Ent.* 665. 59.

*Phalaena Rhediella.* Linn. *Syst. Nat.* 2. 898. 444.

*Fn. Sv.* 1405.

*Clerk. phal.* tab. 12. fig. 12.

Habitat in Europae borealis nemoribus.

83. T. alis nigro auratis, punctis nouem argenteis conue- *Roesella.*  
 xis submarginalibus. *Syst. Ent.* 665. 60.

*Phalaena Roesella.* Linn. *Syst. Nat.* 2. 898. 445. *Fn.*

*Sv.* 1406.

*Phalaena* antennis filiformibus, alis eleuatis auratis, argenteo maculatis absque puncto nigro. *Degeer Inf. Vers. Germ.* 2. 1. 366. 6.

*Frisch. Inf.* 3. 31. tab. 4.

*Clerk. phal.* tab. 12. fig. 13.

*Degeer Inf.* tab. 30. fig. 10 — 12.

Habitat in Mali foliis subcutanea, nitidissima.

*Larua* 14poda flaua, capite testaceo.

*Puppa* nuda, oblonga, brunnea.

*Linneella*. 84. T. alis fusco auratis, punctis quatuor argenteis eleuatis. *Syst. Ent.* 666. 61.

*Tinea* flauo aurata punctis quatuor argenteis antennarumque apice albo. *Geoff. Inf.* 2. 200. 45.

*Phalaena Linneella* alis fuscis, punctis tribus argenteis eleuatis. *Linn. Syst. Nat.* 2. 898. 446. *Fn. Sv.* 1408.

*Clerk. phal.* tab. 12. fig. 8.

Habitat in Pomonae arboribus.

*Rajella*. 85. T. alis auratis, maculis septem argenteis, secunda tertiaque connatis. *Syst. Ent.* 666. 62.

*Phalaena Rajella*. *Linn. Syst. Nat.* 2. 898. 447. *Fn. Sv.* 1407.

*Phalaena* antennis filiformibus, alis eleuatis auratis, argenteo maculatis punctoque apicis atro. *Degeer Inf. Vers. Germ.* 2. 1. 366. 95.

*Degeer Inf.* 1. tab. 31. fig. 11. 12.

Habitat in Alni foliis subcutanea.

*Larua* pallida, capite vtrinque puncto testaceo.

*Puppa* oblonga mucrone dentato intra folliculum ex-  
tus excrementis tectum.

*Schreberella*. 86. T. alis auratis, basi fasciis duabus maculisque duabus apicis argenteis nitidissimis.

Habitat in Anglia. *Mus. Dom. Teats.*

Parua, at nitidissima. Alae aureae basi fasciis duabus in medio, maculisque duabus oppositis argenteis. Pedes cinereae. Pedes argento nitent.

*Klemanella*. 87. T. alis auratis argento subfasciatis, apice maculis duabus atris, interiori oblonga tripunctata.

Habitat

Habitat in Angliae hortis. *Muf. Dom. Teats.*

Media. Corpus aureum, antennis nigris. Alae aurcae nitidae postice strigis duabus integris tribusque abbreviatis argenteis. Puncta duo atra alam terminant, alterum in medio simplex, alterum ad angulum ani oblongum et in hoc puncta tria argentea.

88. T. alis aureo argenteis, apice obtuso vtitato subocel- *Harrisella*  
lato. *Syst. Ent.* 666. 63.

*Phalaena Harrisella. Linn. Syst. Nat.* 2. 399. 449. *Fn.*  
*Sv.* 1432.

Habitat in Europa, Anglia.

89. T. alis fusco argenteis, fascia postica aurea. *Syst. Ent.* *Drurella*  
666. 64. \*

Habitat Hafniae in Arboretis.

90. T. alis pallide argenteis punctoque terminali atro. *Cramere-*  
*Gen. Inf. Mant.* 297. \*

Habitat in Angliae hortis. *Muf. Dom. Teats.*

91. T. alis auratis, lineola apicis maculisque septem mar- *Blancar-*  
ginalibus argenteis. *della.*

Habitat in Angliae nemoribus. *Muf. Dom. Teats.*

Parua. Antennae supra fuscae, subtus argenteae. Tho-  
rax aureus limbo argenteo. Alae anticae aurcae ni-  
tidae linea longitudinali bascos argentea, quae cum  
limbo thoracis coit. Maculae quatuor ad marginem  
crassior, tres ad tenuiorem argenteae, omnes po-  
ne medium. Alae posticae valde villosae, argenteae,  
immaculatae.

92. T. alis auratis postice atris, fascia argentea. *Syst. Ent.* *aurella*  
666. 65. \*

Habitat Lipsiae in Pomonae arboribus.

Minuta.

93. T. alis argenteis postice subcaudatis, fusco striatis, pun- *Clerkella*  
ctoque atro. *Syst. Ent.* 666. 66. \*

*Phalaena Clerkella* alis argenteis, postice fusco auratis,  
caudatis, ocello terminali atro. *Linn. Syst. Nat.* 2.  
899. 450. *Fn. Sv.* 1411.

Habitat in Europae nemoribus.

*Fabr. Spec. Inf. T. II.*

V

94. T.

- Lichenella*. 94. *T. aptera laevis nigra*.  
*Phalaena Lichenella*, femina aptera laevis nigra. *Linn. Syst. Nat.* 2. 899. 452. *Fn. Sv.* 1451.  
*Phalaena* antennis filiformibus elinguis, femina aptera. *Degeer Inf. Vers. Germ.* 2. 1. 276. 4. tab. 11. fig. 1. 2. 3.  
 Habitat in Europae borealis Lichenibus.  
 Species omnino aptera videtur, mihi haud rite nota, forte ad *Hepialos* amandanda.  
*Larva* 16poda, laevis, obscure grisea, postice incrassata, intra cucullum triangulari prismaticum ex arena constructum.

## 151. ALUCITA.

- Xylostellula*. 1. *A. alis cinereo fuscis, vitta dorsali communi sinuata alba*. *Syst. Ent.* 667. 1.  
*Phalaena Xylostellula*. *Linn. Syst. Nat.* 2. 890. 389. *Fn. Sv.* 1390.  
*Tinea* cinerea dorso vitta longitudinali alba. *Geoff. Inf.* 2. 195.  
*Roef. Inf.* 1. phal. 4. tab. 10.  
 Habitat in Lonicera Xylosteo, Cheiranthi floribus.
- ephippella*. 2. *A. alis pallide auratis, vitta dorsali niuea, fascia aurea*. *Gen. Inf. Mant.* 297. \*  
 Habitat in Europae hortis. *D. Schutz.*
- dentella*. 3. *A. alis fuscis, apice adscendentibus, vitta dorsali communi videntata alba*. *Syst. Ent.* 667. 2. \*  
*Phalaena* antennis filiformibus, alis vitta longitudinali flava. *Degeer Inf. Vers. Germ.* 2. 1. 364. 1.  
 Habitat in Sueciae Caprifolio.  
*Larva* glabra viridis, linea dorsali purpurea.  
*Puppa* flava intra folliculum fusiformem vtrinque apertum.
- styluella*. 4. *A. alis flavo auratis, fasciis duabus fuscis obliquis*. *Syst. Ent.* 667. 3.  
*Phalaena styluella*. *Linn. Syst. Nat.* 2. 893. 413.  
 Habitat in Psaliae nemoribus.

5. A. alis flauis immaculatis, capite thoraceque niueis. *lucella*.  
*Syst. Ent.* 667. 4. \*  
 Habitat in Sueciae arboretis.
6. A. alis fuscis, vitta communi dorsali abbreviata macu- *flauella*.  
 laque flauis. *Syst. Ent.* 667. 5.  
 Habitat in Sueciae arboretis.
7. A. alis cinereis, vitta dorsali margineque postico ni- *vittella*.  
 gris. *Syst. Ent.* 668. 6. \*  
*Phalaena vittella* alis cinereis, vitta dorsali communi  
 nigra, margine postico atro punctato. *Linn. Syst.*  
*Nat.* 2. 390. 382. *Fn. Sv.* 1366.  
*Clerk. phal.* tab. 3. fig. 10.  
 Habitat in Lipsiae hortis.
8. A. alis fusco nitidis, marginibus niueis. *marginel-*  
 Habitat in Iuniperetis Angliae. *Mus. Dom. Teats.* *la*.  
 Media. Palpi crassi, bifidi, interne niuei, externe fu-  
 sci. Caput niueum antennis fuscis. Alae anticae fu-  
 scae, nitidae margine interiori et exteriori late niueo.  
 Posticae exalbidae, immaculatae.
9. A. alis fuscis, vitta communi dentata alba, thorace ni- *bipunctel-*  
 uco, punctis duobus atris. *Syst. Ent.* 668. 7. \* *la*.  
 Habitat in Lichene Prunastri Saxoniae.
10. A. alis albo nigroque variis, capite niueo. *Syst. Ent.* *granella*.  
 668. 8.  
*Phalaena granella*. *Linn. Syst. Nat.* 2. 889. 377. *Fn.*  
*Sv.* 1413.  
*Tinea* tota fusco nebulosa, capite albo. *Groff. Inf.* 2.  
 136. 11.  
*Lewenh. Epist.* 1692. Mart. 7.  
*Reaum. Inf.* 3. tab. 20. fig. 14. 16.  
*Roef. Inf.* 1. phal. 4. tab. 12.  
*Back. microsc.* tab. 13. fig. 13.  
*Degeer Art. Holm.* 1746. 48.  
 Habitat in frumentorum granariis femina rodens glo-  
 meransque, hyeme parietes adscendens.  
*Larua* 16poda, nuda alba, capite brunneo.  
*Puppa* folliculata, attenuata, brunnea.



*niueella*. 11. *A. alis niueis*, maculis duabus marginalibus fasciaeque media nigris, capite albo. *Gen. Inf. Mant.* 298. \*  
Habitat in Anglia. *Muf. Dom. Teats.*

*Lappella*. 12. *A. alis pallidis*, puncto nigro apice adscendentibus. *Syst. Ent.* 668. 9.

*Phalaena Lappella*. *Linn. Syst. Nat.* 2. 889. 378. *Fn. Sv.* 1425.

*Clerk. phal.* tab. 11. fig. 15.

*Aët. Holm.* 1752. 66.

Habitat in Aretii capitulis intra semina.

*costella*. 13. *A. niuea*, alis rufo auratis, macula costali bascos alba fusco punctata. *Syst. Ent.* 668. 10. \*

Habitat in Angliae nemoribus. *Dom. Drury.*

*scabrella*. 14. *A. alis fusco cinereis*, punctis nigris eleuatis scabris. *Gen. Inf. Mant.* 297. \*

*Phalaena scabrella* alis albis, dorso nigro striatis exasperatis, palpis spinosis. *Linn. Syst. Nat.* 2. 891. 396. *Fn. Sv.* 1446.

Habitat in Europa boreali.

*aristella*. 15. *A. albida*, alis linea argentea. *Syst. Ent.* 669. 11.

*Phalaena aristella*. *Linn. Syst. Nat.* 2. 894. 416.

Habitat in Europa boreali.

*caudella*. 16. *A. alis subcaudatis testaceis*, linea postica fusca. *Syst. Ent.* 669. 12.

*Phalaena caudella* alis testaceis, caudali linea fusca, palpis porrectis. *Linn. Syst. Nat.* 2. 894. 417.

Habitat in Europa australiori.

*Swammerdam-* 17. *A. alis pallidis immaculatis*, antennis longissimis. *Syst. Ent.* 669. 13.

*mella*. *Phalaena Swammerdamella*. *Linn. Syst. Nat.* 2. 895. 424. *Fn. Sv.* 1391.

*Clerk. phal.* tab. 12. fig. 1.

Habitat in Europa boreali frequens.

*Robertella*. 18. *A. antennis longissimis albis*, alis fuscis, angulo ant. albo. *Syst. Ent.* 669. 14.

CLASS VI. GLOSSATA. Alucita. 309

- Phalaena Robertella*. Linn. Syst. Nat. 2. 896. 429. Fn. Sv. 1394.  
Habitat in Europae Pomona.
19. A. antennis mediocribus, apice albis, alis fusco auratis. *Frischella*.  
Syst. Ent. 663. 48.  
*Phalaena Frischella*. Linn. Syst. Nat. 2. 896. 433. Fn. Sv. 1396.  
Habitat in Europa boreali frequens.
20. A. alis atro aurcis, fascia flava, antennis longis. *Syst. Degeerella*.  
Ent. 669. 15.  
*Phalaena Degeerella*. Linn. Syst. Nat. 2. 895. 426. Fn. Sv. 1393.  
*Phalaena* antennis filiformibus triplo longioribus, alis deflexis aeneis, fascia flava, coeruleo marginata. *Degeer Inf. Vers. Germ.* 2. 1. 359. 6.  
*Tinea nigra*, alis superioribus lineis longitudinalibus fascia lata transversa inferneque radiis plurimis aurcis, antennis corpore longioribus. *Geoff. Inf.* 2. 193. tab. 12. fig. 5.  
*Degeer Inf.* 1. tab. 32. fig. 13.  
β. *Phalaena Sulzella* antennis mediocribus, alis nigris superioribus fascia aurea. Linn. Syst. Nat. 2. 896. 427.  
Habitat in Europa boreali frequens.  
*Variat* antennis longitudine corporis et triplo longioribus.
21. A. antennis mediocribus, alis nigris, fascia alba. *Syst. Podarella*.  
Ent. 670. 16.  
*Phalaena Podarella*. Linn. Syst. Nat. 2. 896. 428.  
*Tinea nigra*, fascia transversa alba. *Geoffroy Inf.* 2. 194. 32.  
Habitat in Europa boreali.
22. A. alis nigris extrorsum deauratis, antennis longissimis. *Reaumur*.  
Syst. Ent. 670. 17.  
*Phalaena Reaumur*ella. Linn. Syst. Nat. 2. 895. 425. Fn. Sv. 1392.  
*Tinea nigra*. *Geoff. Inf.* 2. 193. 28.  
Habitat in Salice.

310 CLASS. VI. GLOSSATA. Pterophorus.

*fasciella.* 23. A. alis auratis, fascia fusca, antennis apice albis. *Syst. Ent.* 670. 18. \*

Habitat in Angliae nemoribus. *Mus. Dom. Monson.*

Variat antennis longissimis capite thoraceque nigris et antennis mediocribus, capite ferrugineo, thorace aureo.

*friatella.* 24. A. alis auratis, flauo striatis, fascia media flaua eupro marginata.

Habitat in Angliae hortis. *Mus. Dom. Teats.*

Variat capite nigro et fuluo, antennis longissimis albidis et antennis corpore paullo longioribus, basi hirtis atris, apice albis. Alae aureae inprimis apice flauo striatae. In medio cupreae nitidissimae fascia flaua, Alae posticae fusco aureae, nitidae.

*sulphurella.* 25. A. alis anticis auratis, maculis duabus sulphureis oppositis, posticis flauis. *Syst. Ent.* 670. 19. \*

Habitat in Angliae hortis. *Mus. Dom. Monson.*

*oppositella.* 26. A. alis fuscis, maculis duabus oppositis flauis, posticis fuscis. *Syst. Ent.* 670. 20. \*

Habitat in Angliae nemoribus.

Varietatem fascia antica flaua misit. *Dom. Lund.*

152. PTEROPHORVS.

*monodactyla.* 1. P. alis patentibus linearibus indiuisis. *Syst. Ent.* 671. 1. *Phalaena monodactyla.* *Linn. Syst. Nat.* 2. 899. 453. *Fn. Sv.* 1452.

*Reaum. Inf.* 1. tab. 20. fig. 12 — 16.

Habitat in Europae hortis.

*didactylus.* 2. P. alis fisis fuscis, strigis albis, anticis bifidis, posticis tripartitis. *Syst. Ent.* 671. 2.

*Pterophorus* fuscus, alis superioribus apice bipartitis, inferioribus tripartitis. *Geoff. Inf.* 2. 92. 2.

*Phalaena didactyla.* *Linn. Syst. Nat.* 2. 899. 454. *Fn. Sv.* 1453.

*Phalaena* fusca albofstriata, alis anticis didactylis, posticis tridactylis. *Degeer Inf. Vers. Germ.* 2. 1. 136. 1. tab. 4. fig. 9.

*Raf.*

CLASS. VI. GLOSSATA. Pterophorus. 311

*Raj. Inf.* 205. 101.

*Petiv. Gazoph.* tab. 66. fig. 10.

*Merian. Europ.* tab. 22.

*Schaeff. Elem.* tab. 104.

— — *Icon.* tab. 93. fig. 7.

Habitat in Conuoluulo; Géo riuali.

*Larua* 16poda, viridis, fasciculato pilosa.

*Puppa* elongata lineis duabus dorsalibus tuberculorum quadrispinosorum.

3. *P.* alis fissis pallidis, lineis albis, anticis bifidis, posticis tridactylus. *Syst. Ent.* 671. 3.

*Phalaena tridactyla.* *Linn. Syst. Nat.* 2. 899. 455.

*Acta Soc. Berol. Nat. cur.* 3. tab. 1. fig. 8.

Habitat in Europae hortis.

4. *P.* alis patentibus fissis, cinereo nebulosis, posticis fustigeradactylus. *Syst. Ent.* 671. 4.

*Phalaena tesseradactyla.* *Linn. Syst. Nat.* 2. 900. 456.

*Fn. Sv.* 1454.

Habitat in Europae pratis.

5. *P.* alis patentibus fissis testaceis, puncto fusco. *Syst. Ent.* 671. 5. *pterodactylus.*

*Phalaena pterodactyla.* *Linn. Syst. Nat.* 2. 900. 458.

*Fn. Sv.* 1456.

*Scop. carn.* 674.

*Wien. Verz.* 297. tab. 1. a. 3.

— — — tab. 1. b. 8.

Habitat in Europae Conuoluulo.

*Larua* subpilosa coerulea, linea dorsali purpurea.

6. *P.* alis niueis, anticis bifidis, posticis tripartitis. *Syst. Ent.* 672. 6. *pentadactylus.*

*Phalaena pentadactyla* alis patentibus fissis quinque-

partitis niueis, digito quinto distincto. *Linn. Syst.*

*Nat.* 2. 900. 459. *Fn. Sv.* 1257.

*Roef. Inf.* 1. phal. 4. tab. 5.

*Reaum. Inf.* 1. tab. 20. fig. 1. 2.

*Petiv. Gazoph.* tab. 67. fig. 6.

*Ammir. Inf.* tab. 23.

*Sulz. Inf.* tab. 16. fig. 10.

— — —

312 CLASS. VI. GLOSSATA. Pterophorus.

Habitat in Europa frequens.

*Larva* 16poda, pilosa, viridis nigro punctata lineaque dorsali alba.

*Puppa* pilosa, viridis, nigro punctata.

*hexada-* 7. *P. alis fissis cinereis, singulis sexpartitis. Syst. Ent.*  
*stylus.* 672. 7.

*Pterophorus cinereus fusco maculatus, alis superiori-*  
*bus octopartitis, inferioribus tripartitis. Geoff. Inf.*  
2. 92. 3.

*Phalaena hexadactyla. Linn. Syst. Nat. 2. 900. 460.*  
*Fn. Sv. 1458.*

*Reaum. Inf. 1. tab. 19. fig. 19 — 21.*

*Harris Inf. angl. tab. 2. fig. 7.*

*Frisch. Inf. 7. tab. 73.*

*Petiv. Gazoph. tab. 67. fig. 7.*

Habitat in Lonicera Xylosteo.





CLASSIS VII.  
R Y N G O T A.

153. FVLGORA.

1. **F** fronte rostrata recta, alis lividis, posticis ocella- *laterna-*  
tis. *Syst. Ent.* 673. 1. *Linn. Syst. Nat.* 2. 703. 1. *ria.*  
*Laternaria phosphorea.* *Linn. Mus. Lud. Vlr.* 152.  
*Cucuius peruianus.* *Grew. Mus.* 158. tab. 13.  
*Laternaria.* *Merian. Sur.* tab. 49.  
*Reaum. Inf.* 5. tab. 20. fig. 6. 7.  
*Deger Inf.* 3. 195. 1.  
*Roef. Inf.* 2. *Gryll.* tab. 28. 29.  
*Seb. Mus.* 4. tab. 77. fig. 3. 4.  
Habitat in America meridionali.  
Prominente fronte noctu lucem viuacissimam spargit.
2. **F.** fronte rostrata adscendente, quadrifarie serrata. *serrata.*  
*Naturf.* 13. tab. 3. fig. 1. 2.  
*Seb. Mus.* 4. tab. 77. fig. 5. 6.  
Habitat Surinami.  
Frons rostrata adscendens, dentibus validis acutis qua-  
drifarie serrata. Elytra fusca. Alae apice ocellatæ.
3. **F.** fronte rostrata muricata, apice trifido, alis nigris *Diade-*  
rubro maculatis. *Syst. Ent.* 673. 2. *Linn. Syst. Nat.* *ma.*  
2. 703. 2.  
*Naturf.* 13. tab. 3. fig. 3.  
*Seb. Mus.* 4. tab. 77. fig. 7. 8.  
Habitat in India.
4. **F.** fronte rostrata adscendente, elytris viridibus luteo *candela-*  
maculatis, alis flavis apice nigris. *Syst. Ent.* 673. 3. *ria.*  
*Linn. Syst. Nat.* 2. 703. 3.  
*Laternaria candelaria.* *Mus. Lud. Vlr.* 153.  
*Aff. Holm.* 1746. 63. tab. 1. fig. 5. 6.

*Edw. Av.* tab. 120.

*Degeer Inf.* 3. 197. 2.

*Roef. Inf.* 2. *Gryll.* 189. tab. 30.

*Sulz. Inf.* tab. 10. fig. 62.

Habitat in China.

- phosphorea.* 5. *F. fronte rostrata subulata adscendente, corpore griseo glauco. Syst. Ent.* 674. 4. *Linn. Syst. Nat.* 2. 704. 4.

*Cicada filirostris* griseo fusca, subtus flavescent, fronte rostrata filiformi adscendente. *Degeer Inf.* 3. 201. 4. tab. 32. fig. 6.

Habitat in America meridionali.

- noctuida.* 6. *F. fronte rostrata acuminata adscendente, corpore viridi, alis hyalinis. Syst. Ent.* 674. 5. *Linn. Syst. Nat.* 2. 704. 5.

*Cicada conirostris* viridis, fronte rostrata conico adscendente, alis hyalinis. *Degeer Inf.* 3. 202. 5. tab. 32. fig. 4.

Habitat in America meridionali.

- lucernaria.* 7. *F. fronte rostrata prominente, corpore supra viridescente, subtus flauo. Syst. Ent.* 674. 6. *Linn. Syst. Nat.* 2. 704. 6.

*Cicada breuirostris* supra viridis, subtus flaua, fronte rostrata obtusa, breui. *Degeer Inf.* 3. 202. 6. tab. 32. fig. 6.

Habitat in America meridionali.

- flammea.* 8. *F. fronte rostrata adscendente tereti truncata. Syst. Ent.* 674. 7. *Linn. Syst. Nat.* 2. 704. 7. *Amoen. acad.* 6. 399. 39.

Habitat in America meridionali.

- tenebrosa.* 9. *F. fronte rostrata recta truncata, elytris griseis scabris. Syst. Ent.* 674. 8. \*

*Cicada laternaria fusca* fronte rostrata subulata recta, apice truncata, alis superioribus punctis nigris. *Degeer Inf.* 3. 200. 3. tab. 32. fig. 1.

Habitat in Guinea. *Mus. Dom. Banks.*

CLASS. VI. RYNGOTA. Fulgora. 315

10. F. fronte rostrata recta truncata, elytris cinereis, nigro maculatis. *Syst. Ent.* 674. 9.\* *obscurata.*  
Habitat in noua Hollandia.
11. F. fronte obtusiuscula, alis truncatis viridibus. *Syst. truncata.*  
*Ent.* 674. 10. *Linn. Syst. Nat.* 2. 704. 8.  
*Cicada truncata* viridi albicans, alis deflexis compressis, postice truncatis. *Linn. Amoen. acad.* 6. 399. 40.  
Habitat in India.
12. F. fronte conica inaequali, elytris hyalinis, striga atra. *hyalina-*  
Habitat in Bengal. *Mus. Dom. Banks.* *ta.*  
Maior F. obscurata. Frons rostrata, breuis, conica, supra inaequalis, subtus fulcata, pallida atomis strisque nigris. Oculi magni, globosi, albi. Thorax pallidus nigro irroratus. Scutellum magis obscurum. Elytra hyalina nervis albo nigroque punctatis strigae in medio atra. Alae hyalinae macula apicis atra.
13. F. fronte conica, elytris fuscis, margine exteriori virescente, punctis nigris fulvisque, alis basi rubris. *festiva.*  
Habitat in Coromandel. *Mus. Dom. Banks.*  
Caput supra planum, fuscum, subtus flauescens margine atro. Thorax fuscus, immaculatus. Elytra fusca, margine exteriori virescente, qui color tamen apicem alae haud attingit. In hoc margine puncta quinque atra, quorum quatuor posteriora intus puncto fuluo terminantur. Alae sanguineae, apice fuscae.
14. F. fronte conica bifulcata, elytris hyalinis, macula fuscis, *fenestrata.*  
marginali fusca.  
Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*  
Minor F. europaea. Frons conica, flauescens, immaculata dorso fulcis duobus exarata. Elytra hyalina nervis nigris maculaque magna marginali fusca. Alae hyalinae, immaculatae.
15. F. fronte conica, corpore viridi, alis hyalinis reticulatis. *Syst. Ent.* 674. 11. *Linn. Syst. Nat.* 2. 704. 9. *europaea.*  
*Naturforsch.* 9. tab. 3.  
*Sulz. Hist. Inf.* tab. 9. fig. 5.  
Habitat in Europa australiori.



316 CLASS. VII. RYNGOTA. Membracis.

154. MEMBRACIS.

*rhombea*. 1. M. foliacea, folio rhombeo postice latiore. *Syst. Ent.* 675. 1.\*

*Cicada rhombea*. Linn. *Syst. Nat.* 2. 704. 1.

*Aët. Angl.* 1765. 55. tab. 6.

Habitat in America.

*fronditia*. 2. M. thorace foliaceo, dorso repando sinuato, apice porrecto obtusissimo.

*Cicada fronditia* foliacea, dorso repando, rostro obtusissimo. Linn. *Syst. Nat.* 2. 705. 3.

*Cicada foliata sinuata* fusca, thorace compresso foliaceo, margine sinuoso, apice rostrato. *Degeer Inf.* 3. 208. 11. tab. 32. fig. 15. 16.

Habitat Surinami.

*foliata*. 3. M. thorace foliaceo rotundato atro, arcubus albis. *Syst. Ent.* 675. 2.\*

*Cicada foliata*. Linn. *Syst. Nat.* 2. 705. 2.

*Cicada foliata fasciata* nigro fusca, thorace compresso foliaceo verticali, fascia arcuque albis. *Degeer Inf.* 3. 205. 8. tab. 32. fig. 9. 10.

*Gronov. Zooph.* 677.

*Merian. Surin.* tab. 5. fig. vlt.

Habitat Surinami.

*squamigera*. 4. M. thorace foliaceo antice posticeque acuto griseo. *Syst. Ent.* 675. 3.\*

*Cicada squamigera*. Linn. *Syst. Nat.* 2. 705. 4.

*Cicada hastata* griseo livida, thorace magno elevato, antice posticeque acuto, pedibus quatuor primoribus latis compressis. *Degeer Inf.* 3. 209. 12. tab. 32. fig. 17. 18.

Habitat in America meridionali.

*spinosa*. 5. M. thorace tricorni, postice producto longitudine alarum. *Syst. Ent.* 675. 4.\*

*Sulz. Hist. Inf.* tab. 9. fig. 6.

Habitat in Indiis.

**CLASS VII. RYNGOTA, Membracis. 317**

6. M. thorace tricorni, cornu intermedio longiori com- *acumina-*  
presso. *Syst. Ent. 675. 5.\** *ta.*  
Habitat in Pennsylvania. *Mus. Dom. Banks.*
  
7. M. thorace biaurito, capitis clypeo antrorsum dilatato, *aurita.*  
rotundato. *Syst. Ent. 676. 6.*  
*Cicada aurita. Linn. Syst. Nat. 2. 706. 7.*  
*Cicada* thorace obtuse bicorni. *Geoff. Inf. 422. 17.*  
*Cicada* cristata, thorace bicorni, corniculis subrotun-  
dis. *Schreb. Inf. 8. fig. 1. 2.*  
*Schaeff. Icon. tab. 96. fig. 3.*  
*Sulz. Hist. Inf. tab. 9. fig. 7.*  
Habitat in Quercu, Corylo.
  
8. M. thorace bicorni albo punctato postice producto ab- *punctata.*  
domine longiori. *Syst. Ent. 676. 7.\**  
Habitat in Brasilia. *Mus. Dom. Banks.*
  
9. M. thorace bicorni nigro, postice subulato longitudine *cornuta.*  
abdominis, alis fuscis. *Syst. Ent. 676. 8.*  
*Cicada cornuta. Linn. Syst. Nat. 2. 705. Fn.*  
*Sv. 879.*  
*Cicada* thorace acute bicorni, pone producto. *Geoff.*  
*Inf. 2. 423. 18.*  
*Cicada cornuta. Schreb. Inf. 11. fig. 3. 4.*  
*Cicada* fusca, thorace bicorni postice subulato longitu-  
dine abdominis. *Degeer Inf. 3. 181. 9. tab. 11.*  
*fig. 22.*  
*Ranata cornuta. Petiv. Gazoph. tab. 47. fig. 2. 3.*  
*Sulz. Inf. tab. 10. fig. 63.*  
*Schaeff. Icon. tab. 96. fig. 2.*  
*Scop. carn. 340.*  
Habitat in Carduis, Salicibus.
  
10. M. thorace bicorni fusco, postice producto filiformi *Taurus.*  
longitudine abdominis, cornubus arcuatis. *Syst.*  
*Ent. 676. 9.\**  
Habitat in Coromandel. *D. Koenig.*
  
11. M. thorace bicorni viridi, postice subulato longitudine *Vitalus.*  
abdominis, alis albis. *Syst. Ent. 677. 10.\**  
Habitat in America. *Dom. Drury.*

318 CLASS. VII. RYNGOTA. Tettigonia.

- spunctata*. 12. M. thorace subipermi postice producto, abdomine brevius testaceo, elytris macula bascos nigra. *Syst. Ent.* 677. 11.\*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

- Bonasia*. 13. M. thorace bicorni postice producto, margine albo, alis puncto bascos albo. *Syst. Ent.* 677. 12.\*

Habitat in America. *Mus. D. Hunter.*

- conuoluta*. 14. M. thorace inermi laeui virescente, margine flavo, postice producto abdomine longiori.

Habitat in Brasilia. *Mus. Dom. Banks.*

Media. Caput flavum, laeue immaculatum. Thorax laevis, valde productus alas abdomenque involvens, viridescens margine anteriori et laterali flavescentis. Apex acutus, ater.

- mutica*. 15. M. thorace inermi ferrugineo, postice producto carinato longitudine abdominis. *Gen. Inf. Mant.* 297.\*

Habitat in America boreali.

- inermis*. 16. M. thorace inermi virescente, postice subulato longitudine abdominis. *Syst. Ent.* 677. 13.\*

Habitat in Americae plantis.

- Genistae*. 17. M. thorace inermi fusco, postice producto, abdomine dimidio brevius. *Syst. Ent.* 677. 14.\*

*Cicada* thorace inermi pone producto. *Geoff. Inf.* 1. 424. 19.

Habitat in Genista tinctoria Angliae. *Dom. Lee.*

155. TETTIGONIA.

- grossa*. 1. T. thorace viridi nigro sublineata, alis albis, postice macula bascos flava. *Syst. Ent.* 678. 1.\*

Habitat in Brasilia. *Mus. Dom. Banks.*

- grisea*. 2. T. grisea, elytris aqueis, margine postico nigro punctato, costa antice alba. *Syst. Ent.* 678. 2.\*

Habitat in Americae insulis. *Mus. Tottianum.*

2. T. thoracis margine dilatato acuto, alis atris, margi- *limbata*,  
ne albo. *Syst. Ent.* 678. 3.\*  
Habitat in America. *Mus. Dom. Banks.*
4. T. thoracis margine dilatato, atra alis albidis. *Syst. dilatata*.  
*Ent.* 678. 4.\*  
Habitat in Iamaica. *Mus. Dom. Banks.*
5. T. scutello emarginato, alarum costa virescente. *Syst. Tibicen*.  
*Ent.* 679. 5.\*  
*Cicada Tibicen* elytris anastomosibus ferrugineis, scu-  
telli apice emarginato. *Linn. Syst. Nat.* 2. 707. 19.  
*Mus. Lud. Vlr.* 160.  
*Cicada Lyricus* thorace antice viridi, postice fuluo ma-  
culis nigris, alis hyalinis, venis fuscis, margine di-  
latato viridi. *Degeer Inf.* 3. 212. 14. tab. 22. fig. 23.  
*Tibicen. Merian. Sur.* tab. 49.  
*Panorpa. Brown. Iam.* 436. tab. 43. fig. 15.  
*Guerucu Erementi. Margr. Bras.* 256.  
*Seb. Mus.* 4. tab. 85. fig. 9. 10.  
Habitat in America.  
Variat thorace nigro et nigro brunneoque variegato.
6. T. nigra, elytris albis, costa flava. *Syst. Ent.* 679. 6.\* *septende-*  
*Cicada septendecim* nigro virescens elytris margine fla- *decim*.  
uescente, capite vtrinque octofriato. *Linn. Syst.*  
*Nat.* 2. 708. 20.  
*Cicada* maxilla vtraque lincis octo transuersis concavis,  
alarum margine inferiore lutescente. *Kalm. Aët.*  
*Holm.* 1756. 101.  
*Collins. Aët. angl.* 1765. 65. tab. 8.  
Habitat in America.
7. T. thorace variegato, elytris hyalinis antice nervis *catena*.  
punctatis, postice strigis duabus vndatis fuscis. *Syst.*  
*Ent.* 679. 7.\*  
Habitat in Capite Bon. Spei. *Mus. Dom. Banks.*
8. T. atra, thorace, elytris alisque flauo maculatis. *Syst. maculata*.  
*Ent. Append.* 831.\*  
*Cicada maculata. Drury Inf.* 2. tab. 37. fig. 1.  
Habitat in China.

320 CLASS. VII. RYNGOTA. Tettigonia.

- fridula*. 9. *T. villosa*, elytris griseis, alis flavis, omnibus margine hyalino. *Syst. Ent.* 679. 8.\*  
*Cicada fridula* elytris margine dilatato, punctis septem albis. *Linn. Syst. Nat.* 2. 706. 12. *Maj. Lud. Vir.* 157.  
*Cicada catenata*. *Drury Inf.* 2. tab. 37. fig. 2.  
Habitat in Capite Bon. Spei.
- cingulata*. 10. *T. thorace maculato*, elytris hyalinis, costa maculataque fuscis, abdomine atro, fasciis flavis. *Syst. Ent.* 680. 9.\*  
Habitat in noua Zelandia. *Mus. Dom. Banks.*
- villosa*. 11. *T. thorace fusco viridique variegato*, pectore albo villoso.  
Habitat ad promontorium Bon. Spei. *Mus. Dom. Banks.*  
Minor, frons viridis vtrinque strigis atris. Thorax viridis fusco varius. Pectus lana alba tectum. Scutellum nigrum lineis quatuor flavis. Abdomen viride. Alae hyalinae nervis nigris puncto vno alteroue albo.
- cruentata*. 12. *T. thorace variegato*, elytris hyalinis, costa fluorescente, abdomine atro, fasciis sanguineis. *Syst. Ent.* 680. 10.\*  
Habitat in noua Hollandia. *Mus. Dom. Banks.*
- conspurcata*. 13. *T. atra*, elytris basi flauo maculatis, ano flauo. *Gen. Inf. Mant.* 298.\*  
Habitat in India. *D. Fothergill.*
- haematodes*. 14. *T. nigra immaculata*, abdominis incisuris sanguineis. *Syst. Ent.* 680. 11.\*  
*Cicada haematodes*. *Linn. Syst. Nat.* 2. 707. 14.  
*Scop. carn.* 347.  
*Petiv. Gazoph.* tab. 15. fig. 7.  
*Gron. Zooph.* 675.  
*Fuesl. Inf.* 24. 452.  
Habitat in Europa australi.
- Orni*. 15. *T. nigra flauo maculata*, elis albis, basi flavis, maculis duabus nigris. *Syst. Ent.* 680. 12.\*

CLASS. VII. RYNGOTA. Tettigonia. 321

*Cicada orni* clytris intra marginem punctis sex concatenatis anastomosibusque interioribus fuscis. *Linn. Syst. Nat.* 2. 707. 16.

*Scop. carn.* 346.

*Seb. Mus.* 4. tab. 85. fig. 4. 5.

*Roef. Inf.* 2. Gryll. tab. 25.

*Reaum. Inf.* 5. tab. 5. fig. 16. 17. 18. 19.

*Sulz. Hist. Inf.* tab. 10. fig. 65.

*Schaeff. Icon.* tab. 4. fig. 14.

Habitat in Europa australi.

16. *T. clytris linea flexuosa, alis margine hyalino.* *repanda.*

*Cicada repanda.* *Linn. Syst. Nat.* 2. 707. 17.

*Cicada nigra* linea fulva, thoracē nigra linea, alis linea flexuosa fusca, maculis margineque hyalinis.

*Degeer Inf.* 3. 219. 15. tab. 33. fig. 1.

Habitat in India.

17. *T. grisea, thoracis linea alba, clytris albo reticulatis.* *Syst. Ent.* 681. 13. *reticulata.*

*Cicada reticulata.* *Linn. Syst. Nat.* 2. 707. 18.

*Cicada fusco testacea, capite obtuso, thorace linea alba, alis albo reticulatis, tibiis posticis nigro maculatis.* *Degeer Inf.* 3. 227. 24. tab. 23. fig. 15. 16.

Habitat in America meridionali.

18. *T. atra, alis albis basi nigris, venis testaceis.* *Syst. atrata.*  
*Ent.* 681. 14.\*

Habitat in China. *Dom. Gunnerus.*

19. *T. nigra, ore thoracis maculis duabus abdomineque sanguineis.* *Syst. Ent.* 681. 15.\* *sanguinolenta.*

*Cicada sanguinea* alis superioribus fuscis, fronte abdomine thoracisque maculis binis sanguineis. *Degeer Inf.* 3. 221. 18. tab. 33. fig. 17.

Habitat in China. *Dom. Drury.*

20. *T. clytris fusco aureis, tibiis anticis incrassato densitatis rufis.* *Syst. Ent.* 681. 16.\* *splendidiola.*

Habitat in China. *Dom. Drury.*

*Fabr. Sp. Inf. T. II.*

X

21. *T.*

322 CLASS. VII. RYNGOTA. Cicada.

*muta*. 21. T. elytris hyalinis, costa sanguinea, abdomine linea dorsali sanguinea. *Syst. Ent.* 681. 17.\*

Habitat in noua Zelandia. *Mus. Dom. Banks.*

*violacea*. 22. T. violacea, alis apice fuscis. *Syst. Ent.* 682. 18.

*Cicada violacea*. Linn. *Syst. Nat.* 2. 708. 21. *Mus.*

*Lud. Vlr.* 162.

*Fuesly Inf. Helv.* 24. 455.

Habitat in Europa australiori.

156. C I C A D A.

*perspicillata*. 1. C. atra, alis macula ocellari albo hyalina, abdomine apice flauo.

Habitat in India. *Mus. D. Schulz.*

Gigas in hoc genere. Caput et thorax atra oculis antennisque fuluis. Abdomen atrum vltimis segmentis flauis striga atra. Elytra atra, immaculata. Alae atrae macula magna ocellari alba hyalina versus marginem posticum. Pedes nigri, posticis geniculis flauis.

*ocellata*. 2. C. elytris compressis adscendentibus viridibus, punctis ocellaribus ferrugineis. *Syst. Ent.* 682. 1.

*Degeer Inf.* 3. tab. 33. fig. 2.

Habitat Tranquebariae. *D. Koenig.*

*limbata*. 3. C. elytris deflexis viridibus, margine rubro, interiori basi nigro punctato.

Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*

Magna. Caput ferrugineum, vtrinque inter oculos eleuatum, antennis breuibus nigris. Thorax glaber, lacus lobo antico virescente. Elytra lata, deflexa, virescentia, margine rubro, qui postice lineola tenuissima nigra terminatur. Sutura antica nigro punctata puncto distincto vtrinque ad scutellum, postice rubra. Alae albae immaculatae.

Aliam simillimam vidi, at totam exalbidam sutura basi nigro punctata margineque elytrorum postico nigro.

*marginata*. 4. C. flaua, elytris deflexis nigris, linea laterali alba. *Syst. Ent.* 682. 2. Linn. *Syst. Nat.* 2. 710. 39.

Habi-

CLASS. VII. RYNGOTA. Cicada. 323

Habitat in America.

Alae pallidae.

5. C. exalbida, alis deflexis patulis, elytris antice fusco *phala-*  
punctatis. *Syst. Ent.* 682. 3. *Linn. Syst. Nat.* 2. *noides.*  
710. 40.

*Cicada flava*, alis amplissimis deflexis albidis nigro pun-  
ctatis. *Degeer Inf.* 3. 222. 19. tab. 33. fig. 6.

*Sulz. Hist. Inf.* tab. 9. fig. 10.

Habitat in America meridionali.

6. C. exalbida, elytris deflexis, thoracis margine antico *bipuncta-*  
punctis duobus compressis. *Syst. Ent.* 682. 4. *Linn.* *ta.*  
*Syst. Nat.* 2. 710. 41.

Habitat in America meridionali.

7. C. grisea, elytris deflexis, capite apice puncto nigro. *grisea.*  
*Syst. Ent.* 683. 5.\*

Habitat in Americae meridionalis insulis. *Dom. Rohr.*

8. C. elytris deflexis viridibus, punctis tribus albidis. *3puncta-*  
Habitat in Africa aequinoctiali. *Muf. Dom. Banks.* *ta.*  
Maior C. viuida. Corpus totum viride fronte fuleata.  
Elytra magna, deflexa, viridia punctis tribus albis  
transuersim positis. Margo posticus sanguineus.  
Alae albae.

9. C. viridis, elytris deflexis, alis albis, fronte conica. *viuida.*  
*Syst. Ent.* 683. 6.\*

Habitat in Americae meridionalis insulis, *Dom. Rohr,*  
in noua Hollandia. *Muf. Dom. Banks.*

10. C. elytris deflexis viridibus, dorso scabro, capitis *minuta.*  
margine postico thoraceque linea dorsali sanguinea.  
*Syst. Ent.* 683. 7.\*

Habitat in noua Hollandia. *Muf. Dom. Banks.*

11. C. alis deflexis hyalinis, friga, fascia punctisque fuscis, *cunicula-*  
ano lanato. *Syst. Ent.* 683. 8. *Linn. Syst. Nat.* 2. *ria.*  
711. 44.

Habitat in India orientali.  
Frons ouata ferruginea.



324 CLASS. VII. RYNGOTA. Cicada.

- lanata*. 12. C. alis nigris, punctis coeruleis, fronte lateribusque rubris, ano lanato. *Syst. Ent.* 683. 9. *Linn. Syst. Nat.* 2. 711. 42. *Mus. Lud. Vir.* 163.  
*Cicada lanata*. *Drury Inf.* 2. tab. 37. fig. 11.  
*Sulz. Hist. Inf.* tab. 9. fig. 11.  
 Habitat in India.  
 Oculi spinosi.
- tomento-  
sa*. 13. C. elytris deflexis viridibus, fuluo subfasciatis, alis albis, fasciis duabus atris. *Syst. Ent.* 683. 10.\*  
 Habitat in India orientali. *Mus. Tottianum*.
- barbata*. 14. C. fusca abdomine virescente, ano lanato niueo. *Syst. Ent.* 684. 11.\*  
 Habitat in noua Hollandia. *Mus. Dom. Banks*.
- villosa*. 15. C. viridis, elytris deflexis exalbidis, ano lanato. *Syst. Ent.* 684. 12.\*  
 Habitat in America meridionali. *Dom. Rohr*.
- elongata*. 16. C. thorace rubro, flauo lineato, elytris longissimis nigro punctatis.  
 Habitat in noua Hollandia. *Mus. Dom. Banks*.  
 Corpus paruum, gibbum. Frons inter oculos eleuata, canaliculata, flaua. Thorax lacte ruber margine antico lineisque flauis. Abdomen flauum, fusco varium. Elytra corpore triplo longiora, obscura, nigro punctata. Alae obscurae.
- vittata*. 17. C. flaua, vitta ferruginea longitudinali duplici repando dentata. *Syst. Ent.* 684. 13. *Linn. Syst. Nat.* 2. 710. 36. *Fn. Sv.* 889.  
*Fuesl. Inf. Helv.* 24. 463.  
 Habitat in Europae hortis.
- lateralis*. 18. C. nigra, elytris latere albis. *Syst. Ent.* 684. 14. *Linn. Syst. Nat.* 2. 709. 29. *Fn. Sv.* 886.  
*Locusta pulex* fusca vtrinque in lateribus alarum arcu albicante. *Raj. Inf.* 68. 2.  
*Geoff. Inf.* 1. 416. 3.  
*Fuesl. Inf. Helu.* 24. 460.  
 Habitat in Europae plantis.

CLASS. VII. RYNGOTA. Cicada. 325

19. *C. supra nigra*, scutello elytrorum maculis duabus dor- *variega-*  
salibus margineque exteriori flavescentibus. *Syst. ta.*  
*Ent. 684. 15. \**  
Habitat in Brasilia. *Mus. Dom. Banks.*
20. *C. supra nigra*, capite scutelloque flavis, elytrorum *fenestra-*  
margine hyalino. *Syst. Ent. 684. 16. \** *ta.*  
Habitat in Insula Otaheiti oceani pacifici. *Mus. Dom.*  
*Banks.*
21. *C. nigra*, oculis albis. *Syst. Ent. 685. 17. Linn. Syst. leuco-*  
*Nat. 2. 709. 26. Fn. Sv. 883.* *phthalma.*  
Habitat in Europae borealis plantis.
22. *C. nigra flavo maculata*, elytris fuscis, puncto margi- *bigutta-*  
nali albo. *ta.*  
Habitat in Germania. *Dom. de Hattorf.*  
Statura parva praecedentis. Caput nigrum punctis ali-  
quot flavis. Thorax niger margine antico flavo.  
Elytra laevia fusca puncto albo versus posteriora.  
Abdomen atrum ano rufo.
23. *C. capite thoracisque basi flavescentibus. Syst. Ent. leucoco-*  
*685. 18. Linn. Syst. Nat. 2. 709. 28. Fn. Sv. 885. phala.*  
*Geoff. Inf. 1. 421. 13.*  
Habitat in Europae plantis.
24. *C. elytris flavis*, linea nigra duplici interrupta longi- *inter-*  
tudinali. *Syst. Ent. 685. 19. Linn. Syst. Nat. 2. 710. rupta.*  
*35. Fn. Sv. 889.*  
*Cicada* alis superioribus flavis, fasciis binis nigris in-  
interruptis longitudinalibus, pedibus flavis. *Degeer Inf.*  
*3. 186. 7. tab. 12. fig. 6.*  
*Ranatra* bicolor ex fusco et pallido striata. *Petiv. Ga-*  
*zoph. tab. 61. fig. 10.*  
*Geoff. Inf. 1. 419. 9.*  
*Fuesly Inf. Helv. 24. 462.*  
Habitat in Europae plantis frequens.
25. *C. nigra*, capitis margine postico thoraceque flavis. *flavicollis.*  
*Syst. Ent. 685. 20. \**

326 CLASS. VII. RYNGOTA. Cicada.

- Cicada flavicollis* fusca, capite thoraceque flavis. *Lin.*  
*Syst. Nat.* 2. 709. 33. *Fn. Sv.* 891.  
 Habitat in Europae graminibus.
- viridis.* 26. *C. elytris viridibus, capite flavo, punctis nigris. Syst.*  
*Ent.* 2. 685. 21. *Lin.* *Syst. Nat.* 2. 711. 46. *Fn. Sv.*  
 896.  
*Locusta pulex* paullo minor. *Raj. Inf.* 68. 3.  
*Ranatra viridescens. Petiv. Gazoph.* 73. tab. 76. fig. 6.  
*Geoff. Inf.* 1. 417. 5.  
*Fuesly Inf. Helv.* 24. 465.  
 Habitat in Europae plantis.
- Lanio.* 27. *C. viridis, capite thoraceque carneis. Syst. Ent.* 685.  
 22. *Lin.* *Syst. Nat.* 2. 710. 37.  
*Fuesly Inf. Helv.* 24. 464.  
 Habitat in Europae plantis.
- turca.* 28. *C. atra, abdomine flavo, alis apice fuscis, lunula hya-*  
*lina. Syst. Ent.* 685. 23. \*  
 Habitat in Europae plantis. *Mus. Dom. Banks.*
- maura.* 29. *C. nigra, pectore abdomineque sanguineis. Syst. Ent.*  
 686. 24. \*  
 Habitat in noua Hollandia. *Mus. Dom. Banks.*
- nebulosa.* 30. *C. nigra, elytris hyalinis, basi fuscis.*  
 Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*  
*Media. Corpus totum atrum, immaculatum. Elytra*  
*basi late fusca punctis duobus ferrugineo hyalinis,*  
*reliqua pars hyalina macula marginali atomisque fu-*  
*scis. Alae breuiore albohyalinae.*
- hyalina.* 31. *C. fusca, elytris fascia abbreviata margineque postico*  
*hyalinis. Syst. Ent. App.* 832. \*  
*Cercopis fenestrata* fusca, elytris fascia media maculis  
 que duabus marginalibus fenestratis. *Syst. Ent.*  
 688. 1. \*  
 Habitat in India orientali. *Dom. Edler.*
- pellucida.* 32. *C. grisea, elytris fascia media hyalina. Syst. Ent.*  
 686. 25. \*  
 Habitat in noua Hollandia. *Mus. Dom. Banks.*

33. *C. fusca*, alis albo hyalinis, margine fusco punctato. *Cynosbat-  
Syst. Ent. 686. 26.\* tis.*  
Habitat in Daniae Rosis.
34. *C. flava*, elytris albis, puncto fasciisque duabus ni- *Serratu-  
gris. Syst. Ent. 686. 27.\* lae.*  
Habitat in Angliae Carduis. *Mus. Dom. Banks.*
35. *C. alis fuscis, nervis albo nigroque punctatis. Syst. nervosa.  
Ent. 686. 28.\**  
*Cicada nervosa* fusca, alis hyalinis fusco maculatis, ner-  
uis punctatis. *Linn. Syst. Nat. 2. 709. 25. Fn. Sv.*  
882.  
*Cicada nervosa punctata* griseo fusca, alis hyalinis fu-  
sco maculatis, nervis punctatis. *Degeer Inf. 3. 182.*  
4. tab. 12. fig. 1. 2.  
*Geoff. Inf. 1. 415. 1.*  
*Scop. carn. 341.*  
*Fuesly Inf. Helv. 24. 459.*  
Habitat in Europae plantis.  
*Variat elytris fascia media fusca. Dom. de Hattorf.*
36. *C. atra viridi variegata, alis hyalinis, punctis tribus variega-  
costalibus nigris. Syst. Ent. 686. 29.\* ta.*  
Habitat in Germania, *Dom. de Hattorf.*
37. *C. fronte thoraceque glaucis, punctis quatuor ocel- lyncea.  
laribus nigris, elytris hyalinis, margine flavescente.*  
*Syst. Ent. 687. 30.\**  
Habitat in India orientali, *D. Koenig.*
38. *C. flava, elytris fusco auratis. Syst. Ent. 687. 31.\* fulgida.*  
Habitat in Angliae plantis. *Mus. Dom. Banks.*
39. *C. elytris flavescentibus, fusco punctatis. Syst. Ent. punctata.  
687. 32.\**  
*Cicada fusca pallida, elytris membranaceis virescenti-  
bus, scutello macula duplici triangulari. Geoff. Inf.*  
1. 417. 4.  
Habitat in Europae nemoribus.
40. *C. flava, alis albis apice fusco striatis. Syst. Ent. 687. Rosae.  
33. Linn. Syst. Nat. 2. 712. 50. Fn. Sv. 902.*

328 CLASS. VII. RYNGOTA. Cercopis.

*Cicada musciformis Rosae* viridi lutea, abdomine luteo, alis albidis apice hyalinis. *Degeer Inf.* 3. 103. 9.

*Geoff. Inf.* 1. 428. 28.

*Reaum. Inf.* 5. tab. 20. fig. 10 — 14.

*Frisch. Inf.* 11. 13. tab. 20.

*Fuesly Inf. Helv.* 24. 467.

Habitat in Rosae foliis.

*aurata.* 41. *C. flava*, elytris fulvo pictis, maculis quatuor nigris posticeque auratis. *Syst. Ent.* 687. 34. *Linn. Syst. Nat.* 2. 711. 48. *Fn. Sv.* 899. *Fuesl. Inf. Helv.* 24. 466.  
Habitat in Europae borealis plantis.

*Vlmi.* 42. *C. alis viridi luteis*, apicibus nigricantibus inauratis. *Linn. Syst. Nat.* 2. 711. 49. *Fn. Sv.* 900.

*Cicada musciformis Vlmi* viridi lutea, abdomine nigricante, alis superioribus apice nigricantibus. *Degeer Inf.* 3. 189. 8.

*Geoff. Inf.* 1. 427. 27.

Habitat in Europae Vlmis.

*cuspidata.* 43. *C. grisea*, capite depresso plano, apice fusco. *Syst. Ent.* 687. 35. \*  
Habitat in Angliae plantis. *Mus. Dom. Banks.*

*Quercus.* 44. *C. flaescens*, elytris sanguineo apice fusco maculatis. *Gen. Inf. Mant.* 298. \*.  
Habitat in Quercu, frequens in praedio comitis *Holt. Lekhoff.* prope Kiel.

157. CERCOPIS.

*grossa.* 1. *C. elytris fusco griseis*, macula marginali fulva cinerea.

Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*

Magna in hoc genere. Caput pallidum punctis aliquot bascos nigris. Thorax antice pallidus punctis nigris, postice emarginatus, fuscus. Elytra griseo fuscoque mixta macula marginali fulva majorique cinerea. Alae albiae fusco nebulosae. Abdomen fuscum ano supra nigro.

CLASS. VII. RYNGOTA. Cereopis. 329

2. C. thorace atro, fascia fulva, elytris fulvis, maculis *maculata*.  
apiceque nigris.  
Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*  
Statura praecedentis. Caput atrum, apice fulvum. Thorax lacuis alis fascia vndata fulva in medio. Elytra fulva atomis baseos, fascia abbreviata in medio apiceque late nigris. Abdomen supra rubrum, subtus nigrum.
3. C. atra, elytris maculis duabus fasciaque sanguineis. *sanguinolenta*.  
*Syst. Ent.* 688. 2.  
*Cicada sanguinolenta.* Linn. *Syst. Nat.* 2. 708. 22.  
*Cicada nigra*, elytris maculis sex rubris. *Geoff. Inf.* 1. 418. tab. 8. fig. 5.  
*Naturf.* 6. tab. 2.  
*Scop. carn.* 330.  
*Fuesly Inf. Helv.* 24. 456.  
Habitat in Europae Salicibus.
4. C. rufa, elytris nigris, fasciis duabus flavis. *Syst. Ent. cruentata*.  
688. 3. \*  
*Cicada rubra* alis deflexis fuscis, fasciis duabus flavis.  
Linn. *Syst. Nat.* 2. 111. 45.  
*Degeer Inf.* 3. 224. 20. tab. 33. fig. 8.  
Habitat in America meridionali.
5. C. sanguinea, thorace macula, elytris fasciis duabus *Carnifex*.  
nigris. *Syst. Ent.* 688. 4. \*  
Habitat in noua Hollandia. *Mus. Dom. Banks.*
6. C. flava, fasciis quatuor fuscis. *4fasciata*.  
*Cicada 4fasciata.* Linn. *Syst. Nat.* 2. 706. 9.  
*Cicada flava*, fasciis quatuor transuersis nigris. *Degeer*  
*Inf.* 3. 225. 21. tab. 33. fig. 11.  
Habitat in America meridionali.
7. C. fusca, elytris maculis binis albis lateralibus. *Syst. spumaria*.  
*Ent.* 688. 5.  
*Cicada spumaria.* Linn. *Syst. Nat.* 2. 708. 24. *Fn. Sv.* 881.  
*Cicada fusca*, fasciam duplici albida interrupta transuersa, *Geoff. Inf.* 1. 415. 2.

*Cicada spumaria Graminis* fusca, alis superioribus maculis albis. *Degeer Inf.* 3. 163. 1. tab. 11. fig. 1 — 21.  
*Locusta pulex Swammerdamii*, nobis *Cicadula*. *Raj. Inf.* 67.

*Ranatra bicolor*, capite nigricante. *Petiv. Gazoph.* tab. 61. fig. 9.

*Cicada fusca* alis superioribus maculis albis, in spuma quadam viuens. *Degeer Abt. Holm.* 1741. 221.

*Vermes spumans*. *Frisch. Inf.* 8. 26. tab. 12.

*Locusta germanica*. *Roef. Inf.* 2. *Gryll.* tab. 23.

*Sulz. Inf.* tab. 10. fig. 64.

*Schaeff. Elem.* tab. 42.

*Fuesly Inf. Helv.* 450.

Habitat in Europae plantis variis, latitans intra spumam, graminibus, rosis aliisque adhaerens.

*lineata*. 8. *C. flauescens*, coleoptris striis tribus nigris.

Habitat in Germania. *Dom. de Hattorf.*

Media. Caput et thorax flauescencia linea dorsali nigra, quae per elytrorum suturam continuatur. Elytra flauescencia stria longitudinali futuraeque nigris.

*coleoptra*. 9. *C. elytris* totis coriaceis, alas obtegentibus, griseis fusco irroratis.

Habitat in Germaniae plantis.

Media. Caput et thorax flauescencia. Abdomen fuscum. Elytra lata, tota coriacea, pone basin in angulum paruum excurrunt, grisea atomis plurimis punctis 2 — 3 distinctis fuscis. Pedes grisei.

*rustica*. 10. *C. grisea* immaculata, alis albis. *Syst. Ent.* 689. 6.\*

Habitat in Europae plantis.

*gibba*. 11. *C. nigra*, elytris albo maculatis. *Syst. Ent.* 689. 7.\*

Habitat in Daniae nemoribus.

*unifascia*. 12. *C. cinerea*, elytris fascia obliqua fusca.

Habitat in Agro Pedemontano. *D. Allioni.*

Maior praecedente. Corpus cinereum fascia elytrorum obliqua fusca pone medium. Abdomen supra atrum.

*bifascia*. 13. *C. flauescens*, elytris fuscis, fasciis duabus albis. *Syst. Ent.* 689. 8.

*Cicada*

CLASS. VII. RYNGOTA. Notonecta. 331

*Cicada bifasciata* fusca, fasciis alarum binis albis. *Linn. Syst. Nat.* 2. 706. 10. *Fn. Sv.* 898.

*Cicada trifasciata* nigro fusca, fasciis tribus transuersis, capite subtus flauo. *Degeer Inf.* 3. 186. 6. tab. 11. fig. 25.

Habitat in Europae borealis plantis.

14. *C. nebulosa*, vertice punctis duobus abdomineque basi atris. *Syst. Ent.* 898. 9. *Populi.*

*Cicada Populi* subtus flauescens, fronte punctis duobus nigris. *Linn. Syst. Nat.* 2. 709. 32. *Fn. Sv.* 891.

Habitat in Europae borealis plantis.

158. NOTONECTA.

1. *N. elytris* griseis, margine fusco punctato apice bifidis. *glauc.*  
*Syst. Ent.* 689. 1. *Linn. Syst. Nat.* 2. 712. 1. *Fn. Sv.* 903.

*Notonecta* capite luteo, elytris fusco croceoque variegatis, scutello atro. *Geoff. Inf.* 1. 476. 1. tab. 9. fig. 6.

*Nepa Notonecta* cylindracea, elytris griseo flauis margine fusco punctatis, scutello nigro. *Degeer Inf.* 3. 382. 5. tab. 18. fig. 16. 17.

*Cicada aquatica* *Mouffeti. Raj. Inf.* 58.

*Notonecta* vulgaris nigro pallidoque mixta. *Petiv. Gazoph.* tab. 72. fig. 6.

*Cimex aquaticus* angustior. *Frisch. Inf.* 6. 28. tab. 13. *Cimex* aquaticus. *Roesl. Inf.* 3. tab. 27.

*Bradl. nat.* tab. 26. fig. 2. E.

*Huffnag. Inf.* tab. 12. fig. 19.

*Swiz. Inf.* tab. 10. fig. 67.

*Schaeff. Elem.* tab. 90.

— — *Icon.* tab. 33. fig. 5. 6.

*Fuesly Inf. Helv.* 24. 468.

Habitat in Europae aquis, piscibus molesta.

2. *N. grisea*, postice nigra, scutello atro, vtrinque puncto *americana.*  
bafcos flauo. *Syst. Ent.* 690. 2. \*

Habitat in Americae aquis. *Mus. Tottianum.*

3. *N.*



*niuea.* 3. *N. elytris albis immaculatis, apice rotundatis.* *Syst. Ent.* 690. 3.\*

Habitat in India. *D. Koenig.*

*minutissima.* 4. *N. grisea, capite fusco, elytris truncatis.* *Syst. Ent.* 690. 4.

*Notonecta minutissima* elytris cinereis, maculis fuscis longitudinalibus. *Linn. Syst. Nat.* 2. 713. 3. *Fn. Sv.* 905.

*Notonecta* cinerea anelytra. *Geoff. Inf.* 1. 477. 2.

*Fuesly Inf. Helv.* 24. 470.

Habitat in Europae aquis.

Laruae *N. glaucae* simillima, et perfecta.

### 159. S I G A R A.

*friata.* 1. *S. elytris pallidis, lineolis transuersis undulatis numerosissimis fuscis.* *Syst. Ent.* 691. 1.

*Notonecta friata* elytris pallidis lineolis transuersis undulatis striatis. *Linn. Syst. Nat.* 2. 712. 2. *Fn. Sv.* 904.

*Notonecta vulgaris* compressa fusca. *Petiv. Gazoph.* tab. 72. fig. 7.

*Corixa.* *Geoff. Inf.* 1. 478. 1. tab. 9. fig. 7.

*Nepa friata* oblonga nigro fusca, lineolis transuersis numerosissimis griseis striata. *Degeer Inf.* 3. 389. 6. tab. 12. fig. 1. 2.

*Roes. Inf.* 2. 177. tab. 19.

*Ioblot. microsc.* 1. 1. tab. 7. fig. 2. 3.

*Scharff. Eiern.* tab. 50.

— — *Icon.* tab. 97. fig. 2.

Habitat in Europae aquis frequens.

Magnitudinis valde variat.

*recepta.* 2. *S. elytris totis coriaceis fuscis, margine exteriori flavo.* *Gm. In. Mant.* 208.\*

Habitat in Kilise aquis. *Dum. de Schest.*

### 160. N E P A.

*granulata.* 1. *N. crandata* testacea, flavo maculata. *Syst. Ent.* 691. 1.

*Nepa granulata* testacea, scutello laevi, alis albis maculis virentibus. *Linn. Syst. Nat.* 2. 713. 1. *Mus. Lud.* 1. 1. 104.

*Nepa*

*Nepa surinamensis* grisea, fusco nebuloſa, corpore oblongo plano, thorace lacui, pedibus maculatis.

*Degeer Inf.* 3. 379. 4.

Inſectum magnitudinae bruſci. *Margr. Braſ.* 258. fig. 259.

*Merian, Sur.* tab. 66.

*Roef. Inf.* 3. tab. 26.

*Gronov. Zooph.* 685.

Habitat in Americae aquis.

Antennae palmato quadrifidae, laciniae arcuatis.

2. *N. ecaudata* subrotunda, pallide fusca, tibiis anticis *annulatis*, pallidis fusco annulatis.

Habitat in Coromandel. *Mus. Dom. Banks.*

Reliquis magis rotundata, latior, depressa plana, pallide fusca, thorace scutelloque antice obscurioribus.

Pedes fusco pallidoque varii tibiis anticis pallidis annulis fuscis.

3. *N. ecaudata* fusca, thoracis elytrorumque margine antice albido. *rustica*.

*Notonecta rustica* ecaudata, corpore fusco immaculato.

*Syst. Ent.* 691. 2.

Habitat in Coromandel. *Mus. Dom. Banks.*

Reliquis minor. Caput fuscum antice pallidum. Thorax fuscus margine laterali et postico pallidis. Elytra fusca margine antice pallido.

Ova aggregata, erecta in dorso saepius circumfert.

4. *N. ecaudata*, scutello rugoso, alis niueis. *Syst. Ent. fusca.* 692. 3. *Linn. Syst. Nat.* 2. 713. 5. *Mus. Lud. Vlr.* 166.

Habitat in Americae meridionalis aquis.

5. *N. cauda bifeta* corpore ovato. *Syst. Ent.* 692. 4. *cinerea*.

*Nepa cinerea* thorace inaequali, corpore oblongo ovato. *Linn. Syst. Nat.* 2. 714. 5. *Fn. Sv.* 906.

*Nepa* corpore ovato. *Geoff. Inf.* 1. 481. 2.

*Nepa Scorpio aquaticus* grisea corpore ovato depresso, thorace inaequali. *Degeer Inf.* 3. 361. 1. tab. 18. fig. 1.

*Scorpio palustris* ad cimices referenda. *Raj. Inf.* 58.

*Scorpio palustris.* *Mouff. Inf.* 321.

*Scorpio-*

*Scorpiones* aquatici. *Jonf. Inf. tab. 25. fig. 1. 2.*

*Scorpius* vulgaris aquaticus. *Petiv. Gazoph. tab. 74. fig. 4.*

*Gmex* aquaticus. *Roef. Inf. 3. tab. 21. fig. 6. 7. 8.*

*Bradl. nat. tab. 26. fig. 2. 6.*

*Swammerd. Bibl. 1. tab. 3. fig. 4.*

*Huffn. pict. tab. 11. fig. 2.*

*Frisch. Inf. 7. tab. 15.*

*Sulz. Inf. tab. 10. fig. 68.*

*Schaeff. Elem. tab. 69.*

— — *Icon. tab. 33. fig. 7. 9.*

*Fuesly Inf. Helv. 24. 471.*

Habitat in Europae aquis.

Antennae palmato bifidae. Abdomen supra rubrum.

*maculata*. 6. *N. nigra*, thorace scutelloque ferrugineo maculatis, alis fuscis. *Syst. Ent. 692. 5. \**

Habitat in Indiae orientalis aquis. *D. Koenig.*

*linearis*. 7. *N. cauda bifeta corpore lineari. Syst. Ent. 692. 6.*

*Nepa linearis*, manibus spina laterali pollicatis. *Linn.*

*Syst. Nat. 2. 714. 7. Fn. Sv. 908.*

*Nepa* corpore lineari. *Geoff. Inf. 1. 480. tab. 10. fig. 1.*

*Nepa linearis* corpore angustissimo elongato, thorace

longo, tibiis anticis in medio spina laterali. *Deger*

*Inf. 3. 369. 2. tab. 19. fig. 1. 2.*

*Locusta* aquatica *Mouffeti. Raj. Inf. 59.*

*Swammerd. Bibl. nat. 1. 233. tab. 3. fig. 9.*

*Roef. Inf. 3. 141. tab. 23.*

*Sulz. Hist. Inf. tab. 10. fig. 4.*

*Fuesly Inf. Helv. 25. 473.*

*Frisch. Inf. 7. 24. tab. 16.*

*Gronov. Zooph. 683.*

*Schaeff. Icon. tab. 5. fig. 5. 6.*

Habitat in Europae aquis.

Antennae mihi haud visae.

## 161. N A V C O R I S.

*cimicoides*. 1. *N. abdominis margine serrato. Syst. Ent. 693. 1. Geoff. Inf. 1. 474. 1. tab. 9. fig. 5.*

*Nepa cimicoides. Linn. Syst. Nat. 2. 714. 6. Fn. Sv. 907.*

*Nepa*

**CLASS. VII. RYNGOTA. Acanthis. 335**

*Nepa Naucoris* ovata fusca viridi variegata, abdominis margine serrato. *Degeer Inf.* 3. 375. 3. tab. 19. fig. 8. 9.

*Cimex aquaticus* lator. *Frisch. Inf.* 6. 31. tab. 14.

*Roef. Inf.* 3. 176. tab. 28.

*Gronov. Zooph.* 686.

*Schaeff. Elem.* tab. 87.

— — *Icon.* tab. 33. fig. 3. 4.

*Sulz. Hist. Inf.* tab. 10. fig. 3.

*Fuesly Inf. Helv.* 24. 472.

Habitat in Europae aquis.

2. N. abdominis margine integro. *Syst. Ent.* 693. 2. \* *nepaeformis.*  
Habitat in Antigua Insula. *Dom. Drury.*

**162. ACANTHIA.**

1. A. aptera. *Syst. Ent.* 693. 1. *leclularia.*  
*Cimex leclularius.* *Linn. Syst. Nat.* 2. 715. 1. *Fn. Sv.* 909.

*Geoff. Inf.* 1. 434. 1.

*Fuesly Inf. Helv.* 25. 474.

*Degeer Inf.* 3. tab. 17. fig. 9.

*Raj. Inf.* 7.

*Mouff. Inf.* 259.

*Bonann. Microsc.* tab. 65.

*Ledermüll. Microsc.* tab. 52. 63.

*Sulz.* tab. 10. fig. 69.

Habitat in Europae domibus.

Animal foetidum, nocturnum, sanguisugum pellitur nido carbonum, oleo therebinthinae accenso, *Mentha arvensis*, oleo vitrioli, tabaci, *Thlaspi*, *Sambuco*, *Cannabi*.

2. A. nigra, elytris coriaceis, abdomine longioribus apice hyalino striatis. *Gen. Inf. Mant.* 298. \* *Zosteras.*

*Acanthia pellucens* elytris pellucido punctatis, apice striatis. *It. Norwag.* d. 12. Iul. \*

Habitat ad littora maris, velocissime supra fueos et *Zosteras* cursitans Kiliae frequens. *Dom. Lund.*

3. A. aptera, elytris coriaceis nigro griseis. *Gen. Inf. coriacea.* *Mant.* 299. \*

*Cimex*

*Cimex grylloides* apterus niger, elytris ouatis margine luteis, abdomine breuioribus. *Linn. Syst. Nat.* 2. 717. 13. *Fn. Sv.* 910.

Habitat Kiliac. *Dom. Lund.*

*pedefris.* 4. *A.* aptera elytris coriaceis connatis abbreviatis cinereis, ano setoso.

Habitat in Agro Pedemontano. *D. Allioni.*

An potius Puppa alius speciei? Corpus paruum totum cinereum. Coleoptra lata abdomine dimidio breuiora in medio connata, canaliculata. Pedes pallidi. Anus setis plurimis nigris longitudine dimidii abdominis fasciculatis.

*clauicornis.* 5. *A.* elytris reticulato punctatis, antennis clauatis. *Syst. Ent.* 694. 2.

*Cimex clauicornis* coleoptratus, elytris neruoso carinatis reticulato punctatis, antennis clauatis. *Linn. Syst. Nat.* 2. 717. 16. *Fn. Sv.* 911.

*Cimex* ex albo fuscoque cinereus elytrorum thoracique margine punctato, antennis subclauatis. *Geoff. Inf.* 1. 461. 56.

*Reaum. Inf.* 3. tab. 34. fig. 1 — 4.

*Fuesly Inf. Helv.* 25. 480.

Habitat in Europae hortis.

*Serratulae.* 6. *A.* nigra, elytris pallescentibus, alis apice fuscis. *Syst. Ent.* 694. 3. \*

Habitat in Angliae Serratulis. *Mus. Dom. Banks.*

*littoralis.* 7. *A.* elytris fordidis, albo punctatis, corpore nigro. *Syst. Ent.* 694. 4.

*Cimex littoralis.* *Linn. Syst. Nat.* 2. 717. 14. *Fn. Sv.* 914.

*Cimex* ouatus nigricans, dorso gibbo, elytris maculis dilute fuscis diaphanis, alis breuissimis. *Degeer Inf.* 3. 278. 21. tab. 14. fig. 17. 18.

Habitat in Europae littoribus.

*rugosa.* 8. *A.* elytris pallidis, corpore oblongo, femoribus anticis crassissimis. *Syst. Ent.* 694. 5.

*Cimex rugosus.* *Linn. Syst. Nat.* 2. 717. 15. *Fn. Sv.* 916.

Habitat in Europa boreali.

CLASS. VII. RYNGOTA. Acanthia. 337

9. A. thorace lunato, margine prominente, abdomine fer- *lunata*.  
rato. *Syst. Ent.* 694. 6. \*  
Habitat in Indiis.
  
10. A. membranacea, abdominis margine imbricatim se- *corticalis*.  
cto, corpore nigro. *Syst. Ent.* 695. 7.  
*Cimex corticalis*. *Linn. Syst. Nat.* 2. 718. 17. *Fn. Sv.*  
917.  
*Schaeff. Icon.* tab. 51. fig. 6. 7.  
Habitat in Europae arboribus syluaticis.
  
11. A. nigra, abdomine laeui brunneo, alis pallidis. *Syst. laeuis*.  
*Ent.* 695. 8. \*  
Habitat in Anglia. *Mus. Dom. Banks.*
  
12. A. membranacea, thorace denticulato, capite spinoso, *Betulae*.  
elytris antice dilatatis. *Syst. Ent.* 695. 9. \*  
*Cimex Betulae*. *Linn. Syst. Nat.* 2. 718. 18. *Fn. Sv.* 918.  
*Cimex depressus Betulae* ouatus depressus nigro fuscus,  
thorace denticulato, capite muricato, elytris faminae  
abdomine angustioribus. *Degeer Inf.* 3. 305. 36. tab.  
15. fig. 16. 17.  
Habitat in Europae borealis Betula alba.
  
13. A. abdomine marginato fusco, apice flauescente, tibiis *crassipes*.  
anticis incrassatis. *Syst. Ent.* 695. 10. \*  
Habitat Dresdae Saxonum.
  
14. A. membranacea, abdomine flauo, fascia nigra, tho- *erosa*.  
racis margine sinuato, tibiis anticis incrassatis.  
*Cimex erosus* membranaceus, abdomine flauo, fascia ni-  
gra, thorace rugoso, tibiis anticis incrassatis. *Linn.*  
*Syst. Nat.* 2. 718. 19.  
*Cimex Scorpio* ouatus depressus angulatus, antennis  
clauatis, abdomine flauo, fascia nigra, pedibus anti-  
cis cheliformibus. *Degeer Inf.* 3. 350. 25. tab. 35. fig.  
13. 14.  
Habitat in America meridionali.
  
15. A. ferruginea, thorace marginato, margine antice scu- *umbracu-*  
telloque apice albis. *Syst. Ent.* 695. 11. \* *lata*.  
Habitat Hafniae. *D. Torstenfson.*

*ferrata*. 16. A. thoracis margine ferrato, corpore griseo, alis albis fusco maculatis. *Syst. Ent.* 695. 12. \*

Habitat - - - - *Mus. Tottianum*.

*Cardui*. 17. A. scutello lineis tribus eleuatis, antennis apice nigris. *Syst. Ent.* 696. 13.

*Cimex Cardui*. *Linn. Syst. Nat.* 2. 718. 21. *Fn. Sv.* 920.

*Cimex depressus* ouatus griseus nigro maculatus, antennis clauatis, thorace lineis tribus longitudinalibus eleuatis. *Degeer Inf.* 3. 309. 38. tab. 6. fig. 1.

Habitat in *Cardui* capitulis *Europae borealis*.

*Pyri*. 18. A. thorace trialato, scutello foliato, elytris reticulatis basi gibbis. *Syst. Ent.* 696. 14. \*

*Cimex* antennis clauatis, thorace elytrisque corpore multo latioribus diaphanis reticulatis, fascia duplici transuersa. *Geoff. Inf.* 1. 461. 57.

Habitat in foliorum *Pyri* communis pagina inferiori, ca maculans. *Prof. Hermann*.

*rhombica*. 19. A. thorace acutangulo, abdomine dilatato rhombico, postice sexdentato. *Syst. Ent.* 696. 15.

*Cimex rhombeus*. *Linn. Syst. Nat.* 2. 718. 22.

Habitat in *Africae arboribus*.

## 163. C I M E X.

### \* Scutellati scutello longitudine abdominis.

*Stockerus*. 1. C. scutellaris ouatus, corpore viridi, maculis nigris, abdomine ferrugineo. *Syst. Ent.* 696. 1.

*Cimex Stockerus* scutellaris, corpore viridi, maculis nigris. *Linn. Syst. Nat.* 2. 715. 2.

*Stockerus benghalensis* viridis nigro maculatus. *Petit. Gazoph.* 34. tab. 21. fig. 12.

Habitat in *China*.

Affinis sequenti, subtus ferrugineus lateribus cyaneis.

*nobilis*. 2. C. scutellaris oblongus coerulco auratus nigro maculatus. *Syst. Ent.* 697. 2. *Linn. Syst. Nat.* 2. 716. 3. *Amoen. acad.* 6. 400. 46.

*Sulz.*

*Sulz. Hist. Inf. tab. 11. C.*

*Schroet. Abhandl. 1. tab. 1. fig. 9.*

Habitat in Aſia.

Subtus rufus faſciis lateralibus coeruleo auroque nientibus.

3. *C. scutellaris*, thorace ſcutelloque aureis, thorace punctis duobus, ſcutello maculis duabus coeruleſcentibus. *Syſt. Ent. 697. 3. \** *regalis.*

Habitat in noua Hollandia. *Muf. Dom. Banks.*

4. *C. scutellaris*, thorace ſcutelloque rufis, abdomine cyaneo, margine ſanguineo. *Syſt. Ent. 697. 4. \** *imperialis.*

Habitat in noua Hollandia. *Muf. Dom. Banks.*

5. *C. scutellaris niger obſcurus*, abdominis margine ferrugineo, pedibus pallidis. *raſticus.*

Habitat in Italia. *D. Allioni.*

Statura praeſcedentium, at paullo minor, obſcurus minime nitens. Caput piceum, thorax antice piceus, poſtice niger punctatus. Scutellum nigrum, immaculatum. Abdomen nigrum margine ferrugineo, ano tamen nigro. Pedes omnes pallidi.

6. *C. scutellaris viſcens*, maculis annularibus nigris. *annulus.*  
*Syſt. Ent. 697. 5. \**

Habitat in Senegalia. *Mauduit.*

Alae apice nigrae.

Variat colore corporis obſcure pallido.

7. *C. scutellaris ſupra teſtaceus*, thorace punctis duobus, ſcutello quatuor cyaneo nigris. *6punctatus.*

Habitat in America meridionali. *Muf. D. Hunter.*

Medius. Antennae nigrae. Caput rufo teſtaceum lineis duabus apice coeuntibus nigris. Thorax teſtaceus punctis duobus magnis dorſalibus nigris. Scutellum teſtaceum punctis duorum parium nigris. Subtus niger abdomine fuluo maculato. Pedes nigri femoribus fuluis.

8. *C. scutellaris ſupra rufus*, maculis diſformibus nigris. *Drurasi.*  
*Syſt. Ent. 697. 6.*



240 CLASS. VII. RYNGOTA. Cimex.

*Cimex Druraci* scutellaris sanguineus, punctis maculisque nigris, subtus atro coerulescens. *Linn. Mant.* 534.

*Drury Inf.* 1. tab. 42. fig. 1. 5.

*Sulz. Hist. Inf.* tab. 10. fig. 5.

Habitat in America meridionali.

**Fabricii.** 9. *C. scutellaris* subpurpurascens fuluo punctatus. *Syst. Ent.* 698. 7.

*Cimex Fabricii* scutellaris purpureus punctis sanguineis, subtus atro coerulescens. *Linn. Mant.* 534.

Habitat in Caienna.

Alter, sexus supra rufescens, alter obscure coerulescens.

**Schulzii.** 10. *C. scutellaris* atro aeneus, scutello vtrinque macula bascos coccinea.

Habitat Caiennae. *D. Schulz.*

Praecedentibus minor. Statura et magnitudo *C. scarabaeoidis*. Corpus totum atrum aeneo colore splendens macula magna ovata coccinea vtrinque ad basin scutelli. Antennae et pedes fuscii.

**paganus.** 11. *C. scutellaris* cyaneus, scutello abdomineque rufis, cyaneo maculatis. *Syst. Ent.* 698. 8.

Habitat in noua Hollandia. *Mus. Dom. Banks.*

Variat thorace macula rufa et abdomine punctis quatuor cyaneis in medio.

**illustris.** 12. *C. scutellaris* glaucus, thorace scutelloque punctis duobus fuscis.

Habitat ad Cap. bon. sp. *Dom. D. Vahl.*

Magnus. Caput glaucum linea media abbreviata maculisque duabus bascos fuscis. Thorax subangulatus glaucus vtrinque macula albida obsoleta et in hac punctum fuscum. Scutellum magnum glaucum in medio puncto vtrinque fusco. Antennae et pedes fuscii.

**lineola.** 13. *C. scutellaris* supra cyaneus, capite thoraceque linea dorsali, scutello punctis duobus rubris.

Habitat ad Cap. bon. sp. *Mus. Dom. Banks.*

Medius. Antennae nigrae. Caput cyaneum linea dorsali rubra, quae tamen apicem haud attingit. Thorax

rax cyaneus, nitidus, linea dorsali rubra. Scutellum abdomen totum tegens puncto vtrinque baseos rubro. Elytra cyanea alis fuscis. Pectus flavum lateribus cyaneis. Abdomen basi rufescens, tunc nigrum apice flavescentia. Pedes cyanei femoribus basi flavis.

14. *C. scutellaris niger*, lineis tribus flavescentibus. *trilineatus*.  
 Habitat in America meridionali. *Mus. D. Hunter*.  
 Minor *C. lineato* supra totus niger lineis tribus flavescentibus, quae in capite magis rubrae fere coeunt. Subtus niger viridi colore tinctus abdomine vtrinque linea punctorum rubrorum.
15. *C. scutellaris ruber*, thorace lineis quinque, scutello *nigrolitibus* nigris, abdomine flavo, punctis nigris. *Syst. neatus*.  
*Ent.* 698. 9. *Linn. Syst. Nat.* 2. 716. 6.  
*Cimex rotundatus ruber* supra fasciis longitudinalibus infra punctis nigris, scutello amplo totum fere abdomen obtegentē. *Geoff. Inf.* 1. 468. 68.  
*Scop. carn.* 352.  
*Mouff. Inf.* 174. fig. 5.  
*Gronov. Zooph.* 689.  
*Schaeff. Elem.* tab. 44. fig. 1.  
 — — *Icon.* tab. 2. fig. 3.  
*Sulz. Hist. Inf.* tab. 10. fig. 6.  
*Fuesly Inf. Helv.* 25. 477.  
 Habitat in Europa australiori.
16. *C. scutellaris supra rufus*, thorace punctis decem, scutello *semipunctato* lineis quatuor nigris. *Syst. Ent.* 698. 10. \* *status*.  
 Habitat in America, in Europa australi. *D. Allioni*.
17. *C. scutellaris cyaneus*, scutello basi apiceque flavescente, abdominis margine flavo nigro punctato. *Syst. anchorago*.  
*Ent.* 699. 11. \*  
 Habitat in America boreali. *D. Solander*.  
 Variat sexu, mas rufus, vbi femina flava.
18. *C. scutellaris*, corpore flavescente, supra lituris longitudinalibus nigris. *Linn. Syst. Nat.* 2. 716. 7. *Syst. grammicus*.  
*Ent.* 699. 12.  
 Habitat in Africa.

*Pedemontanus*. 19. *C. scutellaris* fuscus, atomis numerosissimis albis.  
Habitat in Agro Pedemontano. *D. Allioni*.  
Paullo minor præcedente. Caput obscure fuscum im-  
maculatum. Thorax et scutellum fusca atomis siue  
punctis parvis numerosissimis sparsis albis. Pedes  
nigri,

*Maurus*. 20. *C. scutellaris* cinereus; scutello basi punctis duobus  
albis. *Syst. Ent.* 699. 13. *Linn. Syst. Nat.* 2. 716. 5.  
*Fn. Sv.* 913.

*Schaeff. Icon.* tab. 43. fig. 3. 4. 15. 16.

*Sulz. Hist. Inf.* tab. 4. fig. d.

*Fuesly Inf. Helv.* 25. 476.

*Scop. carn.* 352.

Habitat in Oriente, apud nos quadruplo minor.

*Hottentotta*. 21. *C. scutellaris* ferrugineus, immaculatus. *Syst. Ent.*  
699. 14. \*

*Cimex* fuscus, scutello totum abdomen obtegente im-  
plissimo. *Geoff. Inf.* 1. 435. 3. ?

Habitat in Arabia. *Prof. Forskahl*.

*albolineatus*. 22. *C. scutellaris* griseus albo striatus.  
*tus*. Habitat in Italia. *D. Allioni*.

Minor. Caput griseum, stria media albida. Thorax  
griseus stria quinque fere obsoletis inprimis inter  
mediis duabus. Scutellum griseum stria quinque  
marginibusque albis. Abdomen fuscum margine pu-  
sco, punctis aliquot nigris. Pedes fusci.

*irroratus*. 23. *C. scutellaris* virescens fusco irroratus. *Syst. Ent.* 699.  
15. \*

Habitat in Americae meridionalis Insulis. *Dom. &*  
*Rohr*.

*lanatus*. 24. *C. scutellaris* aeneo niger griseo hirtus.

*Cimex lanatus*. *Pall. It.* 2. 729. 32.

Habitat in Sibiria. *Pallas. Mus. Dom. Banks*.

Corpus totum oblongum, aeneo nigrum, immaculatum,  
pilis elongatis griseis hirtum.

*scarabaeoides*. 25. *C. scutellaris*, corpore atro aeneo. *Syst. Ent.* 700. 16.  
*Linn. Syst. Nat.* 2. 716. 4. *Fn. Sv.* 912.

*Cimex*

CLASS. VII. RYNGOTA. Cimex. 343

*Cimex hemisphaericus* nigro aeneus, scutello totum abdomen obtegente, amplissimo. *Geoff. Inf.* 1. 435. 2.

*Sulz. Inf.* 26. tab. 11. fig. 70.

*Fuesly Inf. Helv.* 25. 475.

Habitat in Ranunculi floribus.

26. *C. scutellaris* nigro aeneus scutelli abdominisque margine pedibusque pallidis.

Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*

Affinis sequenti, at paullo maior, et caput et thoracis margo corpori concolora.

27. *C. scutellaris* nigro aeneus, margine omni pedibusque flavis. *Syst. Ent.* 700. 17. \*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

28. *C. scutellaris*, scutello fuliginoso, lituris quinque nigris, postica alba. *Syst. Ent.* 700. 18. *Linn. Syst. Nat.* 2. 716. 8. *Fn. Sv.* 914.

*Schaeff. Icon.* tab. 11. fig. 10 — 12.

*Fuesly Inf. Helv.* 25. 478.

Habitat in Europa, in Oriente. *Forskahl.*

29. *C. scutellaris* niger, scutello lituris tribus albis. *Syst. Ent.* 700. 19. \*

Habitat in Arabia. *Prof. Forskahl.*

30. *C. scutellaris* obscurus, scutello scabro ante apicem tuberculato.

Habitat in Italia. *D. Allioni.*

Praecedente paullo minor. Corpus totum fuscum obscurum paullo ferrugineo mixtum. Thorax fascia media impressa inaequali. Scutellum magnum, punctis eleuatis sparsis scabrum, et ante apicem tuberculo magno eleuato, obtuso. Pedes nigricantes genubus pallidis tibiisque ferratis.

31. *C. scutellaris*, ater, scutello basi pedibusque griseis. *Syst. Ent.* 700. 20. \*

Habitat in Angliae nemoribus.

- Arabs.* 32. *C. scutellaris*, thorace spinoso, corpore ovato fluido, ano bidentato. *Syst. Ent.* 700. 21. *Linn. Syst. Nat.* 2. 716. 9. *Mus. Lud. Vlr.* 168.  
*Cimex sylvestris foetens*, viridis, triangularis. *Sloan. Jamaic.* 2. tab. 237. fig. 15. 16.  
 Habitat in America meridionali.

- Indicus.* 33. *C. scutellaris*, thorace subangulato, corpore supra viridi, subtus flavo, ano bidentato. *Syst. Ent.* 701. 22. *Linn. Syst. Nat.* 2. 717. 11. *Mus. Lud. Vlr.* 170.  
*Degeer Inf.* 3. tab. 34. fig. 3.  
 Habitat in Indiis.

**\*\* Spinosi, thorace vtrinque spina armato.**

- taurus.* 34. *C. thorace spinoso griseus*, thoracis spinis porrectis compressis truncatis.  
 Habitat in Coromandel. *Mus. Dom. Banks.*  
 Magnus. Corpus totum supra griseum, subtus flavescens, pedibus nigro punctatis. Spinæ thoracis crassæ, compressæ, obtusissimæ.

- bidens.* 35. *C. thorace acute spinoso, griseus, antennis rufis.* *Syst. Ent.* 701. 23. *Linn. Syst. Nat.* 2. 718. 23. *Fn. Sv.* 921.  
*Cimex ovatus fuscus*, thorace spinis longioribus, subtus maculis rubris, antennis rufis, abdomine subtus macula nigra. *Degeer Inf.* 3. 259. 6. tab. 13. fig. 9.  
*Fuesly Inf. Helv.* 25. 482.  
 Habitat in Europa.  
 Puncta aliquot rufa in pectore.

- sanguinipes.* 36. *C. thorace obtuse spinoso, fuscus, scutello apice albidio, abdominis margine maculato, pedibus rufis.*  
 Habitat in Italia. *D. Allioni.*  
 Statura omnino *C. grisei*. Antennæ nigrae articulo ultimo basi pallido. Caput et thorax obtuse spinosus, fusca, immaculata. Scutellum fuscum macula parva, vtrinque ad basin flava, apiceque album. Abdomen subtus flavum, fusco maculatum margine albo nigroque vario. Pedes rufi.

- vulpes.* 37. *C. thorace obtuse spinoso, griseus, pedibus rubris.* *Syst. Ent.* 701. 24. *Linn. Syst. Nat.* 2. 719. 24. *Fn. Sv.* 922.  
*Cimex*

CLASS. VII. RYNGOTA. Cimex. 345.

*Cimex* ouatus fuscus, thorace angulato spinoso scutelli apice pedibusque rubris. *Degeer Inf.* 3. 253. 2.

*Cimex* sylvestris, corpore breviorè fuscus, scapulis magis exstantibus, macula e flavo rubente centro crucis dorsalis. *Raj. Inf.* 54. 2.

*Schaeff. Icon.* tab. 57. fig. 6. 7.

*Fuesly Inf. Helv.* 25. 483.

Habitat in Europa frequens.

38. *C.* thorace obtuse spinoso subuirefcente, elytris griseis, *luridus*, macula fusca, clypeo emarginato. *Syst. Ent.* 701. 25. \*

Habitat in Angliae nemoribus. *Mus. Dom. Banks.*

39. *C.* thorace obtuse spinoso, subferrugineus, spinis antennisque nigris. *Syst. Ent.* 701. 26. \* *nigricornis.*

Habitat in Saxonia.

40. *C.* thorace spinoso, oblongus supra rufescens, subtus *ictericus*, flavus. *Syst. Ent.* 702. 27. *Linn. Syst. Nat.* 2. 719.

25. *Amen. acad.* 6. 399. 41.

Habitat in America.

41. *C.* thorace subspinoso, fuscus, abdominis margine variegato, tibiis annulo albo. *punctatus.*

*Cimex punctatus* ouatus punctatus, thorace subspinoso, tibiis annulo albo. *Linn. Syst. Nat.* 2. 720. 34. *Fn. Sv.* 924.

*Cimex* ouatus niger subaeneus, thorace obtuse spinoso tibiis annulo albo. *Degeer Inf.* 3. 269. 14.

Habitat in Europa boreali.

Antennae nigrae. Corpus supra obscure fuscum immaculatum subtus griseum, abdominis margine flavo maculato.

42. *C.* thorace subspinoso supra nigricans, thoracis margine scutellique apice albis. *albipes.*

Habitat in Italia. *D. Allioni.*

Statura praecedentis. Corpus supra nigricans, obscurum, subtus pallidum. Antennae albo nigroque variae. Thoracis margo albidus. Scutellum apice album. Abdominis margo albo nigroque varius. Pedes pallidi.

*dentatus*. 43. C. thorace acute spinoso ferrulato, corpore cinerea nigroque vario. *Syst. Ent.* 702. 28. \*

Habitat in India orientali. *D. Koenig.*

*floridanus*. 44. C. thorace spinoso, niger rubro varius, scutello maculis tribus rubris. *Syst. Ent.* 702. 29. \* *Linn. Syst. Nat.* 2. 719. 26.

Habitat in America.

*armatus*. 45. C. thorace acute spinoso, scutello nigro, punctis duobus apiceque testaceis, antennis pedibusque rubris. *Syst. Ent.* 702. 30. \*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

*haemorrhous*. 46. C. thorace spinoso, niger, abdomine rufo, elytris punctis linearibus quinque nigris. *Syst. Ent.* 702. 31. *Linn. Syst. Nat.* 2. 719. 27. *Amoen. acad.* 6. 400. 45.

Habitat in America.

*apustulatus*. 47. C. thorace spinoso, niger, elytris liuidis, capite punctis duobus coccineis. *Syst. Ent.* 703. 32. \*

*Cimex apustulatus* oblongiusculus niger, thorace spinoso, punctis verticis duobus coccineis. *Linn. Syst. Nat.* 2. 719. 29.

*Cimex balteatus* oblongus niger, thorace spinoso, abdomine rubro, elytris antice testaceis nigro maculatis. *Degeer Inf.* 3. 334. 8. tab. 34. fig. 10.

Habitat in America meridionali.

*punicus*. 48. C. thorace spinoso, niger, scutello lunula apiceque rubris. *Syst. Ent.* 703. 33. *Linn. Syst. Nat.* 2. 719. 34.

Habitat in Africa.

*Ypsilon*. 49. C. thorace spinoso, liuidus, scutello y flauo inscripto. *Syst. Ent.* 703. 34. *Linn. Syst. Nat.* 2. 720. 32.

*Cimex Ypsilon aeneus* ouatus griseus, thorace spinoso, punctis binis albis, elytris fusco aeneis, puncto albo, scutello ypsilon flauesciente inscripto. *Degeer Inf.* 3. 332. 6. tab. 34. fig. 7. 8.

Habitat in America meridionali.

*clypeatus*. 50. C. thorace spinoso, viridis, fascia flauicante, capite clypeato. *Syst. Ent.* 703. 35. *Linn. Syst. Nat.* 2. 720. 33.

*Cimex*

*Cimex* ouatus, thorace vtrinque aculeato, fasciâ transuersali flauicante, scutello gibboso, capite clypeato, ano ferrato. *Gronov. Zooph.* 691.

Habitat in China.

51. *C. thorace dentato spinoso supra viridis, capite thoracisque antico scutellique basi flauis. albicollis.*

*Cimex flauicollis. Drury Inf.* 2. tab. 36. fig. 4.

Habitat in Iamaica.

Corpus subtus flaescens.

52. *C. thorace obtuse spinoso, subvirescens, antennis nigris, sterno porrecto. Syst. Ent.* 703. 36. *Liann. Syst. rhoidalis. Nat.* 2. 720. 35. *Fn. Sv.* 925.

*Cimex subouatus viridis, angulis thoracis acutis rubris, apice nigris, abdomine subtus acuto. Geoff. Inf.* 1. 465. 63.

*Cimex ouatus viridis, subtus pallide flauus, elytris macula fusca, abdomine supra rubro maculis nigris. Degeer Inf.* 3. 254. 3. tab. 35. fig. 7.

*Sulz. Hist. Inf.* tab. 11. fig. 72.

*Fuesly Inf. Helv.* 25. 483.

*Schaeff. Icon.* tab. 57. fig. 8.?

Habitat in Europae graminosis.

53. *C. thorace spinoso griseus, scutello punctis duobus flauis, tarsis anticis apice compresso membranaceis. oculatus. Syst. Ent.* 703. 37. \*

Habitat in China. *Dom. Drury.*

54. *C. thorace spinoso griseus, tibiis albo annulatis. Syst. annulatus. Ent.* 704. 38. \*

Habitat in Virginia. *Dom. Drury.*

55. *C. thorace crenato obtuse spinoso scutelloque punctis apusulatus. duobus rufis. Syst. Ent.* 704. 39. \*

Habitat in America. *Dom. Drury.*

56. *C. thorace obtuse spinoso, virescens, thorace maculis quatuor, scutello elytrisque apice fuscis. Syst. Ent. maculatus. 704. 40. \**

Habitat in America. *Dom. v. Rohr.*

Variat rarius elytrorum apice concolore.

57. *C.*



*pugnax.* 57. *C. oblongus*, thorace acute spinoso, virescens antennis rufis. *Syst. Ent.* 704. 41. \*

Habitat in America. *Dom. Drury.*

*emeritus.* 58. *C. thorace* acute spinoso, virescens, abdomine lineis duabus albis. *Syst. Ent.* 705. 42. \*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

*gladiator.* 59. *C. thorace* acute spinoso scutelloque flavescentibus nigro punctatis, elytris punctis albis scabris. *Syst. Ent.* 705. 43. \*

Habitat in America. *Dom. v. Rohr.*

Simillimum e Cap. bon. sp. misit *Dom. Vahl.* vix diuersum.

*furcatus.* 60. *C. thorace* acute spinoso ferrato, fuscus, clypeo acuminato bifido. *Syst. Ent.* 705. 44. \*

Habitat in Patagonia. *Mus. Dom. Banks.*

Variat colore toto griseo.

*pugillator.* 61. *C. thorace* acute spinoso, fuscus, margine flauo, subtus fuluus nigro punctatus.

Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*

Magnus. Caput deest. Thorax fuscus margine spinisque acutissimis flauis. Scutellum et elytra fusca, elytrorum margine flavescente. Alae atrae. Subtus fuluus lineis sex punctorum nigrorum.

*Vittor.* 62. *C. thorace* acute spinoso fuscus, scutello apice rufo, pedibus pallidis nigro punctatis. *Syst. Ent.* 705. 45. \*

Habitat in Americae Insulis. *Dom. v. Rohr.* In Brasilia *Mus. Dom. Banks.*

*ciliatus.* 63. *C. thorace* ciliato obtuse spinoso nigro, margine fasciaque postica flauis, femoribus posticis ferratis. *Syst. Ent.* 706. 46. \*

Habitat in America. *Dom. v. Rohr.*

*armiger.* 64. *C. thorace* acute spinoso griseus, scutello punctis duobus, antennis pedibusque pallidis.

Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*

Corpus supra griseum s. flavesces punctis numerosissimis nigris impressis. Thoracis spinac acutissimae, nigrae.

nigrae. Scutellum punctis duobus bascos pallidis. Abdomen pallidum vittis duabus fuscis. Antennae et pedes pallidi.

65. C. thorace obtuse spinoso, fuscis, pedibus pallidis nigro punctatis. *Syst. Ent.* 706. 47. \* *scorbuticus.*  
Habitat in America.

66. C. thorace acute spinoso obscure ferrugineus, spinis atris, abdomine atro, vittis flauescentibus. *melacanthus.*

Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*

Paruus. Corpus totum supra obscure ferrugineum, solis thoracis spinis acutissimis atris. Abdomen atrum vittis quatuor flauescentibus exterioribus marginalibus. Pedes flauescentes.

67. C. thorace acute spinoso, ano quadridentato, corpore flauescente, spinis subconcoloribus. *Syst. Ent.* 706. 48. \* *aggressor.*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

68. C. thorace subspinoso, virescens, elytris vitta submarginali flaua. *vittatus.*

Habitat ad Promontorium bon. sp. *Mus. Dom. Banks.*

Antennae ferrugineae. Caput virescens. Thorax virescens, punctatus margine antico flauescente. Scutellum punctis nigris impressum basi obscurum apice virescens. Elytra virescentia punctis numerosis nigris impressis vitta ante marginem flaua, quae intus puncto eiusdem coloris notata. Alae albae. Subtus virescens ano obtusissimo.

69. C. thorace acute spinoso, ano quadridentato, corpore viridi, spinis nigris. *Syst. Ent.* 706. 49. \* *defensor.*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

70. C. thorace obtuse spinoso, abdomine marginato acuto, antennis rufis. *Syst. Ent.* 707. 50. *marginatus.*

*Cimex marginatus* oblongo ouatus, griseus, thorace obtuse spinoso, antennis medio rubris. *Linn. Syst. Nat.* 2. 719. 28. *Fn. Sv.* 923.

*Cimex* oblongus fuscus immaculatus, thorace vtrinque obtuse angulato, capite prope antennis externe denticulato. *Geoff. Inf.* 1. 446. 20.

*Cimex auriculatus* oblongo ovatus fuscus, thorace obtuse angulato, antennis grossis clauatis medio rubris, abdomine supra rubro. *Degeer Inf.* 3. 272. 17.

*Scop. carn.* 363.

*Fuesl. Inf. Helv.* 25. 484.

*Schaeff. Icon.* tab. 41. fig. 4. 5.

Habitat in Europae plantis.

*spiniger*. 71. *C.* thorace obtuse spinoso dentatoque, capite quadri-spinoso.

Habitat in Italia. *D. Allioni.*

Statura omnino *C. marginati*, at duplo minor. Antennae rufae primo articulo apice spinoso cinereo, ultimo nigro. Caput cinereum antice spinis vtrinque duabus erectis, acutis. Thorax griseo fuscus margine eleuato in spinam obtusam postice emarginatam vel potius bidentatam. Scutellum griseum apice puncto albo. Elytra grisea basi albida. Alae fuscae. Abdomen griseum margine prominente acuto albo maculato.

*fasciculatus*. 72. *C.* thorace subspinoso, cinereus, alis fuscis, pedibus fasciculato tuberculatis.

Habitat ad Promontorium bon. sp.

Corpus duplo minus *C. marginato*, totum griseum alis solis fuscis. Pedes omnes, imprimis tamen quatuor anteriores, fasciculis plurimis pilorum tuberculatis.

*quadratus*. 73. *C.* thorace obtuse spinoso, supra fuscus, subtus flauescens, abdomine quadrato. *Syst. Ent.* 707. 51.

Habitat in Germania.

*velut*. 74. *C.* thorace acute spinoso bipunctato, elytris puncto basico strigaeque flauis. *Syst. Ent.* 707. 52. \*

Habitat in America. *Dom. Drury.*

*abundans*. 75. *C.* thorace subspinoso, elytris fuscis albo reticulatis, abdominis basi spinoso. *Syst. Ent.* 707. 53. \*

*Merian. Surin.* tab. 51.

Habitat Surinami. *Dom. Drury.*

76. C. thorace spinoso, femoribus posticis incurvis tibiisque posticis videntatis. *Syst. Ent.* 708. 54. *Linn.*  
*Syst. Nat.* 2. 720. 36. *Mus. Lud. Vlr.* 171.  
 Habitat ad Cap. bon. sp.
77. C. thorace ferrulato, femoribus posticis incurvis dentatis tibiisque posticis compressis. *Syst. Ent.* 708. 55.\* *femoratus.*  
 Habitat in India. *Mus. Dom. Banks.*
78. C. thorace acute spinoso, femoribus omnibus apice bidentatis, posticis incurvis. *curvipes.*  
 Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*  
 Statura omnino praecedentium. Corpus fuscum abdominis medio pallidiore. Antennae nigrae ultimo articulo rufo. Thorax fuscus, acute spinosus. Elytra obscure rufescentia alis nigris. Femora omnia apice bidentata. Tibiae inermes.
79. C. thorace acute spinoso nigro, fascia flava, tibiis posticis membranaceis dentatis. *membranaceus.*  
 Habitat in Insula St. Iago promontorii viridis. *Mus. Dom. Banks.*  
 Magnus. Antennae nigro rufoque variae. Caput nigrum puncto paruo antico lineolisque duabus baseos ferrugineis, subtus vtrinque vitta pallida. Thorax niger fascia antica flava margine ferrulato acuteque spinoso. Scutellum nigrum puncto apicis flauo. Elytra nigra puncto medio flauo. Alae immacolatae. Corpus subtus nigrum punctis numerosis rubris. Pedes nigri femoribus ferratis. Tibiae posticae vtrinque membrana, exteriori maiore bidentata, interiori puncto flauo auctae.
80. C. thorace spinoso immaculato, nigricans, elytris fascia albicante, tibiis posticis membranaceis dentatis. *phyllopus.*  
*Syst. Ent.* 708. 56. *Linn. Syst. Nat.* 2. 731. 113.  
 Habitat in America australi.
81. C. thorace subspinoso nigro, maculis duabus fultis, elytris fascia flava, tibiis posticis membranaceis dentatis. *auctus.*  
 Habitat Caiennae. *Mus. Dom. Banks.*

Nimis praecedenti affinis. Antennae nigro rufoque variae. Thorax ater maculis duabus magnis fulvis nigro punctatis. Elytra fuluo striata fasciaeque flavescente.

*Gonagra*. 82. C. thorace subspinoso, fascia alba, elytris fuscis, puncto cinereo. *Syst. Ent.* 708. 57. \*

Habitat in Insula St. Thomas Americae meridionalis. *Mus. Tottianum.*

*Kermesi-*  
*nus*. 83. C. thorace subspinoso, oblongus rufus, fascia albidus nigro punctata, femoribus posticis multidentatis. *Linn. Syst. Nat.* 2. 732. 116.

*Cimex rubro balteatus* oblongus obscure ruber, antennis nigris, elytris fascia flavescente nigro punctata, femoribus posticis magnis multidentatis. *Degeer Inf.* 3. 337. 11. tab. 34. fig. 15.

Habitat Surinami.

*australis*. 84. C. thorace subspinoso, oblongus niger, thorace fascia antica rubra, tibiis posticis basi membranaceis. *Syst. Ent.* 708. 58. \*

Habitat in Insula Otaheite maris pacifici. *Mus. Dom. Banks.*

*cruciger*. 85. C. thorace acute spinoso, oblongus supra niger, thorace lineis, elytris cruce ferrugineis. *Syst. Ent.* 709. 59. \*

Habitat in Brasilia.

*sanctus*. 86. C. thorace acute spinoso, oblongus rufus, elytris nigris, cruce ferruginea. *Syst. Ent.* 709. 60. \*

Habitat in Brasilia. *Mus. Dom. Banks.*

*Variat* scutello elytrisque rufis immaculatis, an distinctus?

*balteatus*. 87. C. thorace subspinoso, oblongus ferrugineus, elytris linea transversa flava, femoribus posticis multidentatis.

*Cimex balteatus* oblongus ferrugineus, elytris linea transversa flava, femoribus posticis multidentatis. *Linn. Mant.* 534.

*Cimex balteatus*. *Drury Inf.* 1. tab. 43. fig. 3.

Habitat in America meridionali.

88. C. thorace acute spinoso, oblongus rufus, femoribus *ferripes*.  
posticis dentatis. *Syst. Ent.* 709. 61. \*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

89. C. thorace acute spinoso, elongatus fuscus, femoribus *linearis*.  
posticis elongatis denticulatis. *Syst. Ent.* 710. 62. \*

Habitat in China. *Drury.* In Italia. *D. Allioni.*

90. C. thorace dentato rufo, macula nigra, femoribus *crenulatus*.  
posticis dentatis, corpore atro.

*Cimex pictus.* *Drury Inf.* 1. tab. 45. fig. 1.

Habitat in Americae meridionalis insulis.

Caput nigrum lineolis rubris. Thorax rufus macula magna antica margineque postico nigris. Margo dentatus. Corpus atrum, immaculatum abdomine solo rubro punctato.

91. C. elongatus ruber, thorace quadrispinoso. *Syst. Ent.* 4 *spinosus*.  
710. 63. *Linn. Syst. Nat.* 2. 720. 37.

*Cimex oblongus* ruber rostro arcuato, capite bispinoso, thorace quadrispinoso, abdomine spinis ciliato. *Degeer Inf.* 3. 347. 22. tab. 35. fig. 7.

Habitat in America meridionali.

\*\*\* Ouati, thorace mutico.

92. C. ouatus, supra sanguineus, scutello maculis duabus, *nigripes*.  
elytris vnica nigris. *Syst. Ent.* 710. 64. \*

*Cimex incarnatus.* *Drury Inf.* 2. tab. 36. fig. 5.

Habitat in India orientali. *D. Koenig.*

*Variat* rarius colore dilutiore et scutello immaculato.

93. C. ouatus viridis, capite thoraceque antice flauescen- *torquatus*.  
tibus. *Syst. Ent.* 710. 65. \*

Habitat in India, in Italia. *D. Allioni.*

94. C. viridi aeneus, thorace fascia antica sanguinea, scu- *rutilans*.  
tello elytris antice flauescen-  
tibus.

Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*

Magnus. Caput aeneum, nitidum antennis nigris. Thorax aeneus nitidus fascia antica sanguinea, margine tamen ipso eleuato, aeneo. Scutellum aeneum apice

*Fabr. Sp. Inf. T. II.*

354 CLASS. VII. RYNGOTA. Cimex.

late flavo. Elytra aenea margine interiori sanguineo apiceque terminantur puncto flavescente. Alae aeneae. Corpus subtus flavescentes abdomine punctis lateralibus cyaneis. Pedes nigri.

*viridulus*. 95. C. ouatus supra flavus, punctis viridibus, subtus viridis. *Syst. Ent.* 710. 66. *Linn. Syst. Nat.* 2. 721. 39. *Mus. Lud. Vlr.* 172.  
Habitat in Indiis.

*prasinus*. 96. C. ouatus viridis immaculatus, antennis apice albo fuscoque annulatis. *Syst. Ent.* 711. 67.

*Cimex prasinus* subrotundus totus viridis, scutello concolore. *Linn. Syst. Nat.* 2. 722. 49. *Fn. Sv.* 931.

*Cimex viridis* totus ouatus viridis, subtus viridigriseus, pedibus concoloribus. *Degeer Inf.* 3. 266. 9.

*Fuesly Inf. Helv.* 25. 488.

Habitat in Europae sylvis.

Simillimum e Cap. bon. sp. a *Dom. Vahl* accepti vix distinctum.

*diffimilis*. 97. C. ouatus supra viridis, subtus ferrugineus,

Habitat in Germania. *Dom. de Hattorf.*

Nimis praecedenti affinis videtur, at subtus totus ferrugineus pedibus virescentibus. An praecedentis mera varietas aetatis?

*juni-peri-nus*. 98. C. subrotundus viridis margine vndique scutelloque apice flavo. *Syst. Ent.* 711. 68. *Linn. Syst. Nat.* 2. 722. 48. *Fn. Sv.* 930.

*Cimex* subrotundus viridis. *Geoff. Inf.* 1. 464. 61.

*Cimex* sylvestris viridis. *Raj. Inf.* 53. 1.

*Schaeff. Icon.* tab. 46. fig. 1. 2.

*Fuesly Inf. Helv.* 25. 487.

*Degeer Inf.* 3. 231. tab. 13. fig. 1. 2.

Habitat in Europae Juniperetis.

*smaragdulus*. 99. C. ouatus viridis, scutello punctis tribus bascos flavis. *Syst. Ent.* 711. 69. \*

Habitat in Insula Madera. *Mus. Dom. Banks.*

100. *C. ouatus* fusco griseus, scutello punctis tribus apice coelebs.  
ceque flavescentibus. *Syst. Ent.* 711. 70. \*  
Habitat in noua Hollandia. *Mus. Dom. Banks.*
101. *C. subrotundus* griseus, margine vndique sanguineo. *dumofus.*  
*Syst. Ent.* 711. 71. \* *Linn. Syst. Nat.* 2. 721. 46. *Fn.*  
*Sv.* 929.  
Habitat in Europa, Suecia.
102. *C. rotundatus* flavescent, capite lineis, scutello punctis tripunctatis tribus nigris. *Syst. Ent.* 711. 72. \*  
Habitat in America. *Dom. Mauduit.*
103. *C. ouatus* griseo fuscoque varius, alis albis fusco punctatis. *Syst. Ent.* 772. 73. *atomarius.*  
Habitat in Insula Terre neuve. *Mus. Dom. Banks.*
104. *C. ouatus* griseo nigroque varius, alis albis nigro striatis. *Syst. Ent.* 712. 74. \* *nubilus.*  
Habitat ad Cap. bon. sp. *Mus. Dom. Banks.*
105. *C. rotundatus* luteo nigroque varius, thorace luteo, punctatis sex nigris. *Syst. Ent.* 712. 75. *Linn. Syst. Nat.* 2. 721. 42. *Mus. Lud. Vlr.* 175. *6punctatus.*  
Habitat in Indiis.
106. *C. ouatus* griseus, scutello vitta nigra. *Syst. Ent.* 712. 76. \* *signatus.*  
Habitat in Sierra Leon Africae. *Mus. Dom. Banks.*
107. *C. ouatus* fuscus, scutello punctis duobus flauis, elytris puncto nigro. *grauis.*  
Habitat in noua Zelandia. *Mus. Dom. Banks.*  
Medius. Corpus totum supra fuscum, si potius deulo armato visum, ferrugineum punctis numerosissimis impressis nigris. Caput acutum, prominens, sulcis duobus impressis. Thoracis margo tenuissime flavescent. Scutellum puncto vtrinque bascos apiceque flavescentibus. Elytra margine bascos flavescente punctoque medio magno nigro. Alae striatae, nigrae. Corpus subtus flavescent femoribus posticis apice nigris.



- cinctus.* 108. *C. ovatus viridis*, thoracis abdominisque margine sanguineo. *Syst. Ent.* 712. 77.\*  
Habitat in Oriente. *Prof. Forskahl.*
- eruentus.* 109. *C. ovatus viridis*, thoracis abdominisque marginibus, antennis pedibusque sanguineis. *Syst. Ent.* 713. 78.\*  
Habitat Surinami. *Dom. Drury.*
- griseus.* 110. *C. ovatus griseus*, abdominis lateribus albo nigroque variis, ventre antice spinoso.  
*Cimex griseus* abdominis lateribus albo nigroque variis, alis nebulosis, ventre antice spinoso. *Linn. Syst. Nat.* 2. 721. 43. *Fn. Sv.* 926.  
*Cimex Betulae* ovatus viridis siue rubro griseus, scutello macula nigra, antennis griseis apice nigris abdominis lateribus flavo nigroque variis. *Degeer Inf.* 3. 261. 8. tab. 14. fig. 9.  
Habitat in Europa.  
Antennae nigrae, flavo annulatae. Corpus supra obscure griseum subtus pallidius punctis fuscis irroratum. Alae albae, fusco punctatae.
- interstin-*  
*ctus.* 111. *C. ovatus griseus*, abdominis margine nigro maculato. *Syst. Ent.* 713. 79. *Linn. Syst. Nat.* 2. 721. 44. *Fn. Sv.* 927.  
Habitat in Europae cultis.
- papillo-*  
*sus.* 112. *C. ovatus fuscus*, antennis nigris, sterno gibbo compresso. *Syst. Ent.* 713. 80.\*  
*Cimex papillosus.* *Drury Inf.* 1. tab. 43. fig. 2.  
Habitat in Sierra Leon Africae.
- funebis.* 113. *C. ovatus ater*, antennis alis pedibusque concoloribus. *Syst. Ent.* 713. 81.\*  
Habitat in Sierra Leon Africae.
- ruficornis.* 114. *C. rotundatus ater*, antennis rufis. *Syst. Ent.* 713. 82.\*  
Habitat in India.
- Baccarum.* 115. *C. ovatus subfulvus*, abdominis margine fusco maculato. *Syst. Ent.* 713. 83. *Linn. Syst. Nat.* 2. 721. 45. *Fn. Sv.* 928.

*Cimex fuscus* antennis abdominisque margine nigro croceoque variegatis. *Geoff. Inf.* 1. 466. 64.

*Cimex Verbaschi* ovatus griseo fuscus, antennis albo nigroque maculatis, scutelli apice glauco, abdomine supra nigro, margine flauo maculato. *Degeer Inf.* 3. 257. 4. tab. 14. fig. 5.

*Cimex* scapulis magis exstantibus, macula e flauo rubente in centro crucis dorsalis. *Raj. Inf.* 54. 2.

*Cimex* ex luteo virescenti infuscatus ad alui margines nigris maculis. *List. Loqu.* 396. *mut.* tab. 31. fig. 19. *Ionst. Inf.* tab. 17. fig. 9.

*Schaeff. Icon.* tab. 57. fig. 1. 2.

*Fuesly Inf. Helv.* 25. 486.

Habitat in Europae baccis frequens.

116. *C. ovatus fuscus*, thorace scutelloque punctatis, abdominis punctis genibusque flauis. *cribrum.*

Habitat in America. *D. Schulz.*

Magnus. Caput fuscum antennis nigris. Thorax fuscus punctis impressis. Scutellum punctis impressis. Elytra laeuia, fusca puncto medio flauo. Alae nigrae. Abdomen fuscum margine punctisque flauis. Pedes nigri genibus flauis.

117. *C. ovatus glaucus*, elytris apice puncto nigro. *Syst. glaucus.* *Ent.* 714. 84. \*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

118. *C. rotundatus* supra sanguineus, capite scutelli basi alisque atris. *Syst. Ent.* 714. 85. \*

*Schroet. Abhandl.* 1. tab. 2. fig. 4.

Habitat in America. *Mus. Dom. Banks.*

119. *C. ovatus nigro rubroque varius*, capite alisque nigris. *Syst. Ent.* 714. 86. *Linn. Syst. Nat.* 2. 723. 56. *Fn. Sv.* 937.

*Geoff. Inf.* 1. 469. 69.

*Scop. carn.* 361.

*Sulz. Inf.* tab. 11. fig. 73.

*Schaeff. Icon.* tab. 60. fig. 10.

*Fuesly Inf. Helv.* 26. 495.

Habitat in Europa, in Oriente. *Prof. Forskahl.*

358 CLASS. VII. RYNGOTA. Cimex.

*festivus.* 120. *C. ouatus* nigro rubroque varius, thorace punctis sex nigris, alis fuscis, margine albedo. *Syst. Ent.* 714. 87. *Linn. Syst. Nat.* 2. 723. 57.  
*Cimex dominulus.* *Scop. Carn.* 362.  
*Fuesly Inf. Helv.* 26. 490.  
 Habitat in America boreali.

*rutilus.* 121. *C. ouatus*, thorace abdomineque rufis, thorace maculis quatuor, abdomine fasciis nigris. *Syst. Ent.* 714. 88. \*  
 Habitat in noua Hollandia.

*cruciatus.* 122. *C. ouatus* nigro pallidoque varius, scutello nigro, cruce alba. *Syst. Ent.* 714. 89. \*  
 Habitat in India orientali. *D. Koenig.*

*bioculatus.* 123. *C. rotundatus* supra niger, thorace rufo, punctis duobus nigris, scutello margine rufo. *Syst. Ent.* 714. 90. \*  
 Habitat in America. *Dom. Drury.*

*apunctatus.* 124. *C. ouatus* obscure rufus, scutello punctis duobus, apiceque albis, abdominis margine nigro punctato.  
 Habitat in Italia. *D. Allioni.*  
 Magnitudo *C. oleracei*. Caput fuscum antennis basi pallidis apice nigris. Thorax et elytra obscure rufa. Scutellum concolor punctis duobus elevatis ad basin apiceque albo. Corpus subtus griseum margine abdominis tenuissime albo nigro punctato. Pedes pallidi.

*bicolor.* 125. *C. ouatus* niger, elytris nigro alboque variis, alis albis. *Syst. Ent.* 715. 91. *Linn. Syst. Nat.* 2. 722. 55. *Fn. Sv.* 936.  
*Cimex* ouatus niger, thoracis lateribus elytrorumque maculis quatuor albis. *Geoff. Inf.* 1. 470. 73.  
*Cimex* ouatus niger nitidus, thorace elytris tibiisque spinosis albo maculatis. *Degeer Inf.* 3. 268. 12.  
*Cimex* colore nigro splendente, maculis albis picto. *Raj. Inf.* 54. 5.  
*Cimex* niger maculis candidis notatus. *List. Scar.* 396. 38.

*Cimex*

*Cimex niger albo maculatus.* *Petiv. Gazoph. tab. 14.*  
fig. 7.

*Schaeff. Icon. tab. 41. fig. 8. 9.*

*Fuesly Inf. Helv. 26. 494.*

Habitat in Europae hortis.

126. *C. ouatus coeruleo aeneus, thorace lineola, scutello olivaceus.*  
apice elytris puncto albo rubrouc. *Syst. Ent. 715.*

92. *Linn. Syst. Nat. 2. 722. 53. Fn. Sv. 934.*

*Cimex sylvestris coerulefens. Raj. Inf. 54. 6. 7.*

*Geoff. Inf. 1. 471. 74.*

*Degeer Inf. 3. 266. 10. tab. 15. fig. 22.*

*Schaeff. Icon. tab. 46. fig. 4. 5.*

*Fuesly Inf. Helv. 25. 492.*

Habitat in Europae Tetradyamiis.

127. *C. subrotundus supra ater, thorace lineis tribus, scu- pilus.*  
tello linea punctisque quatuor, elytris margine fer-  
rugineis. *Syst. Ent. 715. 93.\**

Habitat in India orientali. *D. Koenig.*

128. *C. ouatus niger, margine vndique albo, elytris pun- bigutta-*  
cto albo. *Syst. Ent. 716. 94. Linn. Syst. Nat. 2. 722. tus.*

54. *Fn. Sv. 935.*

*Fuesly Inf. Helv. 25. 493.*

*Scop. carn. 356.*

Habitat in Europae plantis.

129. *C. ouatus coeruleus immaculatus. Syst. Ent. 716. 95. coeruleus.*  
*Linn. Syst. Nat. 2. 722. 50. Fn. Sv. 933.*

*Cimex ouatus aeneo coeruleus nitidus totus. Degeer*

*Inf. 3. 268. 11.*

*Geoff. Inf. 1. 472. 75.*

*Fuesly Inf. Helv. 25. 489.*

*Schaeff. Icon. tab. 51. fig. 4.*

Habitat in Europae Thefio.

130. *C. ouatus ater, plantis rufis. Syst. Ent. 716. 96. Linn. morio.*  
*Syst. Nat. 2. 722. 51. Fn. Sv. 932.*

*Sulz. Inf. tab. 11. fig. 77.*

*Schaeff. Icon. tab. 57. fig. 11.*

— — — tab. 82. fig. 6.

360 CLASS. VII. RYNGOTA. Cimex.

*Fuesly Inf. Helv. 25. 491.*

Habitat in Europae plantis oleraceis.

*tristis.* 131. *C. ouatus ater*, clypeo orbiculato, thorace retuso.  
*Syst. Ent. 716. 97.\**

*Cimex ouatus* totus niger, alis pallidis. *Geoff. Inf. 1. 470. 70.*

*Cimex niger spinipes* ouatus niger nitidus, tibiis spinosis, elytrorum parte membranacea fusca, alis albidis. *Degeer Inf. 3. 269. 13.*

Habitat in Europae plantis.

*spinipes.* 132. *C. ouatus ater*, pedibus piceis, tibiis spinosissimis.  
Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*

Corpus magnitudine et statura omnino praecedentium, atrum, nitidum. Alae pallefcentes. Antennae et pedes picea tibiis omnibus spinis numerosissimis acutis cinctis.

*lugens.* 133. *C. ouatus fuscus*, thoracis lineola, scutello abdomineque margine albis. *Syst. Ent. 716. 98.\**

Habitat in America. *Dom. Drury.*

*melanoccephalus.* 134. *C. subrotundus griseus*, capite scutellique basi nigro aeneis. *Syst. Ent. 716. 99.\**

Habitat in Anglia, in Italia. *Mus. Dom. Banks.*

*acuminatus.* 135. *C. ovalis antice attenuatus exalbidus fusco striatus*, antennis apice rufis. *Syst. Ent. 717. 100. Linn. Syst. Nat. 2. 723. 59. Fn. Sv. 939.*

*Cimex rostratus* ouatus griseus punctatus, fasciis longitudinalibus fuscis, capite magno triangulari longitudine thoracis. *Degeer Inf. 3. 271. 16. tab. 14. fig. 12. 13.*

*Musca cimiciformis.* *Raj. Inf. 56. 6.*

*Geoff. Inf. 1. 472. 77.*

*List. mut. tab. 31. fig. 20.*

*Schaeff. Icon. tab. 42. fig. 11.*

*Fuesly Inf. Helv. 26. 498.*

Habitat in Europae plantis.

\*\*\* Oblor.



\*\*\*\* Oblongi.

136. *C. oblongus* fuscus, femoribus posticis arcuato clauatis videntatis, abdominis basi subtus gibbo. *Phasianus.*  
 Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*  
 Corpus magnum, fuscum. Antennae nigrae articulo ultimo basi apiceque ferrugineo. Alae nitidae. Abdomen subtus basi gibbum. Femora quatuor anteriora apice videntata, postica maxima arcuata, eleuata dente valido compresso. Tibiae compressae lineae vtrinque eleuata vnica.
137. *C. oblongus* fuscus, femoribus posticis arcuatis dentatis, abdomine quadrispinoso. *bellicosus.*  
 Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*  
 Statura omnino *C. valgi*, at duplo minor et thorax inermis. Antennae nigrae. Caput et thorax fusca, pube aurea parum nitida. Scutellum apice cinerascit. Elytra fusca alis nigris. Abdomen subtus pube aurea basi vtrinque spinis duabus atris, validis. Pedes quatuor antici pallidi femoribus apice videntatis, postici obscuriores femoribus breuibus arcuatis apice dentatis. Tibiae compressae serrulatae.
138. *C. oblongus*, elytris fuscis flauo striatis, femoribus tibisque posticis vnispinosis. *Gallus.*  
 Habitat Surinami. *Mus. Dom. Banks.*  
 Statura et magnitudo *C. Indi*. Caput flauo nigroque varium antennis rostroque nigris. Thorax inermis virescens punctis numerosissimis impressis nigris, antice flauo nigroque varius. Scutellum nigrum margine maculisque duabus magnis flauescens. Elytra nigra striis quatuor flauescens, quae apice striga transuersa cinguntur. Alae fuscae. Corpus subtus atrum abdomine sanguineo, ano atro. Pedes antici quatuor simplices atri, apice pallidi. Femora postica recta, crassa, atra spina arcuata flaua in medio. Tibiae flauae basi atrae spina recta acutissima.
139. *C. oblongus* niger subtus rubro punctatus, femoribus dentatis. *Syst. Ent. 717. 101.\* Harpagon.*  
 Habitat in America. *Dom. Drury.*

Medius. Caput pallescens antennis rufis articulis ultimis apice nigris. Thorax fuscus margine tenuissime pallido. Elytra fusca margine exteriori pallido. Alae nigrae. Abdomen supra rufescens ano nigro. Corpus subtus pallidum.

*cingulatus*. 150. C. oblongus, thorace testaceo, antice nigro, margine albo, elytris puncto nigro. *Syst. Ent.* 719. 108. \*  
Habitat in noua Hollandia. *Mus. Dom. Banks.*

*superficialis*. 151. C. oblongus rufus, antennis thoracis basi fascia elytrorum alisque nigris. *Syst. Ent.* 719. 109. \*  
Habitat in America. *Mus. Dom. Banks.*

*albofasciatus*. 152. C. oblongus rufus, elytris nigris, fascia alba. *Cimex albofasciatus* oblongus ruber, elytris nigris, fascia alba, femoribus anticis dentatis. *Degeer luf.* 3. 335. 9. tab. 34. fig. 11.  
Habitat Surinami.  
Margo thoracis et elytrorum albidus.

*ruficollis*. 153. C. oblongus glaucus, capite atro, thorace antice rufo. *Syst. Ent.* 719. 110. \*  
Habitat in Brasilia. *Mus. Dom. Banks.*

*scabrosus*. 154. C. oblongus niger, thoracis margine elytrorumque fasciis duabus rufescentibus. *Syst. Ent.* 719. 111. \*  
Habitat in America. *Dom. Drury.*

*leprosus*. 155. C. oblongus, thorace nigro, basi margine punctisque duobus rubris, elytris atris immaculatis. *Syst. Ent.* 719. 112. \*  
Habitat in America.

*Koenigii*. 156. C. oblongus testaceus, elytris puncto nigro, alis atris. *Syst. Ent.* 719. 113. \*  
Habitat Tranquebariae. *D. Koenig.*

*aegyptius*. 157. C. oblongus rubro nigroque varius, elytris rubris puncto nigro. *Syst. Ent.* 719. 114. *Linn. Syst. Nat.* 2. 727. 79. *Mus. Lud. Vlr.* 178.  
Habitat in Aegypto.



158. *C. oblongus* rubro fuscoque varius, alis atris, apice *mendicus*.  
lineolaque baseos albis. *Syst. Ent.* 720. 115.\*  
Habitat in Malabaria. *D. Koenig.*
159. *C. oblongus* supra ater, thorace elytrisque punctis *famelicus*.  
duobus flavescentibus, alis lineola baseos alba.  
Habitat in Promontorio bon. sp. *Mus. Dom. Banks.*  
Medius. Caput atrum punctis duobus anticis lineola-  
que baseos sanguineis. Thorax ater margine anti-  
co sanguineo punctisque duobus magnis flavis. Ely-  
tra nigra, basi apiceque flava. Alae nigrae lineola  
transversa baseos alba. Pectus nigrum fasciis latera-  
libus fulvis. Abdomen nigrum basi fasciisque late-  
ralibus fulvis. Pedes nigri.
160. *C. oblongus*, elytris rufis, fascia nigra, alis punctis *4gutta-*  
duobus albis. *Syst. Ent.* 720. 116.\* *tus.*  
Habitat in noua Hollandia. *Mus. Dom. Banks.*
161. *C. oblongus* niger, elytris rufis, puncto medio atro, *punctato-*  
alis atris, punctis duobus albis. *guttatus.*  
Habitat in Italia. *D. Allioni.*  
Paruus. Caput nigrum, immaculatum. Thorax niger,  
antice rufus. Scutellum nigrum, immaculatum. Ely-  
tra laete rufa puncto medio marginis crassioris atro.  
Alae albae puncto medio apicisque albo. Corpus  
subtus nigrum. Pedes nigri.
162. *C. oblongus*, elytris rufis, striga vndata nigra, alis *bigutta-*  
nigris, puncto medio albo. *Syst. Ent.* 720. 117. *tus.*  
Habitat in noua Hollandia. *Mus. Dom. Banks.*
163. *C. oblongus* sanguineus, antennis, elytris pedibusque *sanguino-*  
fuscis. *Syst. Ent.* 721. 118.\* *lentus.*  
Habitat in Americae meridionalis insulis.
164. *C. oblongus* rufus, antennis alisque nigris, suturis *suturalis.*  
omnibus albis. *Syst. Ent.* 721. 119.\*  
Habitat in Americae meridionalis insulis. *Mus. Tot-*  
*tianum.*
165. *C. oblongus* rufus, antennis pectore ano pedibusque *excaua-*  
atris, scutello canaliculato. *tus.*  
Habitat



Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*  
 Magnus. Antennae atrae. Caput rufum macula oculari nigra. Thorax rufus maculis duabus anticis nigris, linea dorsali marginibusque elevatis. Scutellum valde canaliculatum rufum. Elytra rufa. Alae atrae. Pectus atrum. Abdomen rufum ano atro. Pedes nigri.

*Carnifex.* 168. *C. oblongus rufus, antennis tibiisque fuscis. Syst. Ent.* 721. 120.\*

Habitat in Cap. bon. sp. *Mus. Dom. Banks.*

*Augur.* 167. *C. oblongus rufus, antennis alis pedibusque nigris.*

Habitat in India orientali. *Mus. Dom. Banks.*

Affinis *C. carnifici*, a quo tamen differt colore subtus toto rufo immaculato pedibusque totis nigris.

*maffians.* 168. *C. oblongus supra rufus, thorace scutelloque punctis duobus nigris.*

Habitat in India orientali. *Mus. Dom. Banks.*

Medius. Caput pallide rufescens antennis nigris. Thorax rufus antice pallidior punctis duobus nigris. Scutellum rufum punctis duobus bascos nigris. Elytra rufa, alae nigrae. Subtus flauescens fasciis cyaneis.

*apterus.* 169. *C. oblongus rubro nigroque varius, elytrisque rubris, punctis duobus nigris, alis nullis. Syst. Ent.* 721. 121. *Linn. Syst. Nat.* 2. 727. 28.

*Cimex oblongus ruber, nigro maculatus, elytris abbreviatis, alis nullis. Degeer Inf.* 3. 276. 20.

*Geoff. Inf.* 1. 440. 11.

*Scop. carn.* 370.

*Raj. Inf.* 55. 3.

*Sulz. Hist. Inf.* tab. 10. fig. 14.

*Fueshy Inf. Helv.* 26. 504.

Habitat in Europae Malua sylvestri.

*friatulus.* 170. *C. oblongus flauescens, scutello elytris alisque fuscis, elytris apice cinereis. Syst. Ent.* 721. 122.\*

Habitat in Brasilia. *Mus. Bom. Banks.*

*nugax.* 171. *C. oblongus fuscus, abdominis margine maculato, tibis anticis femoribusque posticis basi pallidis.*

Habitat

Habitat in Italia. *D. Allioni.*

Corpus medium, fuscum. Antennae albo nigroque variae. Rostrum basi fuscum. Abdominis basis albo fuscoque variegata. Pedes fusci. Tibiae anticae pallidae apice fuscae. Femora postica pallida, apice fusca.

72. *C. oblongus* fuscus, abdomine supra sanguineo, femoribus posticis sexdentatis. *Syst. Ent.* 721. 123. *calcaratus*.

*Cimex calcaratus* oblongus niger, capite linea rubra, femoribus posticis subtus sexdentatis. *Linn. Syst. Nat.* 2. 732. 114. *Fn. Sv.* 968.

*Cimex oblongus* nigro fuscus, abdomine supra rubro, femoribus posticis subtus dentatis. *Degeer Inf.* 3. 280. 24. tab. 14. fig. 23. 24.

*Schaeff. Icon.* tab. 123. fig. 2. 3.

Habitat in Europa.

173. *C. oblongus* ater, elytris fuscis, macula rhombea atra. *Syst. Ent.* 722. 124. *Linn. Syst. Nat.* 2. 729. 96. *Fn. Sv.* 956. *Pini.*

*Cimex oblongus* niger, thorace postice cinereo, elytris cinereis, macula nigra alisque nigris. *Geoff. Inf.* 1. 449. 28.

*Cimex oblongus* niger, thorace postice elytrisque fuscis punctatis, macula nigra. *Degeer Inf.* 3. 279. 22. tab. 14. fig. 23.

*Schaeff. Icon.* tab. 42. fig. 12.

Habitat in Europae borealis pinetis frequens.

174. *C. oblongus* ater, elytris fuscis. *Syst. Ent.* 722. 125.\* *sylvaticus*.  
Habitat in Europae borealis nemoribus.

175. *C. oblongus* ater, alis macula rhombea flava. *Syst. Rolandri. Ent.* 722. 126. *Linn. Syst. Nat.* 2. 729. 98. *Fn. Sv.* 957.

*Cimex oblongus* atro fuscus, alarum macula flava. *Geoff. Inf.* 1. 459. 51.

*Cimex fulvo maculatus* oblongus niger, antennis setaceis, elytris macula fulva, pedibus fuscis nigro punctatis. *Degeer Inf.* 3. 294. 33.

*Sulz. Inf.* tab. 11. fig. 76.

*Schaeff.*

*Schaeff. Icon. tab. 87. 7.*

*Fuesly Inf. Helv. 26. 511.*

Habitat in Europae pinetis.

*Forsteri.* 176. *C. oblongus ater*, scutelli apice, thoracis elytrorumque margine sanguineis.

Habitat ad Cap. bon. sp. *Mus. Dom. Banks.*

Magnitudo et statura praecedentis. Caput atrum immaculatum. Thorax planus margine omni sanguineo, quorum posticus in medio dilatatus. Scutellum nigrum, apice sanguineum. Elytra nigra margine bascos sanguineo. Abdomen nigrum margine medioque sanguineis. Pedes nigri.

*clauimanus.* 177. *C. oblongus ater*, thoracis elytrorumque margine sanguineis, femoribus anticis incrassatis dentatis.

Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*

Statura et summa affinitas praecedentis. Caput atrum immaculatum. Thorax planus ater margine omni sanguineo. Scutellum apice vix rufescit. Elytra nigra margine omni sanguineo. Pedes nigri femoribus anticis incrassatis dentatis.

*6pustulatus.* 178. *C. oblongus niger*, thorace elytrisque maculis duabus rufis. *Syst. Ent. 722. 127.\**

Habitat in Iamaica. *Dom. Drury.*

*lynceus.* 179. *C. oblongus niger*, elytris griseis, apice macula nigra, puncto albo. *Syst. Ent. 722. 128.\**

Habitat in Angliae plantis.

*Vrticae.* 180. *C. oblongus niger*, elytris griseis, alis albis, puncto nigro. *Syst. Ent. 723. 129.\**

Habitat in Vrtica Angliae.

*sylvestris.* 181. *C. oblongus niger*, elytris fuscis apice nigro punctatis, alis nigris puncto bascos apicisque albo.

Habitat Hafniae. *Dom. Lund.*

Parvus. Statura praecedentium. Caput, thorax, scutellum, corpus, pedes atra, immaculata. Elytra fusca margine postico punctis aliquot atris. Alae nigrae puncto bascos apicisque distincto albo.

CLASS. VII. RYNGOTA. Cimex. 369

182. *C. oblongus*, elytris fuscis, basi punctisque duobus *podagri-*  
albis, femoribus anticis crassissimis bidentatis. *Syst. cus.*  
*Ent.* 723. 130. \*

*Cimex abietis* oblongiusculus maculato fulvus pedibus  
rufis, femoribus anticis crassioribus dentatis. *Linn.*

*Syst. Nat.* 2. 732. 115. *Fn. Sv.* 969. ?

*Degeer Inf.* 3. tab. 15. fig. 20. ?

Habitat in Anglia. *Mus. Dom. Banks.*

183. *C. oblongus* griseus, rubro punctatus, antennis claua- *crassicorn-*  
tis. *Syst. Ent.* 723. 131. *Linn. Syst. Nat.* 2. 729. 92. *nis.*  
*Fn. Sv.* 952.

*Schaeff. Icon.* tab. 13. fig. 10.

Habitat in Europa.

184. *C. oblongus* niger, elytris cinereis, alis albis. *Syst. arenari-*  
*Ent.* 723. 132. *Linn. Syst. Nat.* 2. 729. 95. *Fn. Sv.* *us.*  
955.

Habitat in Europa boreali frequens.

185. *C. oblongus*, thoracis elytrorumque margine femo- *succinctus.*  
rumque basi rufis. *Syst. Ent.* 723. 133. *Linn. Syst.*  
*Nat.* 2. 727. 82.

*Cimex rubrocinctus* oblongus cinereo niger, rubro  
marginatus, femoribus anticis dentatis, basi rubris.

*Degeer Inf.* 3. 339. 13. tab. 34. fig. 19.

Habitat in Americae borealis plantis.

186. *C. oblongus*, supra ferrugineo fuscus, corpore elytro- *tunicatus.*  
rumque margine flavis.

Habitat in Germania. *Dom. de Hattorf.*

Antennae rufae articulorum apicibus nigris. Caput et  
thorax ferrugineo fusca. Elytra concolora margine  
tenuissime flavo apicibusque rufis puncto nigro.  
Corpus et pedes flava, immaculata.

187. *C. oblongus* rufus, antennis pectore elytris tibiisque *pubicor-*  
nigris. *Syst. Ent.* 724. 134. \* *nis.*

Habitat in America. *Dom. Drury.*

188. *C. oblongus* niger, pedibus antennisque flavis. *Syst. Coryli.*  
*Ent.* 724. 135. *Linn. Syst. Nat.* 2. 733. 121. *Fn. Sv.* 974.

Habitat in Europae borealis Corylo.

*Fabr. Spec. Inf. T. II.*

A a

189. *C.*

*campestris*. 189. *C. oblongus* flauescens, elytris macula ferruginea. *Syst. Ent.* 724. 136. *Linn. Syst. Nat.* 2. 728. 87. *Fn. Sv.* 950.

*Geoff. Inf.* 1. 452. 34.

Habitat in Europae borealis campis.

*pratensis*. 190. *C. oblongus* flauescens, elytris viridibus. *Syst. Ent.* 724. 137.

*Cimex pratensis* oblongus griseus, scutello macula cordata flava, elytris apice puncto fusco. *Linn. Syst. Nat.* 2. 728. 86. *Fn. Sv.* 949.

*Geoff. Inf.* 1. 451. 33.

*Scop. carn.* 386.

Habitat in Europae pratis frequens.

*punctatus*. 191. *C. oblongus* viridis, thorace bipunctato, elytris pallidioribus, puncto apicis flavo. *It. Norw.* d. 9. Aug. \*

Habitat in Norwagiae plantis.

*leucocephalus*. 192. *C. oblongus* niger, capite pedibusque fulvis. *Syst. Ent.* 724. 138. *Linn. Syst. Nat.* 2. 723. 60. *Fn. Sv.* 940.

*Cimex* rotundato ovatus, capite genibusque ferrugineis, pedibus saltatoriis. *Geoff. Inf.* 1. 472. 76.

*Cimex* ovatus niger, antennis setaceis, capite pedibusque rufo flavis. *Degeer Inf.* 3. 290. 28.

Habitat in Europa.

*rosens*. 193. *C. oblongus* sanguineus, capite elytrorumque apicibus albidis, alis fuscis. *Gen. Inf. Mant.* 299. \*

Habitat in Germania. *D. Schulz.*

*tripustulatus*. 194. *C. oblongus* niger, scutello maculisque tribus elytrorum coccineis.

Habitat in racemis Vrticae dioicae, Hafniae frequens. *Dom. Lund.*

Parvus. Caput ferrugineum ore nigro. Thorax ater linea dorsali rufa antice tricuspidata. Scutellum coccineum. Elytra glabra nigra maculis tribus coccineis ad marginem crassiore, quarum vltima terminatur puncto atro. Corpus atrum. Pedes ferruginei.

195. *C. oblongus niger*, elytris alisque apice puncto nigro. *Syst. Ent.* 724. 139. \* *triguttatus*.

*Cimex triguttatus oblongus niger*, elytris testaceis, punctis tribus niueis, interiore minore. *Linn. Syst. Nat.* 2. 729. 94.

Habitat in Europa australiori.

196. *C. oblongus niger*, elytris striatis, alis postice flavo maculatis. *Syst. Ent.* 725. 140. *Linn. Syst. Nat.* 2. 729. 93. *Fn. Sv.* 954. *salto-rinus*.

Habitat in Europa ad littora maris.

197. *C. oblongus ater*, antennis apice capillaribus. *Syst. Ent.* 725. 141. *Linn. Syst. Nat.* 2. 725. 72. *Fn. Sv.* 944. *ater*.

*Cimex oblongus ater*, antennis seta terminatis. *Geoff. Inf.* 1. 460. 54.

Habitat in Europa boreali.

198. *C. oblongus ater*, rostro femoribusque sanguineis, *tyrannus*. antennis apice capillaribus.

Habitat in Italia. *D. Allioni*.

Affinis nimis praecedenti, at rostrum omnino sanguineum. Pedes rufi femoribus inprimis anticis nigro annulatis. Plantae nigrae.

199. *C. oblongus ater*, capite scutello maculisque duabus elytrorum coccineis, antennis apice capillaribus. *Schack.*

Habitat in Italia. *D. Allioni*.

Statura praecedentis. Caput coccineum antennis nigris. Thorax ater nitidus, immaculatus. Scutellum coccineum. Elytra atra macula magna transversali coccinea in medio, alia minori ad apicem. Pedes nigri.

200. *C. oblongus fuscus*, elytris fasciis duabus albis, antennis apice capillaribus. *Syst. Ent.* 725. 142. \* *bifasciatus*.

*Cimex clauatus oblongus niger*, elytris fuscis, fasciis duabus nigris. *Linn. Syst. Nat.* 2. 729. 97. ?

Habitat Lipsiae.

201. *C. oblongus niger*, capite thorace pedibusque rufis, *flavicollis*. antennis apice capillaribus. *Syst. Ent.* 725. 143. \*

Habitat in Anglia. *Mus. Dom. Banks.*

372 CLASS. VII. RYNGOTA. Cimex.

*trifasciatus*. 202. *C. oblongus niger* antennis apice capillaribus, elytris falciis tribus rufis. *Syst. Ent.* 725. 144.

*Cimex trifasciatus* antennis apice capillaribus corpore oblongiusculo nigro rubroque vario. *Linn. Syst. Nat.* 2. 725. 67.

*Schaeff. Icon.* tab. 13. fig. 8.

Habitat in Europa.

*olivaceus*. 203. *C. oblongus elytris fusco ferrugineis, apice coccineis, puncto atro, antennis apice capillaribus.* *Gen. Inf. Mant.* 299. \*

*Schaeff. Icon.* tab. 13. fig. 2.

Habitat in Germania. *D. Schulz.*

*seticornis*. 204. *C. oblongus ater, elytris fuscis, basi pallidis apiceque puncto coccineo, antennis apice capillaribus.* *Syst. Ent.* 725. 145. \*

Habitat Lipsiae.

Variat rarius capite rufo.

*capillaris*. 205. *C. oblongus flaves cens, elytris apice coccineis, antennis nigris, apice capillaribus.* *Syst. Ent.* 725. 146. \*

Habitat in Lipsiae hortis.

*gothicus*. 206. *C. oblongus niger, scutello elytrorumque apicibus coccineis, antennis apice capillaribus.* *Syst. Ent.* 726. 147. *Linn. Syst. Nat.* 2. 726. 73. *Fn. Sv.* 966.

*Cimex oblongus niger, thoracis lateribus scutelloque flavis, elytris, antennis pedibusque flavo variegatis.* *Geoff. Inf.* 1. 445. 19.

*Schaeff. Icon.* tab. 13. fig. 5.

*Fuesly Inf. Helv.* 26. 501.

Habitat in Europa frequens.

*spissicornis*. 207. *C. oblongus niger, pedibus flavis, antennis incrassatis, apice capillaribus.* *Gen. Inf. Mant.* 300. \*

Habitat Hamburgi. *Dr. Schulz.*

\*\*\*\* Antennis fetaceis.

*laevigatus*. 208. *C. oblongus exalbidus, lateribus albidis.* *Syst. Ent.* 726. 148. *Linn. Syst. Nat.* 2. 730. 101. *Fn. Sv.* 951.

*Cimex*

*Cimex oblongus albidus oculis nigris. Geoff. Inf. 1.*  
452. 26.

*Cimex oblongus albo viridi griseus, pedibus antennisque setaceis nigricantibus, thorace fasciis binis nigris, elytris albo cinctis. Degeer Inf. 3. 292. 30.*

*Fuesl. Inf. Helv. 26. 512.*

Habitat in Europae pratis frequens.

209. *C. oblongus niger, lateribus albidis. Gen. Inf. Mant. lateralis.*  
300. \*

Habitat Kiliae Holfatorum. *Dom. Lund.*

210. *C. oblongus viridis immaculatus, alis hyalinis. Gen. pabulinus.*  
*Inf. Mant. 301. \**

*Cimex pabulinus oblongiusculus flauescenti viridis totus. Linn. Syst. Nat. 2. 727. 83. Fn. Sv. 947. Scop. carn. 132.*

*Cimex oblongus totus viridis oculis fuscis. Geoff. Inf. 1. 456. 43.*

*Cimex oblongus viridis, oculis nigris, antennis longis setaceis. Degeer Inf. 3. 293. 31.*

Habitat in Europae pratis.

211. *C. oblongus viridis, plantis antennarumque apicibus rufis. Linn. Syst. Nat. 2. 730. 102.*

Habitat in Europa.

Duplo praecedente maior.

212. *C. oblongus, elytris ferrugineis latere albidis, antennis nigris, linea thoracis alba. Syst. Ent. 726. 149. dolabratus.*

*Linn. Syst. Nat. 2. 730. 103. Fn. Sv. 959.*

Habitat in Europae Festuca natante.

213. *C. oblongus, capite, thorace elytrisque nigris, scutello fulvo. Syst. Ent. 726. 150. Linn. Syst. Nat. 2.*

*730. 104.*

*Gronov. Zooph. 712.*

Habitat in India.

214. *C. oblongus, thorace elytrisque atris, linea flava fasciam elytrorum formante. Syst. Ent. 726. 151. \* Gronovii.*

*Linn. Syst. Nat. 2. 730. 106.*



374. CLASS. VII. RYNGOTA. Cimex.

Gronov. Zooph. 711.

Habitat in India.

- ferus.* 215. *C. oblongus* griseus immaculatus. *Syst. Ent.* 726. 152. *Linn. Syst. Nat.* 2. 731. 108. *Fn. Sv.* 962.  
Habitat in Europae sylvis.

- marginellus.* 216. *C. oblongus* niger, thoracis lineis tribus, elytris marginate omni albis punctoque apicis coccineo.  
Habitat in Italia. *D. Allioni.*  
Statura praecedentium. Caput atrum, immaculatum. Thorax ater lineis tribus albis. Elytra atra margine omni exteriori et interiori albis. Punctum coccineum elytra terminat.

- friatus.* 217. *C. oblongus* niger, elytris flavo fuscoque striatis, apice pedibusque rufis. *Syst. Ent.* 727. 153. *Linn. Syst. Nat.* 2. 730. 105. *Fn. Sv.* 960.  
*Cimex oblongus* niger, thorace fasciis tribus flavis, scutello nigro, elytris lineis flavis, apice fulvo. *Geoff. Inf.* 1. 454. 38.  
*Cimex oblongus* niger, antennis setaceis, pedibus rubris, thorace macula flava, elytris flavo nigroque striatis. *Degeer Inf.* 3. 290. 29. tab. 15. fig. 14. 15. *Petiv. Gazoph.* tab. 26. fig. 1.  
*Schaeff. Icon.* tab. 13. fig. 4.  
Habitat in Vlmi campestris foliis.

- Tiliae.* 218. *C. oblongus* virefcens, fasciis tribus fuscis, media angulata. *Gen. Inf. Mant.* 301. \*  
Habitat Kiliae Holsatorum. *Dom. Lund.*

- Populi.* 219. *C. oblongus* albo fuscoque nebulosus. *Syst. Ent.* 727. 154. *Linn. Syst. Nat.* 2. 731. 109. *Fn. Sv.* 963.  
Habitat in Europae Populo tremula.

- agilis.* 220. *C. oblongus*, thorace atro, margine postico flavo, elytris fuscis basi apiceque pallidis.  
Habitat in Germania. *Dom. de Hattorf.*  
Statura et magnitudo praecedentis. Caput atrum nitidum rostro pallido antennarumque articulo primo rufescente. Thorax ater nitidus margine postico late flavo. Scutellum atrum, apice flavum. Elytra fusca  
vd

CLASS. VII. RYNGOTA. Cimex. 379

vel potius picea macula pone basin pallida. Apex pallidus terminatur puncto atro. Abdomen atrum. Pedes pallidi. Simillimum ex Italia misit *D. Allioni*, at absque thoracis margine pallido.

221. *C. oblongus niger*, scutello maculisque tribus elytrorum flavis. *Gen. Inf. Mant.* 301.\* *6guttatus.*  
Habitat in Europa.
222. *C. oblongus maculato fulvus*, pedibus rufis, femoribus crassis dentatis. *Linn. Syst. Nat.* 2. 732. 115. *Abietis.*  
*Cimex grossipes* depressus rufo fuscus, capite thoraceque nigris, femoribus anticis crassis dentatis. *Degeer Inf.* 3. 308. 37. tab. 15. fig. 20. 21.  
Habitat in Europae borealis Abiete.
223. *C. oblongus supra rubiginosus*, elytris strilis sanguineis, alis postice albo fuscoque variis. *Syst. Ent.* 727. 155. *Linn. Syst. Nat.* 2. 731. 110. *Fu. Sv.* 964. *Vlmi.*  
Habitat in Europae Vlmo.
224. *C. corpore elongato fusco flauoque vario*, femoribus posticis elongatis dentatis. *Syst. Ent.* 727. 156.\* *pedestris.*  
Habitat in India orientali. *D. Koenig.*
225. *C. antennis setaceis ferrugineis*, corpore lineari, pal- lide virescenti. *Syst. Ent.* 727. 157.\* *filiformis.*  
Habitat in Americae meridionalis insulis.

\*\*\*\*\* Linearis corpore angusto.

226. *C. linearis*, supra fuscus, thoracis scutellique marginibus lineaque dorsali flavis. *Syst. Ent.* 727. 158.\* *fossarum.*  
Habitat in India orientali. *D. Koenig.*
227. *C. linearis niger*, pedibus anticis brevissimis. *Syst. lacustris.* *Ent.* 728. 159. *Linn. Syst. Nat.* 2. 732. 117. *Fu. Sv.* 970.  
*Cimex Naias* aquaticus linearis niger, pedibus anticis brevissimis, reliquis longissimis. *Degeer Inf.* 3. 311. 39. tab. 16. fig. 7.  
*Cimex aquaticus* figurae longioris. *Raj. Inf.* 57. 1.

376 CLASS. VII. RYNGOTA. Cimex.

Insectum Tipula dictum. *Bauh. Ball.* 233. fig. 1.

*Geoff. Inf.* 1. 463. 59.

*List. nat.* tab. 5. fig. 4.

*Bradl. nat.* tab. 26. fig. 2. d.

*Sulz. Inf.* tab. 11. fig. 78.

*Scop. carn.* 394.

Habitat in Europae aquis, riuiis frequens.

*flagnorum.* 228. *C. linearis teretiusculus niger, thorace medio punctis duobus globosis. Syst. Ent.* 728. 160. *Linna. Syst. Nat.* 2. 732. 118. *Fa. Sv.* 971.

*Cimex linearis nigricans, compressus, capite cylindrico, pedibus anticis brevissimis. Geoff. Inf.* 1. 463. 60.

*Cimex Acus aquaticus linearis niger, capite tereti longissimo, pedibus omnibus aequalibus. Degeer Inf.* 3. 322. 40. tab. 15. fig. 24.

*Sulz. Hist. Inf.* tab. 10. fig. 17.

*Scop. carn.* 395.

Habitat in Angliae lacubus frequens.

*riuulorum.* 229. *C. linearis niger, albo punctatus, abdomine fuluo. Syst. Ent.* 728. 161. \*

Habitat in riuiis montosis Alsatiae. *Prof. Hermann.*

*pallipes.* 230. *C. linearis niger, pedibus pallidis, pectore bispinoso.*

Habitat in Italia. *D. Allioni.*

Corpus statura praecedentium totum nigrum pedibus folis pallidis. Spina valida prominens vtrinque sub thorace. Abdominis margo parum pallefcit.

*culiciformes.* 231. *C. thorace multispinose griseus, elytris lituris nigris, corpore elongato. Syst. Ent.* 728. 162. \*

Habitat in Americae aquis. *Dom. v. Rohr.*

*flum.* 232. *C. linearis fuscus, alis abbreviatis, pedibus longissimis. Gen. Inf. Mant.* 302. \*

*Cimex longipes linearis fuscus, rostro arcuato, antennis pedibusque longissimis, anticis dentatis. Degeer Inf.* 3. 352. 26. tab. 35. fig. 16. ?

Habitat in India. *D. Fothergill.*

CLASS. VII. RYNGOTA. Redupius. 377

233. *C. linearis ater*, thoracis margine, elytrorumque fascia sanguinea. *Syst. Ent.* 728. 163. \*  
Habitat in America. *Dom. v. Rohr.*  
Femora postica cingulo rubro.
234. *C. linearis supra fuscus*, subtus cinerascens, pedibus *cursitans*. longissimis, ano bidentato. *Syst. Ent.* 729. 164. \*  
Habitat in noua Hollandia. *Mus. Dom. Banks.*
235. *C. linearis exalbidus*, pedibus omnibus longissimis, *tipulari-* femoribus clauatis, antennis bicaudatis. *Syst. Ent.* us. 729. 165. \* *Linn. Syst. Nat.* 2. 733. 120. *Fn. Sv.* 973.  
*Cimex arboreus culiciformis.* *Frijch. Inf.* 7. 28. tab. 20.  
Habitat in Europae borealis muscia.
236. *C. linearis griseus*, pedibus pallidis, anticis breuissimis. *Syst. Ent.* 729. 166. *vagabundus.*  
*Cimex vagabundus linearis variegatus*, pedibus anticis breuissimis crassis inflexis. *Linn. Syst. Nat.* 2. 732. 119.  
*Cimex culiciformis linearis fuscus albo maculatus*, antennis pedibusque quatuor posticis longissimis, anterioribus breuissimis. *Degeer Inf.* 3. 323. 41. tab. 17. fig. 1. 2.  
*Cimex arborum oblongus*, alarum signatura alba. *Frijch. Inf.* 7. 11. tab. 6.  
*Geoff. Inf.* 1. 462. 58.  
Habitat in Europae Muscis.
237. *C. linearis cinereus*, femoribus clauatis, antennis bicaudatis. *Syst. Ent.* 729. 167. \* *clauipes.*  
Habitat in Succia.

164. R E D V V I V S.

1. *R. niger*, thoracis margine elytrorumque fascia obsoleta flexuosa rufis. *Syst. Ent.* 729. 1. \* *Gigas.*  
Habitat in India orientali.
2. *R. antennis apice capillaribus*, corpore subuilloso fusco. *Syst. Ent.* 730. 2. *personatus.*

978 CLASS. VII. RYNGOTA. Reduuius.

*Cimex personatus*. Linn. *Syst. Nat.* 2. 724. 64. *Fn. Sv.* 492. *Geoff. Inf.* 1. 436. 4. tab. 9. fig. 3.

*Cimex quisquilius* oblongus obscure fuscus, antennis setaceis capillaceis, rostro breui arcuato. *Degeer Inf.* 3. 281. 25. tab. 15. fig. 7.

*Cimex maximus* pullus, alis nudis ex toto membranaceis. *List. Scar. Angl.* 397. 38.

*Cimex stercorarius* maior oblongus. *Frisch. Inf.* 10. tab. 10.

*Musca cimiciformis* grauiter olens. *Raj. Inf.* 56.

*Sulz. Inf.* tab. 11. fig. 74.?

*Schaeff. Icon.* tab. 13. fig. 6. 7.

— — — tab. 57. fig. 9.

*Fuesly Inf. Helv.* 26. 499.

Habitat in Europae quisquiliis.

*Larua* horrida personata confumit A. lectulariam.

*barbicornis*. 3. *R. niger*, thorace abdominisque basi oliuaceis. *Syst. Ent.* 730. 3. \*

Habitat in Sierra Leon Africae. *Dom. Drury.*

Antennae in altero sexu barbatae.

Variat rarius thorace nigro.

*maculatus*. 4. *R. rufus*, thorace maculis quatuor elytris tribus nigris. Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*

*C. annulato* minor. Caput sanguineum antennis nigris primo articulo rufo. Thorax cruce media impressa rufus maculis quatuor nigris. Elytra rufa maculis tribus magnis nigris. Alae nigrae. Corpus subrufum vtrinque nigro punctatum. Pedes rufi.

*nitidulus*. 5. *R. ater*, thorace oliuaceo, femoribus anticis rufis.

Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*

Nimis affinis *R. barbicorni*, differt tamen corpore toto nitido. Thorax oliuaceus lobo antico margineque laterali distinctis. Pedes atrii femoribus anticis solis rufis.

*longipes*. 6. *R. ruber*, elytris nigris basi fasciaque rubris. *Syst. Ent.* 730. 4. \*

*Cimex longipes*. Linn. *Syst. Nat.* 2. 724. 65.

*Gronov. Zooph.* 709.

Habitat in Insula St. Thomas Americae meridionalis.

7. *R.*

7. R. supra rufus, alis nigris, abdomine atro, margine *marginatus*.  
rufus.

Habitat in India orientali. *Muf. Dom. Banks.*

Medius. Caput rufum rostro antennisque nigris. Thorax elytraque rufa, immaculata. Alae nigrae. Pedes nigri femoribus anticis rufis.

8. R. supra niger, elytris puncto apicis rufo. *apustulatus*.  
Habitat Surinami. *Muf. Dom. Banks.*

Statura R. annulati. Caput atrum, immaculatum. Thorax lobo antico distincto spinaque parua antica acuta. Elytra nigra puncto apicis rufo. Alae nigrae. Abdomen nigrum, apice rufum. Pedes antici nigri femoribus apice, tibiis basi rufis. Pedes reliqui rufi, femoribus annulo fusco.

9. R. antennis apice capillaribus, corpore nigro subtus *annulatus*.  
sanguineo maculato. *Syst. Ent.* 730. 5.

*Cimex annulatus*. *Linn. Syst. Nat.* 2. 725. 71. *Fn. Sv.* 943. *Geoff. Inf.* 1. 437. 5.

*Cimex niger rufipes* oblongus niger, antennis setaceis, capillaceis, rostro breui arcuato, pedibus abdominisque lateribus rubro maculatis. *Degeer Inf.* 3. 286. 26.

*Sulz. Hist. Inf.* tab. 10. fig. 13.

*Schaeff. Icon.* tab. 5. fig. 9. 10. 11.

*Fuesl. Inf. Helv.* 26. 500.

Habitat in Europae Corylo.

10. R. rufescens, antennis pedibusque flauo nigroque *cinctus*.  
annulatis. *Gen. Inf. Mant.* 302. \*

Habitat in America boreali.

11. R. ater, thorace postice, elytris abdominisque *bicolor*.  
margine obscure pallidis.

Habitat in Africa aequinoctiali. *Muf. Dom. Banks.*

Paruus. Caput atrum, immaculatum. Thorax obscure pallidus lobo antico distincto nigro. Elytra obscure fusca, immaculata. Alae atrae. Abdomen atrum margine pallido. Pedes atrii.

12. R. testaceo nigroque varius, thorace antice testaceo, *attelaboides*.  
dentibus duobus nigris. *Syst. Ent.* 731. 6. \*

Habitat in noua Hollandia. *Muf. Dom. Banks.*

13. R.

**quadridens.** 13. *R. fuscus*, capite abdominis margine pedibusque oliuaceis, thorace antice quadridentato.

Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*

**Medius.** Caput cylindricum, postice attenuatum, oliuaceum. Thorax fuscus lobo antico distincto spinis quatuor eleuatis breuib. Margo postice vtrinque videntatus. Elytra nigra, immaculata. Alae nitidae. Abdomen nigrum cyaneo nitidum margine apicis oliuaceo. Pedes oliuacei genubus nigris.

**nigripes.** 14. *R. elytrorum* basi antennis pedibusque rubris, tibiis anticis hirsutissimis.

*Cimex nigripes.* *Linn. Syst. Nat.* 2. 730. 100.

*Cimex hirtipes* oblongus, thorace nigro, elytrorum basi, antennis pedibusque quatuor rubris, tibiis anticis nigris hirsutissimis. *Degeer Inf.* 3. 344. 20. tab. 35. fig. 1.

Habitat in America meridionali.

**collaris.** 15. *R. ater*, elytris postice albis, capite postice attenuato cylindrico.

Habitat in India orientali. *Mus. Dom. Banks.*

**Statura** *R. attelaboidis*. Caput atrum, nitidum, postice attenuatum, cylindricum. Elytra nigra macula magna postica alba. Pedes simplices, atri.

**varius.** 16. *R. rufo nigroque varius*, abdominis margine dente medio erecto obtuso. *Syst. Ent.* 731. 7.\*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

**sexguttatus.** 17. *R. thorace* subspinoso niger, elytris maculis tribus albis. *Syst. Ent. App.* 832.\*

Habitat in India orientali. *Dom. Edler.*

**quadriguttatus.** 18. *R. thorace* postice testaceo, elytris alisque atris, maculis quatuor laete flauis.

Habitat in India orientali. *Mus. Dom. Banks.*

**Reliquis** minor corpore elongato angustiore. Caput nigrum rostro antennisque testaceis. Thorax niger lobo postico distincto testaceo. Scutellum nigrum basi testaceum. Elytra et alae atra, iuncta maculis quatuor dorsalibus flauis, duae ad basin, tertia in medio

CLASS. VII. RYNGOTA. Reduuius. 381

medio et quarta apicis. Abdomen nigrum margine testaceo. Pedes testacei femoribus anticis incrassatis.

19. R. thorace testaceo, elytris alisque nigris, macula media flava. *punctum.*

Habitat in India orientali. *Mus. Dom. Banks.*

Statura et summa affinitas praecedentis. Caput cum antennis et rostro testaceum. Thorax testaceus lobo postico distincto. Elytra alaeque nigra macula magna communi flava. Abdomen nigrum margine flavo. Pedes testacei femoribus anticis incrassatis macula fusca.

20. R. niger, elytris basi pallidis, alis macula communi rufa, pectore spinoso. *hamatus.*

Habitat in America meridionali. *Mus. D. Hunter.*

Medius. Caput atrum. Thorax ater, immaculatus lobo antico distincto. Pectus spina valida hamata siue incurva inter femora antica. Scutellum nigrum. Elytra margine interiori baseos pallido. Alae nigrae macula communi rufa. Abdominis margo elevatus, acutus rufo maculatus. Pedes obscuri femoribus apice nigris, anticis crassioribus.

21. R. niger, thorace abdominisque marginibus rubro maculatis, elytris rufis. *Syst. Ent. 731. 8. \** *iracundus.*

*Cimex iracundus. Scop. carn. 378.*

*Cimex iracundus* oblongus niger, thorace scabro postice rubro, elytris rubris, abdominis lateribus rubro nigroque variis. *Pod. Mus. 58. 17.*

*Cimex sanguineus* rostro arcuato, corpore oblongo nigro, elytris rubris, abdominis lateribus rubro nigroque variis. *Schrank. Beytr. 81. 37. tab. 3. fig. 17.*

Habitat in Germania australiori.

22. R. thorace scutelloque acute spinosis, elytris puncto albo. *Syst. Ent. 731. 9.* *biguttatus.*

*Cimex biguttatus. Linn. Syst. Nat. 2. 725. 70.*

Habitat in Guinea. *Mus. D. Schloffer.*

23. R. niger, elytris ferrugineis, rostro antennis tibiisque flavis, scutello cristato serrato. *Syst. Ent. 732. 10.* *serratus.*

*Cimex*



382 CLASS. VII. RYNGOTA. Reduuius.

*Cimex cristatus* rostro arcuato, antennis setaceis, corpore fulco ferrugineo, scutello cristato. *Linn. Syst. Nat.* 2. 723. 62. *Amoen. acad.* 6. 399. 42.?

*Cimex carinatus*. *Forst. nou. spec.* 72. 72.

*Sulz. Hist. Inf.* tab. 10. fig. 12.

*Drury Inf.* 2. tab. 36. fig. 6.

Habitat in America meridionali.

*albipes*. 24. *R. fuscus*, antennarum annulis tibiis abdominisque punctis lateralibus albis.

Habitat in Europa media. *Dom. de Hattorf.*

Statura *R. aegyptii*, at paullo maior. Antennae nigrae annulis tribus albis, primo maiore. Corpus fuscum, supra immaculatum, subtus punctis lateralibus albis. Pedes nigri femoribus basi tibiisque totis albis.

*aegypti*- 25. *R. corpore villoso griseo, abdominis margine nigro*  
*us.* variegato. *Syst. Ent.* 732. 11. \*

Habitat in Aegypto. *Prof. Forskahl.*

*diadema*. 26. *R. niger, capite thoraceque spinosis.* *Gen. Inf. Mant.*  
302. \*

Habitat in America boreali.

*5spinosus*. 27. *R. thorace nigro, postice spinis quatuor flauis, scutello*  
apice recuruo spinoso.

Habitat in India orientali. *Mus. Dom. Banks.*

Caput postice attenuatum, nigrum, antennis rostroque pallidis. Thorax niger lobo antico distincto posticeque spinis quatuor erectis validis acutis flauis. Scutellum nigrum, apice eleuatum, recuruum, spinosum. Elytra nigra puncto bascos fasciaque communi in medio flauis. Alae nigrae, apice pallidiores. Abdomen deest. Pedes pallidi.

*maurus*. 28. *R. ferrugineus, abdominis margine nigro maculato,*  
thorace antice subspinoso. *Syst. Ent.* 732. 12. \*

Habitat in Mauritania.

*Acantha-* 29. *R. thorace spinoso, abdomine spinis ciliato.* *Syst. Ent.*  
*ris.* 732. 13. \*

*Cimex Acantharis.* *Linn. Syst. Nat.* 2. 720. 38.

*Acan-*

*Acantharis. Brown. Iam. 434. tab. 44. fig. 11.*

*Sulz. Hist. Inf. tab. 10. fig. 8.*

Habitat in Iamaica.

10. *R. elongatus rufus, antennis pedibusque nigris. elongatus.*  
 Habitat in Africa aequinoctiali. *Mus. Dom. Banks.*  
 Corpus reliquis multo angustius, longius, rufum, im-  
 maculatum rostro apice antennis pedibusque nigris.

# 165. P V L E X.

1. *P. rostro corpore breuiore. Syst. Ent. 732. 1. Linn. irritans.*  
*Syst. Nat. 2. 1021. 1. Fn. Sv. 1695.*

*Pulex. Geoff. Inf. 2. 616. 1.*

*Pulex vulgaris. Raj. Inf. 7.*

*Scop. carn. 1055.*

*Alb. Mant. tab. 41.*

*Bonann. Micrograph. fig. 56.*

*Ioblot. Microgr. 1. 1. tab. 3.*

*Frisch. Inf. 11. 8.*

*Valijn. Oper. 1. tab. 25. fig. 1.*

*Leuwenh. Epist. 76. fig. 1 — 20.*

*Roef. Inf. 2. Musc. tab. 2. 3. 4.*

*Backer. microsc. tab. 13. fig. 6.*

*Ledermüll. microsc. 4. tab. 20.*

*Sulz. Inf. tab. 22. fig. 146.*

*Schaeff. Elem. tab. 105.*

Habitat vbique in Europa, in America, sanguisugus.

Leporibus valde molestus ideoque ramento vestimenti

Ieporini capitur et avertitur a Dalekarlis. Pellitur

Satureia, Pulegio, Persicaria, Cotula, Alno.

Mas minor sub femina iungitur.

2. *P. proboscide corporis longitudine. Syst. Ent. 733. 2. pene-*  
*Linn. Syst. Nat. 2. 1021. 2. trans.*

*Pulex cutem penetrans. Catesb. Carol. 3. 10. tab. 10.*  
*fig. 3.*

*Acarus fuscus sub cute nidulans, proboscide acutiori.*

*Brown. Iam. 418.*

*Tunga. Margrav. Brasil. 249.*

*Sloan. Iam. 2. 191. introd. 125.*

Habitat

Habitat in America pedes hominum intrans, oua c  
nens, cacoethem, saepe mortem caussans, caut  
trahendus.

Abdomen feminae ouis innumeris grauidum, orbi  
tum, vti in Tèrmite fatali ad magnitudinem c  
plam totius corporis intumescit.

## 166. A P H I S.

- Lychni-* 1. A. Lychnidis dioicae. *Syst. Ent.* 733. 1. \* *Linn.*  
*dis.* *Nat.* 2. 734. 7. *Fn. Sv.* 980.  
Habitat frequens in Lychnidis dioicae caulibus.
- Capreae.* 2. A. Salicis Capreae. *Syst. Ent.* 734. 2. \*  
Habitat in Europae Salice.
- Sambuci.* 3. A. Sambuci nigrae. *Syst. Ent.* 734. 3. \* *Linn. Syst.*  
2. 734. 4. *Fn. Sv.* 978.  
*Aphis sambuci* tota coerulea atra. *Geoff. Inf.* 1. 4.  
*Cimex* exiguus nunc niger, nunc viridis, cui ala  
toto membranaceae, praegranda, *List. Angl.* 397  
*Frisch. Inf.* 11. 14. tab. 18.  
*Reaum. Inf.* 3. tab. 8. fig. 5 — 15.  
Habitat in Sambuci nigri ramis.
- Solidagi-* 4. A. Solidaginis virgaureae.  
*nis.* Habitat in Solidaginis Virgaureae caulibus Norwa  
Corpus maiusculum, fuscum, antice angustatum, fi  
te emarginata. Cornicula elongata setacea. St  
corniculis duplo breuior. Pedes pallidi genic  
nigris.  
Iuniores omnino testacei.
- Cerasi.* 5. A. Pruni Cerasi. *Syst. Ent.* 734. 4. \*  
Habitat in Pruni Cerasi foliis.
- Ribis.* 6. A. Ribis rubri. *Syst. Ent.* 734. 5. \* *Linn. Syst. N*  
2. 733. 1. *Fn. Sv.* 977.  
*Leuwenh. arc. epist.* 90. 545. tab. 548.  
*Blank. Inf.* 164. tab. 14. fig. D. 2.  
*Frisch. Inf.* 11. 9. tab. 14.  
*Reaum. Inf.* 3. tab. 22. fig. 7 — 10.  
Habitat frequens sub foliis monstrosis pustulatis Ribis  
bri.

7. A. Arundinis Epigeios. *Syst. Ent.* 734. 6.\* *Arundinis.*  
Habitat in Arundinis Epigeios foliis.
8. A. Populi nigrae. *Syst. Ent.* 735. 7.\* *Linn. Syst. Nat. burfaria.*  
2. 736. 29. *Fn. Sv.* 998.  
*Aphis Populi nigra lanata. Geoff. Inf.* 1. 497. 11.  
*Seb. Mus.* 1. tab. 38. fig. 8.  
*Swammerd. Bibl.* 45. fig. 22 — 25.  
*Reaum. Inf.* 3. tab. 26. fig. 7 — 11.  
Habitat in facculis ouatis Populi nigrae.
9. A. Gallii Aparines. *Syst. Ent.* 735. 8.\* *Aparines.*  
Habitat in Gallii Aparines caulibus.
10. A. Aceris campestris. *Syst. Ent.* 735. 9.\* *Linn. Syst. Aceris.*  
*Nat.* 2. 736. 31. *Fn. Sv.* 999.  
*Aphis Aceris viridis maculis nigris. Geoff. Inf.* 495. 5.  
*Reaum. Inf.* 3. tab. 22. fig. 6 — 10.  
Habitat in Aceris campestris foliis.
11. A. Rumicis Lapathi. *Syst. Ent.* 735. 10.\* *Linn. Syst. Rumicis.*  
*Nat.* 2. 734. 5. *Fn. Sv.* 979.  
Habitat in Rumice acuto.
12. A. Pruni domesticae. *Syst. Ent.* 735. 11.\* *Geoff. Inf. Pruni.*  
1. 497. 10.  
*Aphis Pruni dilute viridis albo farinosa, corniculis*  
*brevibus Pruni. Degeer Inf.* 3. 49. 5. tab. 2. fig.  
1 — 8.  
*Reaum. Inf.* 3. tab. 23. fig. 9. 10.  
Habitat in Pruni domesticae foliis.
13. A. Cardui Acanthoidis. *Syst. Ent.* 735. 12.\* *Linn. Syst. Cardui.*  
*Nat.* 2. 735. 17. *Fn. Sv.* 988.  
Habitat in Cardui Acanthoidis caulibus.
14. A. flavesceus, abdomine viridi, antennis pedibusque *Achilleas.*  
albidis. *Gen. Inf. Mant.* 202.\*  
Habitat in Achillea Millefolio frequens Kiliae.
15. A. Corni sanguineae. *Syst. Ent.* 736. 13.\* *Corni.*  
Habitat in Corni sanguineae foliis.  
*Fabr. Sp. Inf. T. II.* B b 16. A.

*Reaum. Inf.* 3. tab. 25. fig. 4 — 7.

Habitat in *Vlmo campestri*.

- Papaueris.* 33. *A. Papaueris somniferi. Gen. Inf. Mant.* 203.\*  
Habitat in *Hollatiae Papauere somnifero*.

- Tiliae.* 34. *A. Tiliae Europaeae. Syst. Ent.* 738. 27.\* *Linn. Syst. Nat.* 2. 734. 11. *Fn. Sv.* 984.

*Aphis Tiliae* alis antennis pedibusque nigro punctatis.  
*Geoff. Inf.* 1. 495. 6.

*Aphis Tiliae nigro punctata* flauo viridis lineis punctisque nigris, alis nigro maculatis, corniculis brevioribus. *Degeer Inf.* 3. 77. 12. tab. 5. fig. 1 — 5.

*Pediculus arboreus Tiliae. Frisch. Inf.* 11. 1. 3. tab. 17.

*Redum. Inf.* 3. tab. 23. fig. 1 — 8.

Habitat in *Europae Tilia*.

- Iuniperi.* 35. *A. Iuniperi communis.*

*Aphis Iuniperi* obscure fusca cinereo maculata, corniculis brevioribus grossis *Iuniperi. Degeer Inf.* 3. 56. 7. tab. 4. fig. 7. 8.

Habitat in *Iuniperi communis foliis*.

- Brassicae.* 36. *A. Brassicae oleraceae. Syst. Ent.* 738. 28.\* *Linn. Syst. Nat.* 2. 734. 12. *Fn. Sv.* 985.

*Pediculus Brassicae. Frisch. Inf.* 11. 10. tab. 3. fig. 15.

Habitat in *Brassicae oleraceae foliorum inuersa pagina*, vnde folia tabescunt.

- Roboris.* 37. *A. Quercus Roboris. Syst. Ent.* 739. 29.\*

*Aphis Quercus Roboris* nigra, corniculis obsoletis atris medii abdominis. *Linn. Syst. Nat.* 2. 735. 22. *Fn. Sv.* 993.

Habitat in *Quercu Robore*.

- Quercus.* 38. *A. Quercus Roboris. Syst. Ent.* 739. 30.\*

*Aphis Quercus* proboscide longissima. *Linn. Syst. Nat.* 2. 735. 24.

*Aphis fusca* proboscide corpore triplo longiore. *Geoff. Inf.* 1. 498. 14.

*Reaum. Inf.* 3. tab. 28. fig. 5 — 14.

Habitat in *Galliae Quercu*.

39. *A. Pini sylvestris ramulorum.* *Linn. Syst. Nat. 2. Pini.*  
736. 25. *Fn. Sv.* 494.  
*Aphis nuda Pini fusca tuberculata, pedibus nudis Pi-*  
*ni.* *Degeer Inf.* 3. 27. 1. tab. 6. fig. 9 — 16.  
Habitat in ramis iunioribus Pini sylvestris.
40. *A. Pini sylvestris foliorum.* *Pineti.*  
*Aphis tomentosa Pini nigra albo farinosa tuberculata,*  
*pedibus villosa tomentosis.* *Degeer Inf.* 3. 39. 2. tab.  
6. fig. 19. 33.  
Habitat in Pini sylvestris foliis.  
Species distincta corpore nigro albo farinaceo pedibus-  
que posticis elongatis ciliatis.
41. *A. Salicis.* *Syst. Ent.* 739. 31. \* *Linn. Syst. Nat. 2. Salicis.*  
736. 26. *Fn. Sv.* 998.  
*Reaum. Inf.* 3. tab. 22. fig. 2.  
Habitat in Salicibus variis.
42. *A. Pistaciae Lentisci.* *Syst. Ent.* 739. 32. \* *Pistaciae.*  
*Aphis Pistaciae nigra, alis albidis, tibiis longissimis,*  
*thorace verrucoso.* *Linn. Syst. Nat. 2. 737. 33.*  
Habitat in Pistacia Lentisco, in folliculo foliorum in-  
carnato, ventricoso, vtrinque attenuato, spithamaco  
pollice crassiori.
43. *A. Rumicis Acetosae.* *Syst. Ent.* 739. 33. \* *Linn. Syst. Acetosae.*  
*Nat. 2. 734. 6.*  
*Aphis Acetosae atra, fascia transversa viridi.* *Geoff.*  
*Inf.* 1. 496. 9.  
*Reaum. Inf.* 3. 286.  
Habitat in Rumice acetosa.
44. *A. Ligustici Scotici.* *Ligustici.*  
*Aphis Ligustici nigra, antennis tibiis basi pallidis.* *It.*  
*Norwag.* d. 3. Aug.  
Habitat in Norwagiae Ligustico scotico.
45. *A. Pruni Padi.* *Syst. Ent.* 739. 34. *Linn. Syst. Nat. Padi.*  
2. 734. 8. *Fn. Sv.* 981.  
*Reaum. Inf.* 3. tab. 23. fig. 9. 10.  
Habitat in Europae Pruno Pado.

*Vermis fucorius Alni*. *Frisch. Inf.* 8. 48. tab. 13.

*Schaeff. Elem.* tab. 39.

*Sulz. Inf.* tab. 12. fig. 80.

Habitat in Bëtulae Alni ramulis.

*Abietis.* 13. *C. Pini Abietis.* *Syst. Ent.* 741. 11. *Linn. Syst. Nat.* 2. 738. 13. *Fn. Sv.* 1011.

*Aphis Gallarum Abietis tomentosa*, corniculis nullis gallarum Abietis. *Degeer Inf.* 3. 99. 17. tab. 8. fig. 1 — 3.

*Psylla pallide flavescentes*, oculis fuscis, alis aqueis. *Geoff. Inf.* 1. 487. 5.

*Insectum tuberculi muricati arboris Taxi.* *Frisch. Inf.* 12. 10. tab. 2. fig. 3.

*Picea pumila.* *Clus. pann.* 20. 21.

Habitat frequens in gallis strobiliformibus squamatis, imbricatis turionum Abietis.

*Salicis.* 14. *C. Salicis.* *Syst. Ent.* 742. 12. *Linn. Syst. Nat.* 2. 739. 14. *Fn. Sv.* 1012.

Habitat in Europae Salicibus.

*Fraxini.* 15. *C. Fraxini excelsioris.* *Syst. Ent.* 742. 13. *Linn. Syst. Nat.* 2. 739. 15. *Fn. Sv.* 1013.

*Psylla nigro luteoque variegata*, alarum ore in apicibus fuscis. *Geoff. Inf.* 1. 486. 4.

Habitat in Fraxino.

*Aceris.* 16. *C. Aceris platanoidis.* *Syst. Ent.* 742. 14. \* *Linn. Syst. Nat.* 2. 739. 16. *Fn. Sv.* 1014.

Habitat in racemis aceris platanoidis.

*Ficus.* 17. *C. Ficus Caricae.* *Syst. Ent.* 742. 15. *Linn. Syst. Nat.* 2. 739. 17.

*Psylla fusca*, antennis crassis pilosis, alarum nervis fuscis. *Geoff. Inf.* 1. 484. 1.

*Reaum. Inf.* 3. tab. 29. fig. 17 — 24.

Habitat in Europae australioris Ficu.

168. C O C C V S.

1. C. hybernaculorum. *Syst. Ent.* 742. 1. *Linn. Syst. Nat. Hesperidum.*  
2. 739. 1. *Fn. Sv.* 1015.  
*Chermes Hesperidum.* *Geoff. Inf.* 1. 505. 2.  
*Schaeff. Elem.* tab. 4b.  
*Aët. parif.* 1692. 14. tab. 14.  
*Reaum. Inf.* 4. tab. 1.  
*Sulz. Inf.* tab. 12. fig. 81.  
*Frisch. Inf.* 12. 12.  
Habitat in arboribus semper virentibus hybernaculo-  
rum, frequentissimus.
1. C. Indarum arborum. *Gen. Inf. Mant.* 303. 2. *Linn. Adonidum.*  
*Syst. Nat.* 2. 739. 2.  
Habitat in Asiae arboribus semper virentibus.
3. C. Adonidum. *Syst. Ent.* 743. 2. *Adoni-*  
*Coccus adonidum* rufa farinaceo pilola. *Linn. Syst. Nat.* 2. 740. 4. *dum.*  
*Coccus Adonidum* corpore roseo farinaceo, alis setisque  
niueis. *Geoff. Inf.* 2. 511. 1.  
Habitat in arboribus Americae, Africae, Europae austr-  
lioris.
4. C. Quercus Roboris. *Syst. Ent.* 743. 3. *Linn. Syst. Quercus.*  
*Nat.* 2. 740. 5. *Fn. Sv.* 1016.  
*Chermes Quercus reniformis.* *Geoff. Inf.* 1. 508. 13.  
*Reaum. Inf.* 4. tab. 6. fig. 1 — 4.  
Habitat in Quercu Robore.
5. C. Quercus cocciferae. *Syst. Ent.* 743. 4. *Linn. Syst. Ilicis.*  
*Nat.* 2. 740. 6.  
*Garid. aixens.* 250. tab. 53.  
*Reaum. Inf.* 4. tab. 5.  
*Ledermüll. microsc.* 72. tab. 36.  
Habitat in quercu coccifera Galliae tinctorius.
1. C. Vlmi campestris. *Syst. Ent.* 743. 5. *Linn. Syst. Nat. Vlmi.*  
2. 740. 9. *Fn. Sv.* 1019.  
*Chermes Vlmi rotundus.* *Geoff. Inf.* 1. 507. 8.  
*Coccus ouatus Vlmi*, ouatus albus fusco transuerse stri-  
tus Vlmi. *Degeer Inf.* 6. 436. 1. tab. 28. fig. 7.  
Habitat in Vlmo campestri.



394 CLASS. VII. RYNGOTA. Coccus.

- Coryli.* 7. *C. Coryli auellanae.* *Syst. Ent.* 743. 6. *Linn. Syst. Nat.* 2. 741. 10. *Fn. Sv.* 1020.  
*Chermes Coryli hemisphaericus.* *Geoff. Inf.* 1. 507. 10.  
*Reaum. Inf.* 4. tab. 3. fig. 4—10.  
Habitat in *Corylo auellana*.
- Betulae.* 8. *C. Betulae albae.* *Gen. Inf. Mant.* 304. \* *Linn. Syst. Nat.* 2. 740. 7. *Fn. Sv.* 1017.  
Habitat in *Europae Betula alba*.
- Carpini.* 9. *C. Carpin Betuli.* *Gen. Inf. Mant.* 304. *Linn. Syst. Nat.* 2. 740. 8. *Fn. Sv.* 1018.  
Habitat in *Carpino Betulo*.
- Tiliae.* 10. *C. Tiliae Europaeae.* *Syst. Ent.* 743. 7. *Linn. Syst. Nat.* 2. 741. 11. *Fn. Sv.* 1021.  
*Chermes Tiliae hemisphaericus.* *Geoff. Inf.* 1. 507. 9.  
*Reaum. Inf.* 4. tab. 3. fig. 1—3.  
Habitat in *Tilia Europaea*.
- Rusci.* 11. *C. Rusci.* *Syst. Ent.* 743. 8.  
*Coccus Rusci testa octo clypeolis cincta.* *Linn. Syst. Nat.* 2. 741. 12.  
*Lepas noua f. Myrti morbus.* *Column. purp.* 16. tab. 17.  
*Balanus terrestris compositus e multis testis.* *Gerian. Adr.* 60. tab. 3. fig. 27.  
*Bocc. Musf.* tab. 107. fig. 33. h. h.  
Habitat in *Italiae Myrto, Rusco*.
- Salicum.* 12. *C. Capreae.* *Gen. Inf. Mant.* 304. *Linn. Syst. Nat.* 2. 741. 14.  
*Coccus rotundus Salicis subrotundus fuscus linea dorsali nigra Salicis.* *Degeer Inf.* 6. 440. 2. tab. 11. fig. 13.  
Habitat in *Europae Salicibus*.
- Pilosellae.* 13. *C. radices Hieracii Pilosellae.* *Syst. Ent.* 744. 9. *Linn. Syst. Nat.* 2. 742. 18. *Fn. Sv.* 1024.  
*Ad. Vpsf.* 1742. 54. tab. 2.  
Habitat in *Hieracii Pilosellae radicibus*.
- Arbuti.* 14. *C. radices Vucae vrsi.* *Syst. Ent.* 744. 10. *Linn. Syst. Nat.* 2. 742. 19.  
Habitat in *radicibus Arbuti vucae vrsi tinctorius*.

CLASS. VII. RYNGOTA. Coccus, 395

- C. radicum Phalaridis. *Syst. Ent.* 744. 11. *Linn. Syst. Nat.* 2. 742. 20. *Phalaridis.*  
*Coccus* graminis corpore roseo. *Geoff. Inf.* 1. 512. 2.  
 tab. 10. fig. 5.  
 Habitat ad Graminum radices.
- C. Crataegi oxyacanthae. *Syst. Ent.* 744. 12. *Linn. Crataegi.*  
*Syst. Nat.* 2. 742. 21.  
*Reaum. Inf.* 4. tab. 6. fig. 11. 12.  
 Habitat in Crataego oxyacantha.
- C. Serratulae aruenis. *Syst. Ent.* 744. 13. \* *Serratulae.*  
 Habitat in Serratula arueni Angliae.
- C. vitis viniferae. *Syst. Ent.* 744. 14. *Linn. Syst. Nat. Vitis.*  
 2. 741. 16.  
*Chermes* Vitis oblongus. *Geoff. Inf.* 1. 506. 6.  
*Aët. Bonon.* 2. 279. tab. 284.  
*Reaum. Inf.* 4. tab. 6. fig. 5 — 7.  
 Habitat in ramis vitis viniferae.
- C. Zosteræ marinae. *Gen. Inf. Mant.* 304. \* *Zosteræ.*  
 Habitat in Zostera marina maris Balthici, captus d. 18.  
 Maii.
- C. radici Scleranthi perennis. *Syst. Ent.* 744. 15. *polonicus.*  
*Linn. Syst. Nat.* 2. 741. 17. *Fn. Sv.* 1023.  
*Chermes* radicum purpureus. *Geoff. Inf.* 1. 504. 1.  
*Coccus* tinctorius radicum. *Breyn. Aët. Phys. med.* 6.  
 3. *App.* 5. tab. 1.  
*Cochinella* germanica. *Frisch. Inf.* 5. 6. tab. 2.  
*Reaum. Inf.* 4. 1. 143.  
 Habitat in Scleranthi perennis radicibus.
- C. Cacti coccinelliferi. *Syst. Ent.* 744. 16. *Linn. Syst. Cacti.*  
*Nat.* 2. 742. 22.  
*Coccinella* alis destituta, corpore rugoso. *Brown. Iam.*  
 435.  
*Coccus* Cacti cacti coccinelliferi. *Degeer Inf.* 6. 447. 1.  
 tab. 80. fig. 12. 13. 14.  
*Sloan. Iam.* 2. 153. *praef.* tab. 9.  
*Pettv. Gazoph.* tab. 1. fig. 5.  
*Reaum.*

396 CLASS. VII. RYNGOTA. Thrips.

*Reaum. Inf.* 4. tab. 7. fig. 11. 12.

*Rausch. Hist. nat. de la Cochinelle* tab. 1. fig. 1. 2. 3. 4. 5.

Habitat in Cactis Opuntiis Americes.

Corpus totum rugosum, tinctorium.

169. THIRIPS.

*Physapus.* 1. *T. elytris glaucis, corpore atro.* *Syst. Ent.* 745. 1. *Linn. Syst. Nat.* 2. 743. 2. *Fn. Sv.* 1027.

*Physapus ater, alis albis.* *Degeer Aët. Holm.* 1744. 3. tab. 4. fig. 4.

*Thrips atra, alis albis, antennis sexnodiis.* *Degeer Inf.* 3. 6. 1. tab. 1. fig. 1.

*Geoff. Inf.* 1. 385. 2.

*Schaeff. Elem.* tab. 127.

Habitat frequens in Europae floribus compositis agilis, cursitans.

Loti corniculatae flores clausos tumidosque reddit, spicas fecalis inanit. *Linn.*

*Juniperina.* 2. *T. elytris niueis, corpore fusco.* *Syst. Ent.* 745. 2. *Linn. Syst. Nat.* 2. 743. 4. *Fn. Sv.* 1029.

*Thrips griseo fusca, oculis nigris, alis albescentibus.* *Degeer Inf.* 3. 10. 2. tab. 1. fig. 5.

*Physapus fuscus, alis albicantibus.* *Degeer Aët. Holm.* 1744. 6. tab. 1. fig. 2.

Habitat in Juniperi Gallis.

*Vlmi.* 3. *T. nigra, alis niueis ciliatis, ano acuminato.*

*Thrips Corticis nigra, alis hyalinis niueis, barbis longissimis, antennis octonodiis.* *Degeer Inf.* 3. 11. 3. tab. 1. fig. 8 — 13.

*Thrips elytris albidis, corpore nigra abdominali fca.* *Geoff. Inf.* 1. 384. 1. tab. 7. fig. 6.

Habitat in Vlmi corticibus gregaria.

*minutissima.* 4. *T. elytris corporeque glaucis, oculis fuscis.* *Gen. Inf. Mant.* 304. *Linn. Syst. Nat.* 2. 743. 3. *Fn. Sv.* 1028.

Habitat in Europae floribus.

**CLASS. VII. RYNGOTA. Thrips. 397**

**T. flava, elytris albidis.**

*urticae.*

*Schrank. Beytr. 31. tab. 1. fig. 25. 26.*

Habitat in pagina inferiori foliorum *Vrticae*, *Vitis*,  
*Coryli* aliorumque solitaria.

5. **T. elytris albo nigroque fasciatis, corpore fusco. Syst. fasciata.**  
*Ent. 745. 3. Linn. Syst. Nat. 2. 743. 5. Fn. Sv.*  
1036.

*Thrips elytris albis nigrisque fasciis. Geoff. Inf. 1.*  
385. 3.

*Thrips atra, alis fasciis transuersis tribus albis, anten-*  
*nis annulo albo. Degeer Inf. 3. 18. 4.*

*Sulz. Inf. tab. 7. fig. 48. 6.*

Habitat in Europae floribus compositis.



CLASSIS VIII.  
ANTLIATA.

170. OESTRVS.

**O.** *buccatus*. 1. *O.* griseus, facie alba, nigro punctata. *Gen. Inf. Mant.* 304. \*

Habitat in Carolina australi. *Mus. Dom. Teats.*

*Bouis*. 2. *O.* thorace flauo, fascia nigra, alis albis fusco fasciatis. *Syst. Ent.* 746. 1.

*Oestrus Bouis* alis maculatis, thorace flauo, fascia fusca, abdomine flauo apice nigro. *Linn, Syst. Nat.* 2. 969. 1. *Fn. Sv.* 1730. *It. Westrog.* 202.

*Oestrus* thorace flauo, angulo nigro, alis nigra fascia pedibus pallidis. *Geoff. Inf.* 2. 456. 3.

*Oestrus Bouis* niger, alis immaculatis, thorace antice posticeque pilis griseis, abdomine antice villis griseis apiceque flauo fuluis. *Degeer Inf.* 6. 297. 2. tab. 15. fig. 22.

*Derh. Theolog.* 1. 8. l. 6. n. 11.

*Vallin. nat.* 1. tab. 28.

*Reaum. Inf.* 4. tab. 36 — 38.

*Frisch. Inf.* 5. 25. tab. 7.

*Sulz. Inf.* tab. 20. fig. 127.

*Schaeff. Elem.* tab. 91.

*Raj. Inf.* 271.

Habitat intra bouum dorsum, vlcerâ cacoethica causans.

*Tarandi*. 3. *O.* alis immaculatis, thorace flauo, fascia nigra, abdomine fuluo, apice nigro. *Syst. Ent.* 746. 2. *Linn. Syst. Nat.* 2. 969. 2. *Fn. Sv.* 1731.

*Linn. Flor. Lappon.* 360. 517.

*Ad. Holmenf.* 1739. tab. 3. fig. 5. 6.

Habitat intra Cerui Tarandi dorsum, facie ipsis lethalis.

CLASS. VIII. ANTLIATA. Tipula. 399

4. O. alis immaculatis, thorace ferrugineo, abdomine nigro, pilis flavis. *Syst. Ent.* 746. 3. *Linn. Syst. Nat.* 2. 969. 3. *Fn. Sv.* 1722. *nasalis.*  
Habitat in Equorum fauce, per nares intrans.
5. O. alis immaculatis, abdomine nigro, basi albo apiceque fuluo. *Syst. Ent.* 747. 4. *haemorrhoidalis.*  
*Oestrus haemorrhoidalis* alis immaculatis, thorace nigro, scutello pallido, abdomine nigro basi albo apiceque fuluo. *Linn. Syst. Nat.* 2. 970. 4. *Fn. Sv.* 1733.  
*Oestrus villosus* pallide flavescens, abdominis medio cingulo nigro, apice fuluo. *Geoff. Inf.* 2. 455. 1.  
*Oestrus intestinalis* fuluo griseus, thorace fascia transversa nigra, alis maculatis, pedibus fulvis. *Degeer Inf.* 6. 291. 1. tab. 15. fig. 13.  
*Musca bipennis*, *Oestrus dicta*, alis membranaceis punctis crebris nigrioribus velut adpersis. *Raj. Inf.* 271. *Reaum. Inf.* 4. tab. 35. fig. 3 — 5.  
*Frisch. Inf.* 5. tab. 7.  
Habitat in Equorum intestino recto per anum intrans. Thorax antice pallidus, postice niger.
6. O. alis subpunctatis, abdomine albo nigroque versicolore. *Syst. Ent.* 747. 5. *Linn. Syst. Nat.* 2. 970. 5. *Ouis.*  
*Fn. Sv.* 1734.  
*Oestrus cinereus* nigro maculatus et punctatus. *Geoff. Inf.* 2. 456. 2. tab. 17. fig. 1.  
*Schreb. Inf.* 15. 12.  
*Vallijn. nat.* 1. 224. tab. 27.  
*Reaum. Inf.* 4. tab. 35. fig. 21 — 24.  
Habitat in Ouium sinu frontis.

171. T I P U L A.

\* Alis patentibus.

1. T. antennis pectinatis, alis macula nigra, abdomine basi rufo, medio flavo fasciato, apice atro. *Syst. Ent.* 747. 1. *pectinis cornis.*

*Tipula*

400 CLASS. VIII. ANTLIATA. Tipula.

*Tipula pectinicornis* antennis pectinatis, alis macula nigra, thorace flavescente. *Linn. Syst. Nat.* 2. 970. 1. *Fn. Sv.* 1735.

*Tipula nigro crocea* flavo crocea; antennis pectinatis, capite thoraceque supra nigris, abdomine maculis nigris, alis puncto fusco. *Degeer Inf.* 6. 400. 24. tab. 25. fig. 3.

*Reaum. Inf.* 5. tab. 1. fig. 11 — 16.

*Schaeff. Elem.* tab. 13. fig. 8.

— — — tab. 129. fig. 3.

— — *Icon.* tab. 106. fig. 5. 6.

*Raj. Inf.* 73. 12.

*Scop. carn.* 839.

Habitat in Europae humidis.

*riuosa.* 2. *T.* alis hyalinis, riulis maculaque niuea. *Syst. Ent.* 748. 2. *Linn. Syst. Nat.* 2. 971. 2. *Fn. Sv.* 1738.

*Tipula* alis albis fusco variegatis. *Geoff. Inf.* 2. 554. 1.

*Tipula* griseo fusca carneo mixta, antennis setacis brevioribus, alis albis, riulis longitudinalibus fuscis. *Degeer Inf.* 6. 341. 2. tab. 19. fig. 1.

*Tipula* maxima, alis maioribus fusco et albo variegatis. *Raj. Inf.* 72. 2.

*Tipula riuosa.* *Scop. carn.* 846.

*Alf. Holm.* 1739. tab. 9. fig. 8.

*Sulz. Inf.* tab. 20. fig. 128.

Habitat in Europa frequens.

*sinuata.* 3. *T.* alis albis, margine sinuato maculisque fuscis, corpore cinereo, pedibus ferrugineis. *Syst. Ent.* 748. 3. \* *It. Norwag.* d. 8. Ian.

Habitat in Europa boreali.

*maculata.* 4. *T.* alis fusco venosis, margine maculisque quatuor fuscis, abdomine supra flavescente. *Syst. Ent.* 748. 4. *Linn. Syst. Nat.* 2. 971. 3. *Fn. Sv.* 1736.

*Tipula* flavo fulva nigro maculata, antennis filiformibus, pedibus rufis, femoribus maculis binis nigris, alis fusco maculatis. *Degeer Inf.* 6. 343. 5.

Habitat in Europae pratis.

**CLASS. VIII. ANTLIATA. Tipula. 401**

**T. alis macula fusca, abdomine atro, fasciis fulvis. Syst. crocata.**

*Ent.* 748. 5. *Linn. Syst. Nat.* 2. 971. 4. *Fn. Sv.* 1739.

**Tipula corpore nigro fulvo flavoque variegato. Geoff.**

*Inf.* 2. 553. 7. tab. 19. fig. 1.

**Tipula flavo fasciata aterrima, abdomine fasciis transversis flavis, antennis filiformibus, pedibus punctoque alarum fuscis. Degeer Inf.** 6. 349. 10.

**Tipula elegans, dorso et scapulis nigris, ventre croceo, alis macula fusca notatis, laevis et splendens. Raj.**

*Inf.* 72. 4.

*Schaeff. Icon.* tab. 126. fig. 4.

*Scop. carn.* 845.

Habitat in Europa boreali frequens.

**T. alis hyalinis, maculis sparsis obsoletis. Syst. Ent. hortorum.**  
748. 6. *Linn. Syst. Nat.* 2. 971. 6. *Fn. Sv.* 1725.

**Tipula griseo fusca, antennis filiformibus rufis, alis fusco alboque maculatis. Degeer Inf.** 6. 344. 6.

*Schaeff. Icon.* tab. 15. fig. 3. 4.

*Raj. Inf.* 72. 6.

*Alb. Inf.* tab. 61.

Habitat in Europae plantis oleraceis.

**T. alis hyalinis, costa marginali fusca. Syst. Ent. 749. oleraceis.**  
7. *Linn. Syst. Nat.* 2. 971. 5. *Fn. Sv.* 1740.

**Tipula alis exalbidis, linea marginali fusca. Geoff. Inf.**  
2. 555. 3.

**Tipula pratensis griseo fusca, antennis filiformibus, alis pallide fuscis immaculatis, costa marginali fusca.**

*Degeer Inf.* 6. 339. 1. tab. 18. fig. 12. 13.

*Reaum. Inf.* 4. tab. 11. fig. 7.

— — — 5. tab. 2. fig. 11.

— — — tab. 3. fig. 1. 2. 10.

*Schaeff. Icon.* tab. 15. fig. 6.

*Scop. carn.* 854.

*Göed. Inf.* 2. tab. 44.

*Bradl. nat.* tab. 25. fig. 3.

*Frisch. Inf.* 4. tab. 12.

*Leuwenh. nat.* 1693. fig. 5.

— — — 1704. fig. 4.

Habitat ad radices plantarum, oleribus infesta,

*Fabr. Sp. Inf. T. II.*

C c

g. T.



402 CLASS. VIII. ANTLIATA. Tipula.

*tricolor.* 8. T. alis albidis, margine exteriori neruoque apice  
do fuscis. *Syst. Ent.* 749. 8. \*

Habitat in America boreali. *Don. Lewin.*

*triangul- 9. T. alis dimidiato fuscis, macula triangulari alba.*  
*laris. Ent.* 749. 9. \*

Habitat in Scotia. *D. Walker.*

*variega- 10. T. nigra, abdominis basi lateribusque rubris,*  
*ta. maculatis. Syst. Ent.* 749. 10. \*

*Tipula variegata* alis hyalinis, macula pallida ob-  
ta, corpore nigro luteoque variegato. *Linn.*  
*Nat.* 2. 971. 7. *Fn. Sv.* 1742.

*Tipula nigra*, antennis filiformibus, thorace abd-  
nisque lateribus luteo maculatis, alis puncto v-  
que latis fuscis. *Degeer Inf.* 6. 346. 8.

*Tipula* alis lutescentibus cum macula fusca corpor-  
pedibus bicoloribus ex nigro croceo. *Raj. Inf.*  
*Schaeff. Icon.* tab. 15. fig. 7.

Habitat in Europae hortis.

*contami- 11. T. atra, alis albis, fasciis duabus punctoque ni*  
*nata. Syst. Ent.* 749. 11.

*Tipula contaminata* alis nigro maculatis, corpore  
gro. *Linn. Syst. Nat.* 2. 972. 8. *Fn. Sv.* 1743.

*Tipula nigra* pedibus abdominisque maculis vtri-  
liuidis, alis nigro maculatis. *Geoff. Inf.* 2. 558. 6.

Habitat in Europae humidis.

*lunata. 12. T. alis cinerascens, lunula marginali alba. S*  
*Ent.* 750. 12. *Linn. Syst. Nat.* 2. 972. 9. *Fn.*  
1744.

*Tipula* alis cinerascens, lunula alba margin-  
*Geoff. Inf.* 2. 555. 4.

*Tipula* flavo grisea, antennis filiformibus, alis pall-  
fuscis, puncto fusco fasciaque obliqua alba. *Deg.*  
*Inf.* 6. 342. 3.

*Tipula septimae* similis, paullo tamen maior, oculis  
ridescens. *Raj. Inf.* 73. 10.

*Reaum. Inf.* 5. tab. 2. fig. 11.

Habitat in Europae pratis.

**CLASS. VIII. ANTLIATA. Tipula. 403**

**T. thorace variegato, abdomine fusco, lateribus flavo** *pratenfis.*  
maculatis, fronte fulva. *Syst. Ent.* 750. 19.

**Tipula pratenfis** alis subcinereis, vasis fuscis punctoque  
marginali obscuro, fronte fulva. *Linn. Syst. Nat.* 2.  
972. 10. *Fn. Sv.* 1745.

*Frisch. Inf.* 4. tab. 12.

**Habitat** in Europae pratis, graminum radices destruens.

**T. flavescens, dorso fusco, alis hyalinis, macula mar-** *dorsalis.*  
ginali nigra.

**Habitat** in Germania. *Dom. de Hattorf.* In Italia. *D.*  
*Allioni.*

**Statura** et magnitudo praecedentis. **Antennae** porre-  
ctae, obtuse ferratae, nigrae. **Thorax** et abdomen  
flavescentia dorso omni fusco. **Alae** hyalinae macu-  
la ordinaria marginali nigra.

**T. fusco cinerea, alis albis, costa nervisque nigris.** *plumbea.*  
**Habitat** in Italia. *D. Allioni.*

**Corpus** medium totum cinereo fuscum s. plumbeum.  
**Antennae** nigrae. **Alae** albae costa exteriori nervis-  
que nigris. **Pedes** nigri femoribus basi testaceis.

**T. alis hyalinis, puncto marginali fusco, abdomine** *cornicina,*  
flavo, lineis tribus fuscis. *Syst. Ent.* 750. 14. *Linn.*  
*Syst. Nat.* 2. 972. 12. *Fn. Sv.* 1747.

**Tipula** alis subfuscis, thorace flavo, characteribus ni-  
gris, abdomine luteo, punctorum nigrorum lineis  
tribus longitudinalibus. *Geoff. Inf.* 2. 556. 5.

**Tipula flavo maculata** antennis filiformibus, capite tho-  
raceque citreo nigro maculato, abdomine flavo te-  
staceo, lineis nigro punctatis. *Degeer Inf.* 6. 347. 9.  
tab. 19. fig. 2. 3.

*Roef. Inf.* 2. *Musc.* tab. 1.

*Raj. Inf.* 73. 7.

**Habitat** in Europa, plantarum radices exedens,

**T. alis glaucis, puncto marginali corporeque atro, ab-** *atrata.*  
dominis segmento primo pedibusque rufis. *Syst.*  
*Ent.* 759. 15. *Linn. Syst. Nat.* 2. 972. 14. *Fn. Sv.*  
1749.

404 CLASS. VIII. ANTLIATA. Tipula.

*Tipula ichneumonea* atra, antennis filiformibus abdomine antice pedibusque rufis, alis puncto nigro. *Degeer Inf.* 6. 350. 11. tab. 19. fig. 10.

Habitat in Europa.

Femina abdomine recuruo, subulato, acutissimo.

*nigra.* 18. *T. alis fuscis, corpore atro. Syst. Ent.* 750. 16. *Linn. Syst. Nat.* 2. 972. 13. *Fn. Sv.* 1748.

*Geoff. Inf.* 2. 559. 10.

Habitat in Europae plantis.

*clawipes.* 19. *T. fusca, tarsis albo annulatis, medio ouato incrassatis.*

Habitat in America boreali. *Mus. Dom. Teats.*

Statura singularis. Caput nobis deest. Thorax fusco postice cinereus. Abdomen cylindricum, fuscum. Alae albae. Pedes elongati, nigri, tibiis annulo bascos albo. Tarsi itidem annulo bascos albo, in medio ouati incrassati, nigri, tunc albi apice nigri.

*bimaculata.* 20. *T. alis hyalinis, maculis duabus fuscis, abdominis medio maculato ferrugineo, antennis plumosis. Syst. Ent.* 751. 17. *Linn. Syst. Nat.* 2. 972. 15. *Fn. Sv.* 1750.

*Tipula nigra, antennis pectinatis, abdominis lateribus pedibusque flauis, alis maculis duabus fuscis. Deger Inf.* 6. 402. 25.

*Reaum. Inf.* 5. tab. 4. fig. 1. 2.

*Schaeff. Icon.* tab. 3. fig. 5. 6.

Habitat in Europa.

Mas abdomine immaculato.

*annulata.* 21. *T. alis fusco variegatis, femoribus annulo albo. Syst. Ent.* 751. 18. *Linn. Syst. Nat.* 2. 973. 16. *Fn. Sv.* 1752.

*Reaum. Inf.* 5. tab. 4. fig. 1. 2.

Habitat in Europa.

*ocellaris.* 22. *T. alis albidis, maculis ocellaribus plurimis nigricantibus. Syst. Ent.* 751. 19. *Linn. Syst. Nat.* 2. 973. 17. *Fn. Sv.* 1751.

Habitat in Europa boreali haud infrequens.

23. *T. alis albidis, maculis tribus fuscis, corpore cinereo cinerea.*  
 immaculato. *R. Norwag.* d. 29. Jul. \*  
 Habitat in Norwagia, capt. d. 29. Jul.
24. *T. alis albidis, fasciis quatuor fuscis flexuosis, abdo- fasciata.*  
 mine flavescente. *Syst. Ent.* 751. 20. *Linn. Syst. Nat.*  
 2. 973. 18.  
 Habitat in Sueciae paludosis.
25. *T. alis albis, punctis tribus marginalibus fuscis, tho- 6puncta-*  
 race compresso fuluo, linea dorsali nigra. *ta.*  
 Habitat in Italia. *D. Allioni.*
- Statura omnino *T. 3punctatae*. Caput nigrum postice  
 valde attenuatum. Thorax compressus, eleuatus,  
 fuluus linea dorsali atra. Abdomen fuscum ano ru-  
 fo. Alae albae punctis tribus marginalibus distinctis  
 fuscis. Pedes fusci femoribus basi testaceis.
26. *T. alis hyalinis, punctis tribus marginalibus fuscis, tripuncta-*  
 corpore flauo. *ta.*  
 Habitat in Italia. *D. Allioni.*
- Statura et magnitudo omnino *T. flavescentis*, distincta  
 tamen punctis tribus alarum marginalibus fuscis.
27. *T. alis immaculatis, corpore flauo. Syst. Ent.* 751. 21. *flavescentis.*  
*Linn. Syst. Nat.* 2. 973. 19. *Fn. Sv.* 1753.  
*Tipula* flaua oculis nigris. *Geoff. Inf.* 2. 557. 7.  
 Habitat in Europae campis.
28. *T. alis hyalinis nitentibus, corpore cinereo fusco. regelatio-*  
*Syst. Ent.* 751. 22. *Linn. Syst. Nat.* 2. 973. 21. *Fn. nis.*  
*Sv.* 1754.  
 Habitat in Europae terrestribus.
29. *T. alis hyalinis, margine tenuiori recuruato, corpo- replicata.*  
 re fusco, antennis simplicibus. *Linn. Syst. Nat.* 2.  
 973. 22. *Fn. Sv.* 1755.  
*Tipula* fusca, antennis filiformibus, alis pallide fuscis,  
 margine tenuiori recuruato. *Deger Inf.* 6. 351. 12.  
 tab. 20. fig. 15.  
 Habitat in Europae borealis aquis,

406 CLASS. VIII. ANTLIATA. Tipula.

- monoptera.* 30. T. nigra, pedibus palpisque pallidis. *Syst. Ent.* 75. 23. *Linna. Syst. Nat.* 2. 973. 23. *Fa. Sv.* 1756.  
Habitat in Europa boreali.

\*\* Alis incumbentibus.

- plumosa.* 31. T. thorace virecente, alis albis, puncto fusco. *Syst. Ent.* 751. 24. *Linna. Syst. Nat.* 2. 974. 26. *Fa. Sv.* 1758.

*Tipula fusca*, thorace virecente, alis pellucidis, puncto nigro. *Geoff. Inf.* 2. 560. 16.

*Tipula annularis* griseo fusca, antennis filiformibus maris plumosis, abdomine fasciis nigris, alis hyalinis puncto nigro. *Degeer Inf.* 6. 379. 18. tab. 19. fig. 14. 15.

*Frisch. Inf.* tab. 11. fig. 12.

*Goed. Inf.* 3. tab. 10.

*Reaum. Inf.* 5. tab. 5. fig. 10.

— — — 4. tab. 14. fig. 12.

Habitat in Europae paludosis.

- motitatrix.* 32. T. pedibus anticis maximis motatoriis, annulo albo. *Syst. Ent.* 752. *Linna. Syst. Nat.* 2. 974. 29. *Fa. Sv.* 1700.

*Tipula* pedibus anticis maximis antenniformibus motatoribus, annulo albo. *Geoff. Inf.* 2. 562. 18.

*Culex luteo viridis*, pedibus antenniformibus. *Frisch. Inf.* 11. 7. tab. 13.

Habitat in Europa frequens.

- fasciculata.* 33. T. pedibus anticis maximis motatoriis nigra, abdominis lateribus ferrugineo maculatis.

Habitat in Germania. *Dom. de Hatterf.*

Parua. Antennae fuscae fasciculato plumosae. Corpus nigrum, abdominis lateribus ferrugineo maculatis. Pedes nigri anticis longioribus basi ferrugineis. Alae albae puncto paruo fusco.

- tendens.* 34. T. ferruginea, alis albis immaculatis, pedibus anticis longissimis pallidis. *Syst. Ent.* 752. 26. \*

Habitat in Daniae paludosis.

CLASS. VIII. ANTLIATA. Tipula. 407

15. T. pedibus anticis longissimis motoriiis nigris, annu- tremula.  
lis albis. *Syst. Ent.* 752. 27. *Linn. Syst. Nat.* 2. 975.  
31. *Fn. Sv.* 1762.

Habitat in Sueciae paludosis.

16. T. pedibus anticis motoriiis omnibus pallidis, alis flexilis.  
fascia fuscescente. *Syst. Ent.* 752. 28. *Linn. Syst.*  
*Nat.* 2. 975. 32.

Tipula pedibus anticis maximis antenniformibus mota-  
toriiis immaculatis. *Geoff. Inf.* 2. 563. 19.

Habitat in Europae aquosis.

17. T. pedibus anticis maximis motoriiis, apice albis. vibrato-  
ria.  
*Syst. Ent.* 752. 29. *Linn. Syst. Nat.* 2. 975. 30. *Fn.*  
*Sv.* 1761.

Habitat in Europae paludosis.

18. T. nigra, abdominis segmentis apice albis, alis pun- fistica,  
cto fusco.

Habitat in Germania. *Dom. de Hattorf.*

Affinis T. cinctae. Caput paruum, nigrum. Thorax  
niger cinereo lineatus. Abdomen nigrum segmento-  
rum apicibus elevatis, niueis. Alae albae puncto  
unico centrali fusco. Pedes alba nigroque varii.

19. T. pallida pilosa, pedibus nigro punctatis. pallida.

Habitat in Germania. *Dom. de Hattorf.*

Statura et magnitudo praecedentium. Antennae fasci-  
culatae pallidae. Thorax pallidus, immaculatus.  
Abdomen pilosum, pallidum segmentorum apicibus  
obscurioribus. Alae pallidae, immaculatae. Pedes  
pallidi punctis numerosissimis nigris.

20. T. pallida, abdomine nigro annulato, alis maculatis. flabelli-  
cornis.

Habitat in Germania. *Dom. de Hattorf.*

Affinis praecedenti, at alae maculatae et pedes omnino  
immaculati.

21. T. liuida, alis maculis tribus marginalibus nigris, ab- cincta.  
domine nigro albo annulato. *Syst. Ent.* 752. 30. *Linn.*  
*Syst. Nat.* 2. 974. 28.

Habitat in Suecia.

408 CLASS. VIII. ANTLIATA. Tipula.

- Arundineti.* 42. *T. albida*, antennis villosis, oculis atris. *Syst. Ent.* 752. 31. *Linn. Syst. Nat.* 2. 974. 24. *Fn. Sv.* 1757.  
Habitat in Arundinetis choreas agens.
- littoralis.* 43. *T. virescens*, alis immaculatis, pedibus anticis longifimis. *Syst. Ent.* 752. 32. *Linn. Syst. Nat.* 2. 974. 27. *Fn. Sv.* 1759.  
*Tipula viridis* alis albis immaculatis. *Geoff. Inf.* 2. 564. 21.  
*Reaum. Inf.* 5. tab. 6. fig. 12. 13. 14.  
Habitat in Europae maritima.
- monilis.* 44. *T. pedibus albis*, annulis nouem nigris, alis albo cinereoque variis. *Syst. Ent.* 753. 33. *Linn. Syst. Nat.* 2. 975. 33. *Fn. Sv.* 1763.  
*Tipula* pedibus albis, annulis nigris, alis albis cinereo maculatis. *Geoff. Inf.* 2. 564. 20.  
Habitat in Europae hortis.
- zonata.* 45. *T. pallida*, alis fasciis duabus punctisque tribus fuscis, femoribus angulo fusco. *Syst. Ent.* 753. 34.\*  
Habitat Oxoniae.
- macrocephala.* 46. *T. virescens*, oculis thoracisque tergo nigris. *Syst. Ent.* 753. 35.\* *Linn. Syst. Nat.* 2. 975. 36. *Fn. Sv.* 1764.  
Habitat ad Europae paludes et littora humida.
- viridula.* 47. *T. viridis*, antennis verticillato pilosis, pedibus pallidis. *Linn. Syst. Nat.* 2. 975. 35.  
Habitat in Europa boreali.  
Thorax antice parum ferrugineus. Pedum articuli nigri.
- pufilla.* 48. *T. viridis*, thoracis tergo maculis tribus nigris, antennis maris plumosis. *Syst. Ent.* 753. 36. *Linn. Syst. Nat.* 2. 975. 37.  
Habitat ad Europae fossulas, choreas agens.
- breuicornis.* 49. *T. nigra* glabra, alis margine nigricantibus, abdomine fusco, tibiis anticis spinosis. *Syst. Ent.* 753. 37.

*Tipula*

CLASS. VIII. ANTLIATA: Tipula, 409

*Tipula brevicornis* nigra glabra, alis margine nigricantibus, tibiis anticis spina terminatis. *Linn. Syst. Nat.* 2. 976. 42. *Fn. Sv.* 1766.

*Tipula flavicauda* nigra, antennis breuibus subulatis abdomine feminae flavo, alis obscuris, tibiis anticis spina terminatis. *Degeer Inf.* 6. 429. 35.

β. *Tipula ferrugata* atra, glabra, alis fuscis, abdomine fusco ferrugineo. *Linn. Syst. Nat.* 2. 976. 49.

Habitat in Europae hortis, umbrosis.

Quiescit alis expansis abdomineque extenso.

Variat tibiis anticis posticisque ferrugineis.

50. T. alis hyalinis, margine exteriori nigro. *Syst. Ent. hortulana* 753. 38. na.

*Tipula hortulana* alis albis, margine exteriori nigro, thorace abdomineque rubro. *Linn. Syst. Nat.* 2. 977. 46.

*Bibio* alis albis, margine exteriori nigro, thorace abdomineque rubris. *Geoff. Inf.* 2. 571. 3.

*Reaum. Inf.* 5. tab. 7. fig. 7 — 10.

β. *Tipula Marci* nigra glabra, alis nigricantibus, femoribus anticis introrsum sulcatis. *Linn. Syst. Nat.* 2. 976. 38. *Fn. Sv.* 1765.

*Tipula Marci* atra tota, antennis breuibus subulatis, alis margine exteriori nigro, tibiis anticis spina terminatis. *Degeer Inf.* 6. 428. 33.

*Bibio ater* hirsutus, alis albis, margine exteriori nigro. *Geoff. Inf.* 2. 510. 2.

*Schaeff. Icon.* tab. 15. fig. 1. 2.

Habitat in Asparagi et Pomonae floribus.

Variat thorace abdomineque rubris et nigris, quas saepius copula iunctas vidimus.

51. T. nigra glabra, alis albis, puncto nigro, antennis breuibus, pedibus nigris. *Syst. Ent. Iohannis.* 754. 39.

*Tipula Iohannis* atra glabra, alis puncto nigro, tibiis pallidis, posticis clauatis. *Linn. Syst. Nat.* 2. 976. 41.

*Tipula atra*, antennis breuibus subulatis, pedibus feminae rufis, alis puncto nigro, tibiis anticis spina terminatis. *Degeer Inf.* 6. 425. 32. tab. 27. fig. 12. 13.

Habitat in Europae umbrosis.



410 CLASS. VIII. ANTIATA. Tipula.

*Pomonae.* 52. *T. nigra* glabra, alis hyalinis, puncto nigro, femoribus ferrugineis. *Syst. Ent.* 754. 40.\*

Habitat in *Pomonae* floribus Angliae, Norwagiae.

*ruficollis.* 53. *T. atra* glabra, thorace rufo.

Habitat ad Promontorium bon. sp. *Mus. Dom. Banks.*

Statura praecedentium. Corpus totum atrum solo thorace gibbo, rufo. Alae ipsae fuscae ad marginem crassiores obscuriores.

*Thomas.* 54. *T. atra* glabra, alis nigris, abdominis lateribus linea crocea. *Syst. Ent.* 754. 41. *Linn. Syst. Nat.* 2. 976. 39.

Habitat Vpsaliae.

*febrilis.* 55. *T. atra* oblonga hirta, alis costa nigricante. *Syst. Ent.* 754. 42. *Linn. Syst. Nat.* 2. 976. 44. *Fn. Sv.* 1761.

*Bibio* ater hirsutus, alis albis, margine exteriori nigro. *Geoff. Inf.* 2. 576. 2.

*Goed. Inf.* 1. tab. 6.

*Petiv. Gazoph.* tab. 14. fig. 4.

Habitat in Europae locis suffocatis.

*rufipes.* 56. *T. atra* hirta, pedibus ferrugineis, posticis elongatis.

Habitat in Insula Terre neuve. *Mus. Dom. Banks.*

Statura omnino praecedentium. Corpus nigrum hirtum. Orbita oculorum ferruginea. Abdomen dorso plano. Pedes omnes ferruginei posticis elongatis femoribus clauatis tibiisque apice compressis. Alae obscurae.

*fuscata.* 57. *T. thorace* lineato, abdomine fusco, alis macula marginali nigra, puncto cinereo. *Syst. Ent.* 755. 44.

*Musca fuscata* nigricans, alis antice albo nigroque variis. *Linn. Fn. Sv.* 2310.

Habitat in Suecia.

*forcipata.* 58. *T. abdomine* cylindrico atro, alis fusco hyalinis, ano appendiculato. *Syst. Ent.* 755. 45.\*

Habitat in Anglia, capt. d. 13. Maii.

*vernans.* 59. *T. cinerea*, thorace nigro lineato, alis albis fuscomaculatis. *Syst. Ent.* 755. 46.\*

Habitat

CLASS. VIII. ANTLIATA. Tipula. 411

Habitat primo vere in Daniae pratis.

Abdomen ferrugineum luteis duabus punctorum fasciis.

60. *T. cinerea*, alis albis, margine villoso. *Syst. Ent. juniperina*. 755. 47.

*Tipula juniperina* alis incumbens glaucis, margine tenuiori villoso. *Linn. Syst. Nat.* 2. 977. 51. *Fn. Sv.* 1774.

*Tipula fusca*, antennis nodosis, alis ovatis hirsutis, costis tribus. *Degeer Inf.* 6. 404. 26. tab. 25. fig. 7-13.

Habitat in Juniperi Trigloshide, quam caussat.

61. *T. hirsuta*, alis deflexis ovatis, ciliatis albo nigroque tessellatis. *Syst. Ent.* 755. 48. *Linn. Syst. Nat.* 2. 977. 48. *Fn. Sv.* 1772.

*Tipula hirsutissima* nigro cinerea, antennis nodosis, alis deflexis ovatis hirsutissimis ciliatis nigro maculatis. *Degeer Inf.* 6. 424. 31. tab. 27. fig. 10. 11.

*Bibio* alis deflexis cinereis ovato lanceolatis nebuloso maculatis. *Geoff. Inf.* 2. 572. 5.

Habitat in Arctio Lappa.

Alae apice albae.

62. *T. alis deflexis cinereis ovato lanceolatis ciliatis.* *Syst. phalaenoides*. *Ent.* 755. 49. *Linn. Syst. Nat.* 2. 977. 47. *Fn. Sv.* 1771.

*Tipula hirsutissima* cinerea, antennis nodosis, alis deflexis ovato lanceolatis hirsutissimis ciliatis immaculatis. *Degeer Inf.* 6. 422. 30. tab. 27. fig. 6.

*Bibio* alis deflexis cinereis ovato lanceolatis ciliatis immaculatis. *Geoff. Inf.* 2. 572. 4.

*Leuwenh. Epist.* 1692. Jan. 24. fig. 2. 3.

*Frisch. Inf.* 11. tab. 11.

Habitat ad Europae sterquiliniam.

63. *T. atra*, alis incumbens subciliatis. *Syst. Ent.* 755. *Perficariae*. 50. *Linn. Syst. Nat.* 2. 977. 49.

Habitat sub *Perficariae* foliis revolutis rubris.

64. *T. cinerea*, pedibus pallidis, alis maculis duabus nigricantibus. *Syst. Ent.* 756. 51. \*

Habitat *Uphalac.*

- palustris*. 65. T. pallida, capite atro, abdomine rufescente. *Syst. Ent.* 756. 52.  
*Tipula palustris* atra, abdomine rufescente. *Linn. Syst. Nat.* 2. 978. 54. *Fn. Sv.* 1775.  
 Habitat in Europae paludosis.
- longicornis*. 66. T. antennis corpore incarnato longioribus. *Syst. Ent.* 756. 53. *Linn. Syst. Nat.* 2. 978. 55. *Fn. Sv.* 1776.  
 Habitat in Europae humidis.
- sericea*. 67. T. nigra, dorso atro, thoracis lateribus caluis, halteribus flauis. *Syst. Ent.* 756. 54. *Linn. Syst. Nat.* 2. 978. 58.  
 Habitat in Suecia, choreas agens in aere, volatu flexuoso rapidoque.
- pulicaris*. 68. T. nigra, thoracis lateribus, scutello abdomineque fluis. *Syst. Ent.* 756. 55. *Linn. Syst. Nat.* 2. 979. 60.  
 Habitat ad Europae folias.
- pennicornis*. 69. T. antennis bipeclinatis, corpore atro, halteribus albis. *Syst. Ent.* 756. 56. *Linn. Syst. Nat.* 2. 979. 61. *Fn. Sv.* 1777.  
 Habitat in Aristolochiae Clematidis floribus, foecundationem promouens.

## 172. BIBIO.

- filata*. 1. B. antennis subelauatis nigra, abdominis segmento secundo lateribus pellucido, femoribus posticis ferratis. *Syst. Ent.* 756. 1.  
*Nemotelus asiloides* niger, antennis cylindricis muticis, abdomine longo cylindrico, segmento secundo rufo flauo, alis fuscis. *Degeer Inf.* 6. 204. 2. tab. 29. fig. 6.  
*Musca clauata*. *Drury Inf.* 1. tab. 44. fig. 1.  
 Habitat in America australi.
- illucens*. 2. B. abdominis nigri segmento secundo lateribus pellucidis, plantis albis. *Syst. Ent.* 757. 2. \*  
*Musca illucens* antennis filatis clauatis, alis coeruleis, corpore coeruleo, abdominis segmento secundo lateribus pellucido. *Linn. Syst. Nat.* 2. 979. 2.

Nemo

- Nemotelus illucens* niger, antennis cylindricis muticis, abdomine oblongo depresso, segmento flauo pellucido, alis fusco violaceis. *Degeer Inf.* 6. 205. 3. tab. 29. fig. 8.  
Habitat in America meridionali.
5. *B. atra*, thorace lineis duabus albis, tibiis basi ferrugineis. *Syst. Ent.* 757. 3. \* *bilineata*.  
Habitat in noua Zelandia. *Mus. Dom. Banks.*
4. *B. nigra* tomentosa, abdomine aeneo. *Gen. Inf. Mant.* 304. \* *aenea*.  
Habitat in Germania. *Dom. de Hattorf.*
5. *B. cinereo hirta*, abdomine conico, segmentis margine albis. *Syst. Ent.* 757. 4. *plebeia*.  
*Musca plebeia* antennis filatis subulatis, corpore cinereo hirta, abdominis incisuris margine albis. *Linn. Syst. Nat.* 2. 979. 1. *Fn. Sv.* 1779.  
Habitat in Europa boreali frequens.
6. *B. atra*, abdomine conico, segmentis margine albis, alis maculatis. *marginalis*.  
Habitat in Italia. *D. Allioni*.  
Statura et magnitudo omnino praecedentis, tota nigra marginibus segmentorum abdominalium solis albis. Alae maculis inprimis ad marginem crassiores nigris.
7. *B. ferrugineo hirta*, abdomine conico fuluo, ano atro. *Syst. Ent.* 757. 5. \* *nobilissima*.  
Habitat in Daniae nemoribus.
8. *B. villosa*, albo canescens, alis hyalinis. *Syst. Ent.* 757. 6. *anilis*.  
*Musca anilis* antennis filatis hirtis muticis, corpore villosa albo canescente. *Linn. Syst. Nat.* 2. 982. 20. *Fn. Sv.* 1791.  
Habitat in Europa boreali.
9. *B. fusca*, alis nigris albo variegatis, scutello piceo. *Syst. Ent.* 757. 7. \* *Nigrita*.

*Nemotelus tigrinus* antennis acutis, corpore lato hirsuto, rufescente atro albo maculato, alis albo nigro maculatis. *Degeer Inf.* 6. 206. 4. tab. 29. fig. 11.  
 Habitat in America. *Mus. Dom. Banks.* In Italia. *D. Allioni.*

*Lar.* 10. *B. hirta* fusca, abdomine vtrinque maculis duabus albis, alis nigris macula marginali apiceque albis.

Habitat in Bengal. *Mus. Dom. Banks.*

*Magna.* Thorax fuscus scutello ferrugineo. Abdomen nigricans maculis vtrinque duabus albis, altera bascos, altera in medio. Anus albicat. Alae magnae nigrae punctis aliquot obsoletis albis. Macula magna ad marginem tenuiorem apexque alba.

*Morio.* 11. *B. hirta atra*, alis nigris, apice hyalinis. *Syst. Ent.* 758. 8.

*Musca Morio* antennis filatis subulatis, corpore hirsuto atris, alis dimidiato nigris. *Linn. Syst. Nat.* 2. 981. 9. *Fn. Sv.* 1784.

*Musca atra hirsuta*, margine alarum tenuiore sinuato albicante. *Geoff. Inf.* 2. 493. 2.

*Nemotelus Morio* antennis acutis, corpore lato nigro hirsuto, lateribus villis nigris, ano fasciculis binis albis, alis dimidiato nigris albisque. *Degeer Inf.* 6. 192. 14. tab. 11. fig. 13.

*Nemotelus septimus.* *Schaeff. Icon.* tab. 76. fig. 7.

*Anthrax Morio.* *Scop. carn.* 971.

*Reaum. Inf.* 6. tab. 27. fig. 19.

*Aldr. Inf.* tab. 2. fig. 13.

Habitat in Europa.

*Maura.* 12. *B. hirta atra albo fasciata*, alis nigris, margine tenuiore sinuato hyalino. *Syst. Ent.* 758. 9.

*Musca maura* antennis filatis subulatis, corpore hirsuto atro albo fasciato, alis nigro fasciatis. *Linn. Syst. Nat.* 2. 981. 11. *Fn. Sv.* 1785.

*Nemotelus maura* antennis acutis, corpore lato nigro, hirsuto, thorace lateribus villis flavo albidis, abdomine fasciis villosis albidis, alis dimidiato nigris albisque. *Degeer Inf.* 6. 191. 13. tab. 11. fig. 11.

*Schaeff.*

*Schaeff. Icon. tab. 76. fig. 8.*

Habitat in Europa.

13. *B. hirta atra*, alis nigris, apice albis, punctis duobus, *Aethiops*,  
ano argenteo.

Habitat in Italia. *D. Allioni.*

Statura omnino praecedentis. Caput et thorax nigra  
immaculata. Alae basi nigrae albo punctatae apice  
albae punctis duobus distinctis nigris. Abdomen  
atrum albo subfasciatum anoque lacte argenteo ni-  
tido.

14. *B. hirta fusca*, abdomine albo maculato, lateribus fer- *capucina*  
rugineis, alis albo nigroque variis.

Habitat - - - - *Mus. Dom. Banks.*

Statura omnino praecedentium, at paullo minor. Ca-  
put et thorax fusca scutello ferrugineo. Abdomen  
nigrum punctis aliquot albis lateribusque ferrugi-  
neis. Alae nigrae punctis duobus bascos macula ma-  
gna transuersa marginis tenuioris apiceque albis.  
Punctum paruum album inter maculam marginis te-  
nuioris et apicis.

15. *B. fusca*, abdomine atro, segmento primo basi, vltimis *Satyris*,  
lateribus albis. *Syst. Ent. 758. 10. \**

Habitat in noua Hollandia. *Mus. Dom. Banks.*

16. *B. hirta flauescens*, alis hyalinis, costa fusca. *Syst. Hotten-*  
*Ent. 758. 11. totta.*

*Musca hottentotta* antennis filatis subulatis, corpore  
hirto lutescente, alis costa fuscis. *Linn. Syst. Nat.*  
2. 981. 13. *Fn. Sv. 1787.*

*Nemotelus hottentottus* antennis acutis, corpore lato  
nigro hirsutissimo villis viridi flauis, alis hyalinis  
costa exteriori fuscis. *Degeer Inf. 6. 190. 12, tab. 11.*  
*fig. 7.*

*Schaeff. Icon. tab. 76. fig. 6.*

Habitat in Europa boreali frequens.

17. *B. fusca*, abdomine albo strigoso, alis basi fuscis, pun- *Sylvanus*,  
cto albo. *Syst. Ent. 758. 12. \**

Habitat in noua Hollandia. *Mus. Dom. Banks.*

Abdo.

416 CLASS. VIII. ANTLIATA. Stratiomys.

Abdomen fuscum strigis aliquot obsoletis albis. Alae albae basi fuscae puncto paruo albido. Pedes nigri.

**Sabaeus.** 18. *B. hirta atra*, abdomine segmentorum marginibus albidis, alis albis, basi exteriore nigris.

Habitat in Italia. *D. Allioni.*

Magnitudo et statura *B. Hottentottae*. Corpus hirtum atrum, halteres apice terminantur puncto flavo. Abdomen segmentorum marginibus parum albicantibus. Alae albae margine exteriori ultra medium nigro. Pedes nigri.

**Lucifer.** 19. *B. hirta flavescent*, alis fuscis basi ferrugineis. *Syst. Ent.* 759. 13. \*

Habitat in Americae meridionalis insulis.

**Trogodyta.** 20. *B. hirta fusca*, alis hyalinis, macula marginali basali argentea. *Syst. Ent.* 759. 14. \*

Habitat in India. *D. Koenig.*

173. STRATIOMYS.

**Chamaeleon.** 1. *S. scutello bidentato luteo*, abdomine nigro, fasciis lateralibus luteis. *Syst. Ent.* 759. 1.

*Stratiomys fusca* thorace subhirsuto ferrugineo, abdomine glabro plano ovato, lunulis sex luteis. *Geoff. Inf.* 2. 479. 1. tab. 7. fig. 4.

*Stratiomys Chamaeleon nigra*, scutello luteo, abdomine supra maculis sex lateralibus citreo luteis. *Degeer Inf.* 6. 151. 1.

*Musca Chamaeleon*, *Linn. Syst. Nat.* 2. 979. 3. *Fa. Sv.* 1780.

*Chamaeleo*. *Goed. Inf.* 1. tab. 70.

*Tabanus aquaticus*. *Frisch. Inf.* 5. 28. tab. 10.

*Musca sellata*. *Sulz. Inf.* tab. 20. fig. 130.

*Swammerd. Bibl. nat.* tab. 42. fig. 2.

*Roef. Inf.* 2. *Musc.* tab. 5.

*Reaum. Inf.* 4. tab. 25. fig. 7.

*Schaeff. Elem.* tab. 121.

— — *Icon.* tab. 14. fig. 16.

Habitat larva in Europae aquis dulcibus, imago in floribus.

**CLASS. VIII. ANTLIATA. Stratiomys. 417**

1. S. scutello bidentato, thorace rufo, vtrinque spinoso. *ephippium.*  
*Syst. Ent. 759. 2. \**  
*Stratiomys nigra* thorace sericeo rubro vtrinque spinoso. *Geoff. Inf. 2. 480. 3.*  
*Musca ephippium. Schaeff. monogr. 1753. tab. 1.*  
Habitat in Europa.
  
3. S. scutello bidentato, abdomine atro, strigis lateralibus *microleon.*  
*albis. Syst. Ent. 760. 3.*  
*Stratiomys microleon* thorace scutelloque aeneis, abdomine supra nigro, lineis sex lateralibus flavis. *Degeer Inf. 6. 152. 2. tab. 9. fig. 1.*  
*Musca microlion. Linn. Syst. Nat. 2. 980. 4. Fn. Sv. 1781.*  
Habitat in Europa boreali.
  
4. S. scutello bidentato, abdomine atro, subtus strigis *frigata.*  
*albis.*  
Habitat in Italia. *D. Allioni.*  
Nimis praecedenti affinis. Antennae basi connatae, atrae. Caput et thorax cinereo pubescentia. Dentes scutelli flavi. Abdomen atrum, supra vix maculatum, subtus vero strigis tribus albis.
  
5. S. scutello bidentato, abdomine supra atro, segmento- *pallipes.*  
rum marginibus vtrinque albidis.  
Habitat in America boreali. *Mus. Dom. Blomfield.*  
Medius. Caput nigrum linea frontali flavescente. Thorax fuscus scutelli margine ferrugineo dentibus duobus albis. Abdomen depressum, supra atrum marginibus segmentorum latere pallidis, subtus pallidum. Pedes pallidi.
  
6. S. scutello bidentato atra, abdomine subtus rufo. *Syst. tigrina.*  
*Ent. 760. 4. \**  
Habitat in Daniae aquis.
  
7. S. scutello bidentato nigro, abdomine viridi nigro an- *hydroleon.*  
gulato. *Syst. Ent. 760. 5.*  
*Stratiomys fusca*, abdomine viridi, fascia longitudinali nigra. *Geoff. Inf. 2. 481. 4.*  
*Fabr. Spec. Inf. T. II. D d Stra-*



418 CLASS. VIII. ANTLIATA. *Stratiomys*.

- Stratiomys hydroleon* thorace aeneo, abdomine viridi, fascia lata longitudinali vndata nigra, pedibus flauis. *Degeer Inf.* 6. 154. 3. tab. 9. fig. 4.  
*Musca hydroleon.* *Linn. Syst. Nat.* 2. 980. 5. *Fa. Sv.* 1782.  
*Schaeff. Icon.* tab. 14. fig. 14.  
Habitat in Europae aquis.
- viridula.* 8. *S.* scutello bidentato, viridis, abdomine flauescens, macula apicis atra. *Syst. Ent.* 760. 6. \*  
Habitat Hafniae.
- trilinea-* 9. *S.* scutello bidentato, corpore viridi, thorace lineis, ab-  
*ta.* domine fasciis nigris. *Syst. Ent.* 760. 7.  
*Stratiomys luteo* virefcens, thorace lineis tribus longitudinalibus, abdomine tribus tranauerfis arcuatis nigris. *Geoff. Inf.* 2. 402. 7.  
*Musca trilineata.* *Linn. Syst. Nat.* 2. 980. 6.  
Habitat in Europae aquis.
- hypoleon.* 10. *S.* scutello bidentato flauo, corpore nigro flauo variegato. *Syst. Ent.* 760. 7.  
*Stratiomys atra*, thorace abdomineque maculis fuscis. *Geoff. Inf.* 2. 481. 6.  
*Musca hypoleon* antennis filatis acuminato setaceis, thorace bidentato, margine striisque duabus flauis. *Linn. Syst. Nat.* 2. 980. 7.  
*Musca rara.* *Scop. carn.* 912.  
Habitat in Europa australiori.
- clauipes.* 11. *S.* scutello sexdentato atra, abdomine ferrugineo. *Syst. Ent.* 761. 9. \*  
*Stratiomys nigra*, alis nigris, femoribus abdomineque luteis. *Geoff. Inf.* 2. 403. 8.  
*Musca clauipes* antennis filatis subulatis, corpore luteo atro, abdomine testaceo, plantis posticis clauatis. *Linn. Syst. Nat.* 2. 981. 12.  
Habitat in Suecia.
- sexdentata.* 12. *S.* scutello sexdentato, thorace aeneo, abdomine nigro.  
Habitat in Anglia. *Mus. Dom. Banks.*

Statura oblonga fere Muscae cuprariae, at antennae et os huius generis. Antennae filatae basi connatae absque seta. Thorax aeneus nitidus, scutello dentibus sex elongatis nigris. Abdomen elongatum supra planum pubescens, nigrum, nitidum. Alae albiae macula marginali nigra. Pedes flavi postici elongati plantis fuscis.

13. S. scutello inermi nigra, abdominis margine tibiisque *marginalis* albidis. *Syst. Ent.* 761. 10. \*

*Stratiomys* nigra, tibiis albidis, alarum margine exteriori albo. *Geoff. Inf.* 2. 481. 5.

Habitat in Anglia. *Mus. Dom. Banks.*

24. S. scutello inermi atra, abdomine niueo, apice atro. *mutica.* *Gen. Inf. Mant.* 305. \*

Habitat in Germania. *Dom. de Hattorf.*

#### 174. RHAGIO.

1. R. cinereus, abdomine flavescente, trifariam nigro punctato, alis nebulosis. *Syst. Ent.* 761. 1. *scolopaceus.*

*Musca scolopacea.* *Linn. Syst. Nat.* 2. 982. 16. *Fn. Sv.* 1788.

*Nemotelus scolopaceus* antennis filatis setigeris, thorace nigro, abdomine flavescente supra nigro punctato, pedibus flavis, alis fusco maculatis. *Deger Inf.* 6. 162. 1. tab. 9. fig. 6.

*Reaum. Inf.* 4. tab. 10. fig. 5. 6.

Habitat in Europa boreali frequens.

2. R. cinereus, abdomine flavescente, trifariam nigro punctato, alis immaculatis, thorace unicolore. *Syst. Ent.* 761. 2. *tringarius.*

*Musca tringaria.* *Linn. Syst. Nat.* 2. 982. 18. *Fn. Sv.* 1789.

Habitat in Europae borealis nemoribus frequens.

Nimis praecedenti affinis.

3. R. cinereus, abdomine trifariam nigro punctato, alis *Vermileo.* immaculatis, thorace maculato. *Syst. Ent.* 762. 3.

*Musca Vermileo.* *Ann. Syst. Nat.* 2. 982. 17.

420 CLASS. VIII. ANTLIATA. Rhagio.

*Nemotelus Vermileo* flavus, antennis filatis setigeris, thorace striis quatuor, abdomine quinquefariam nigro punctato, alis immaculatis. *Degeer Inf.* 6. 169. 5. tab. 10. fig. 9.

*Degeer Aët. Holm.* 1752. 180. 260. tab. 5.

*Reaum. Aët. Paris.* 1753. 402. tab. 17.

Habitat in Galliae arena mobili.

*lineatus.* 4. R. thorace lineato, abdomine nigro, alis hyalinis, macula apiceque fuscis.

*Tipula lineata. Syst. Ent.* 754. 43. \*

Habitat in Anglia, capt. in *Musaeo Britann.* d. 21. Iul. Antennae cylindricae, incurvae, longiores quam in reliquis. Pedes antici elongati, nigri.

*fasciatus.* 5. R. testaceus, abdomine nigro fasciato, alis hyalinis, macula apicis fusca. *It. Norwag.* d. 13. Iul. \*

Habitat in Norwagiae nemoribus, capt. d. 13. Iul.

*tipuliformis.* 6. R. fuscus, alis albis, apice nigris. *Gen. Inf. Mant.* 306. Habitat in Germania. *Dom. de Hattorf.*

*flavipes.* 7. R. thorace ferrugineo, abdomine atro, maculis lateralibus flavis.

Habitat in Germania. *Dom. de Hattorf.*

Præcedentibus minor. Caput globosum, atrum. Thorax laevis, ferrugineus, immaculatus. Abdomen cylindricum, incurvum, atrum maculis utrinque tribus lateralibus vltimaque dorsali flavis. Pedes fuscipantis nigri. Alae hyalinae, immaculatae.

*maculatus.* 8. R. abdomine nigro, segmentorum marginibus albis, alis fusco hyalinis, maculis duabus nigris.

Habitat in Germania. *Dom. de Hattorf.*

Caput globosum, nigrum antennis filiformibus, basi laevibus. Thorax fuscus obsolete flavo lineatus. Abdomen atrum segmentorum marginibus flavis. Alae fusco hyalinae maculis duabus nigris, altera in medio, altera versus apicem. Pedes pallidi plantis nigri.

*longicornis.* 9. R. testaceus maculis abdominis lateralibus nigris, antennis longissimis.

Habitat

CLASS. VIII. ANTLIATA. Syrphus. 421

- Habitat in Germania. *Dom. de Hattorf.*  
 Singulare insectum ob antennis corpore triplo longiores setaceas et forte proprii generis mihi haud rite examinatum. Statura huius generis. Caput et thorax pallida, testacea, immaculata. Abdomen cylindricum, pallidum maculis lateralibus atris. Alae albae nigro maculae. Pedes elongati, pallidi.
10. R. cinereus, alis albis, macula marginali fusca. *Syst. diadema.*  
*Ent. 762. 4. \**  
*Musca diadema* antennis filatis apice setigeris cinerea, alis immaculatis, pedibus subferrugineis. *Linn. Syst. Nat. 2. 982. 19.*  
 Habitat in Lipsiae hortis.
11. R. ater immaculatus, alis hyalinis, macula marginali atratus. nigra.  
 Habitat in Italia. *D. Allioni.*  
 Statura praecedentis, at duplo fere maior. Pedes nigri.

175. S Y R P H V S.

\* Antennis seta plumata.

1. S. antennis plumatis tomentosus niger, abdomine postice rufo. *Syst. Ent. 762. 1. bombylans.*  
*Musca Bombylans. Linn. Syst. Nat. 2. 983. 25. Fn. Sv. 1792.*  
*Harris Inf. angl. tab. 10. fig. 3.*  
 Habitat in Europae borealis nemoribus.
2. S. antennis plumatis tomentosus niger, thorace abdominisque apice flavis. *Syst. Ent. 762. 2. mystacea.*  
*Musca mystacea. Linn. Syst. Nat. 2. 983. 26. Fn. Sv. 1793.*  
*Musca Bombylius* antennis plumatis tomentosa nigra thorace abdominisque apice rufis, alarum basi fulva.  
*Degeer Inf. 6. 133. 1. tab. 8. fig. 2.*  
*Musca Bombylium* referens, humeris rufis, cauda albicante. *Raj. Inf. 266.*  
*Sultz. Inf. tab. 20. fig. 131.*  
*Harris Inf. angl. tab. 10. fig. 1.*

422 CLASS. VIII. ANTLIATA. Syrphus.

*Schaeff. Elem. tab. 131.*

— — *Icon. tab. 10. fig. 9.*

Habitat in Europae nemoribus.

*mussitans.* 3. S. antennis plumatis tomentosus, thorace fulvo, abdomine virescente. *Gen. Inf. Mant. 306. \**

Habitat in nemoribus Kiloniensibus. *Dom. de Schædt.*

*lapponum.* 4. S. antennis plumatis tomentosus niger, scutello ferrugineo, abdomine cingulis tribus albidis interruptis. *Musca lappona. Linn. Syst. Nat. 2. 983. 27. Fa. Sc. 1794.*

*Musca* antennis plumatis nigris, tomentosa nigra, fronte flava, linea nigra, abdomine cingulis tribus flavis, apice rufo. *Degeer Inf. 6. 141. 6. tab. 8. fig. 14.*

Habitat in Europae borealis nemoribus.

*apiarius.* 5. S. antennis plumatis tomentosus, thorace antice flavo, abdominis apice albo, alis apice ferrugineis.

Habitat in Italia. *D. Allioni.*

Statura et summa affinitas S. mystacei. Antennae plumatae, frons nigra. Thorax antice hirtus flavo, postice niger. Abdomen nigrum apice hirtum, album. Alae obscurae, basi ferrugineae.

*inanis.* 6. S. antennis plumatis nudus, abdomine ferrugineo pellucido. *Syst. Ent. 763. 3. \**

Habitat in America. *Dom. v. Rohr.*

*vacuus.* 7. S. antennis plumatis nudus, thorace flavo lineato, abdomine flavo pellucido, cingulis nigris. *Syst. Ent. 763. 4. \**

Habitat in America. *Dom. v. Rohr.*

*obesus.* 8. S. antennis plumatis nudus, alis maculis duabus fuscis, corpore aeneo. *Syst. Ent. 763. 5. \**

*Musca erythrocephala* antennis plumatis pilosa, corpore breui depresso viridi aeneo, capite testaceo, oculis fuscis. *Degeer Inf. 6. 146. 2. tab. 29. fig. 2. ?*

Habitat in America meridionali.

De Synonymo *Degeer* dubito, cum noster omnino nudus et capite haud rufo.

**CLASS. VIII. ANTLIATA. Syrphus. 423**

**\*\* Antennis seta nuda.**

9. *S. antennis setariis nudus*, thorace nigro albo maculato, abdomine testaceo, fascia alba. *Syst. Ent.* 763. 6. \* *pinguis.*  
Habitat in America. *Dom. Drury.*

10. *S. antennis setariis tomentosus*, thorace lineis quatuor, abdomine fasciis tribus interruptis flavis. *Syst. Ent.* 763. 7. \* *pendulus.*

*Musca pendula.* *Linn. Syst. Nat.* 2. 984. 28. *Fn. Sv.* 1795.

*Musca antennis setariis tomentosa nigra*, capite flauo, fascia nigra, thorace lineis quatuor longitudinalibus, abdomineque fasciis tribus interruptis flavis. *Degeer Inf.* 6. 101. 3.

*Geoff. Inf.* 2. 513.

*Frisch. Inf.* 4. tab. 13.

*Reaum. Inf.* 4. tab. 31. fig. 9 — 11.

Habitat in Europae aquis stagnantibus.

*Larua* tubo filiformi respiratorio suspendenda.

11. *S. antennis setariis tomentosus*, thorace nigro fasciato, abdomine flauo, segmentorum marginibus lineaque dorsali atris. *Syst. Ent.* 764. 8. \* *floreus.*

*Musca florea*, antennis setariis tomentosa, thorace fasciis duabus inaequalibus nigris, abdomine quatuor luteis subabruptis. *Linn. Syst. Nat.* 2. 984. 29. *Fn. Sv.* 1796.

*Musca antennis setariis tomentosa nigra*, capite thoraceque pilis viridi flavis, abdomine fasciis quatuor luteis, prima maiore interrupta. *Degeer Inf.* 6. 100. 2. tab. 6. fig. 2.

*Conops florens.* *Scop. carn.* 959.

*Alb. Inf.* tab. 17. fig. 1.

Habitat in Europae floribus.

12. *S. antennis setariis tomentosus*, abdomine ferrugineo, dorso nigro, alis crassioribus. *Syst. Ent.* 764. 9. \* *subcoleoptratus.*

*Conops subcoleoptratus* antennis setariis, abdomine subferrugineo, alis praemorsis externe crassioribus.

*Linn. Syst. Nat.* 2. 1006. 13.

Habitat in Angliae nemoribus.

424 .GLASS. VII. ANTLIATA. Syrphus.

Varietatem minorem ex Italia abdomine macula bifidos tantum rufa misit *D. Allioni*.

*nemorum*. 13. S. antennis setariis tomentosus, abdomine atro, cingulis tribus albis, primo segmento lateribus luteo. *Syst. Ent.* 764. 10.

*Musca nemorum*. *Linn. Syst. Nat.* 2. 984. 30. *Fa. Sv.* 1797.

*Musca* thorace cinereo, abdominis basi fascia lutes interrupta, marginibus incisurarum albis. *Geoff. Inf.* 2. 511. 36.

*Musca horticola* antennis plumatis tomentosa, abdomine nigro, maculis binis lateralibus magnis, cingulisque tribus flavo testaceis. *Degeer Inf.* 6. 140. 4. tab. 8. fig. 12.

*Reaum. Inf.* 3. tab. 31. fig. 8.

*Merian. Europ.* tab. 2.

*Alb. Inf.* tab. 3. fig. M. L.

Habitat in Europae floribus frequens.

*hortorum*. 14. S. antennis setariis tomentosus, thorace subsfasciatus, abdomine fasciis tribus albis, pedibus posticis incrassatis ferrugineis. *Syst. Ent.* 764. 11. \*

*Musca surinamensis* antennis setariis tomentosa fusa, scutello rufo, abdomine maculis binis magnis cingulisque tribus flavo testaceis. *Degeer Inf.* 6. 145. 1. tab. 29. fig. i.

Habitat in America meridionali.

*arbus-  
torum*. 15. S. antennis setariis tomentosus, abdomine nigro, segmento primo secundoque latere ferrugineis. *Syst. Ent.* 765. 12.

*Musca arbusorum*. *Linn. Syst. Nat.* 2. 984. 31. *Fa. Sv.* 1798.

Habitat in Europae arbutis.

*prato-  
rum*. 16. S. antennis setariis tomentosus niger, thorace fasciis punctisque duobus, abdomine maculis vtrinque tribus flavis. *Syst. Ent.* 765. 13. \*

Habitat in America. *Dom. v. Rohr.*

7. S. antennis setariis tomentosus, thorace lineis duabus fruteto-  
flavis, abdomine ferrugineo, linea dorsali inter- rum.  
rupta nigra. *Syst. Ent.* 765. 14.\*

Habitat in Angliæ arboretis.

8. S. antennis setariis subtomentosa, thorace aeneo, setaribus flavis, abdomine fulvo, striis tribus atris. ericeto-  
rum.

Habitat in Africa æquinoctiali. *Mus. Dom. Banks.*

Statura M. nemorum. Os flavum. Thorax æneus lateribus scutelloque flavescens. Abdomen fulvum strigis tribus atris. Pedes flavi.

9. S. antennis setariis tomentosus, thorace griseo, abdomine fusco, tibiis posticis compresso gibbis. *Syst.* tenax.  
*Ent.* 765. 15.

*Musca tenax.* Linn. *Syst. Nat.* 2. 984. 32. *Fn. Sv.* 1799.

*Musca fusca*, segmentis abdominalibus tribus margine albidis, primo latere flavo, thorace vix maculato. *Geoff. Inf.* 2. 520. 52.

*Musca porcina* antennis setariis tomentosa fusca, capite linea nigra nitida, thorace griseo, abdomine antice maculis fulvis. *Degeer Inf.* 6. 98. 1.

*Musca apiformis* tota fusca, cauda obtusa ex eruca caudata in latrinis degente orta. *Raj. Inf.* 272.

*Swammerd. Bibl. nat.* tab. 38. fig. 9.

*Reaum. Inf.* 4. tab. 20. fig. 7.

*Blanc. Inf.* tab. 10. fig. F.

*Goed. Inf.* 1. tab. 2.

*Albin, Inf.* tab. 63. fig. e. f. g.

Habitat in fimetis, cloacis, aquis putrescentibus.

*Larva* tenacissima vix prelo destruenda.

20. S. antennis setariis tomentosus, thorace pallido, lineis nigris, abdomine atro, fascia bascos interrupta trilinea-  
tus.  
ferruginea. *Syst. Ent.* 766. 16.\*

Habitat in noua Zelandia. *Mus. Dom. Banks.*

21. S. antennis setariis tomentosus, thorace quinquelinea- quinqueli-  
to, abdomine nigro, segmentis anticis basi albis. neatns.

Habitat in Africa æquinoctiali. *Mus. Dom. Banks.*

Medius. Frons cinerea lineola vtrinque nigra. Thorax lineis quinque nigris albisque alternis. Scutellum



426 CLASS. VIII. ANTLIATA. Syrphus.

lum obscure pallidum. Abdomen nigrum segmento 1. 2. 3. basi albis, 4 immaculato, fuscus basi late album. Pedes quatuor antici albi femoribus nigris, postici nigri tibiis tarsisque basi albis.

- intricarius.* 22. S. antennis setariis tomentosus lutescens, abdomine nigro, apice genubusque albis. *Syst. Ent.* 766. 17.  
*Musca intricaria.* Linn. *Syst. Nat.* 2. 985. 33. *Fn. Sv.* 1800.  
*Musca* antennis plumatis lenticularibus tomentosa nigra, capite thoraceque villis rufis, abdominis apice villosa griseo. *Degeer Inf.* 6. 138. 3. tab. 8. fig. 10.  
 Habitat in Europae borealis floribus.

- oesitraceus.* 23. S. antennis setariis nigra, scutello albido, abdominis apice lutescente. *Syst. Ent.* 766. 18.  
*Musca oesitracea* antennis setariis tomentosa nigra scutello albido, abdominis apice lutescente. Linn. *Syst. Nat.* 2. 985. 34. *Fn. Sv.* 1801.  
*Musca* antennis plumatis tomentosa nigra, scutello albido, capite antice thoraceque postice villis albis, abdominis apice villosa rufo, tarsis flavis. *Degeer Inf.* 6. 139. 4.  
*Schaeff. Icon.* tab. 10. fig. 6.  
 Habitat in Europae borealis floribus.

- afiliformis.* 24. S. antennis setariis, thorace tomentoso flavescente, abdomine atro, segmento primo secundoque albicantibus. *Gen. Inf. Mant.* 306. \*  
 Habitat in Germania. *Dom. de Hattorf.*

- lucorum.* 25. S. antennis setariis tomentosus, thorace griseo, abdomine atro, segmento primo toto albo. *Syst. Ent.* 766. 19.  
*Musca lucorum.* Linn. *Syst. Nat.* 2. 985. 36. *Fn. Sv.* 1803.  
 Habitat in Europae lucis.

- fallax.* 26. S. antennis setariis tomentosus fuscus, fronte flava, abdominis apice tomentoso fulvo, squama halterum lactea.  
*Musca fallax.* Linn. *Syst. Nat.* 2. 985. 36. *Fn. Sv.* 1802.

*Musca*

CLASS. VIII. ANTLIATA. Syrphus: 427

*Musca ruficaudis* antennis setariis tomentosa nigra, fronte flava, abdominis apice villosissimo fulvo. *Degeer Inf.* 6. 127. 7.

Habitat in Europae borealis floribus.

27. S. antennis setariis tomentosus niger, abdominis vlti. *sylluarum*, mo segmento tomentoso flavo. *Syst. Ent.* 767. 20.

*Musca sylvarum*. Linn. *Syst. Nat.* 2. 985. 37. *Fn. Sv.* 1804.

Habitat in Europae borealis nemoribus.

28. S. antennis setariis elongatis nudus ater, alis cyaneis. *nigrita*.

Habitat in Iamaica. *Mus. Dom. Teats*.

Magnus totus ater, immaculatus. Antennae elongatae, setariae. Alae cyanae margine tenuiori albido.

29. S. antennis setariis tomentosus cinereus, thorace fascia nigra, abdominis apice fulvo, femoribus posticis arcuatis. *clausipes*.

Habitat in Italiae floribus. *D. Allioni*.

Corpus magnum, elongatum. Antennae setariae nigrae. Frons pubescens, cinereus. Thorax dense tomentosus cinereus fascia media lata atra. Abdomen cylindricum, tomentosum basi cinereum, apice fulvum. Alae albae immaculatae. Pedes nigri cinereo subtomentosi femoribus posticis valde incrassatis arcuatis apice vnidendatis.

30. S. antennis setariis elongatis niger, thorace lateribus punctis abdomineque cingulis duobus flavis. *bicinctus*. *Syst. Ent.* 767. 21.

*Musca bicincta*. Linn. *Syst. Nat.* 2. 985. 38. *Fn. Sv.* 1805.

*Musca* antennis setariis elongatis tomentosa nigra, abdomine cingulis duobus flavis, alis macula nigricante: *Degeer Inf.* 6. 126. 16. tab. 7. fig. 16.

Habitat in Europae borealis floribus.

31. S. antennis setariis elongatis niger, thorace maculis lateralibus, abdomine cingulis quatuor arcuatis flavis. *arcuatus*. *Syst. Ent.* 767. 22.

*Musca*

428 CLASS. VIII. ANTLIATA. Syrphus.

- Musca arcuata*. Linn. *Syst. Nat.* 2. 985. 39. *Fn. Sv.* 1806.  
*Musca nigra*, flauo variegata, alis fuscis, antennis capite longioribus. *Geoff. Inf.* 2. 506. 28.  
*Musca* antennis setariis elongatis tomentosa nigra pedibus flauis, thorace maculis lateralibus, abdomine cingulis quatuor interruptis arcuatis flauis. *Deger Inf.* 5. 725. 15. tab. 7. fig. 14.  
 Habitat in Europae floribus.
- cingulatus*. 32. S. antennis setariis tomentosus, abdomine atro, segmento primo basi, reliquis punctis albis. *Syst. Ent.* 767. 23.\*  
 Habitat in noua Zelandia. *Muf. Dom. Banks.*
- mutabilis*. 33. S. antennis setariis elongatis tomentosus, thorace immaculato, abdomine nigro griseoque mutabili. *Syst. Ent.* 767. 24.  
*Musca mutabilis*. Linn. *Syst. Nat.* 2. 985. 40. *Fn. Sv.* 1807.  
 Habitat in Europa boreali.
- deuius*. 34. S. antennis setariis elongatis tomentosus, thorace corulefcente, abdomine obscure violaceo. *Syst. Ent.* 767. 25.\*  
*Musca deuia*. Linn. *Syst. Nat.* 2. 985. 41. *Fn. Sv.* 1804.  
 Habitat in Europa, Anglia.
- leucopus*. 35. S. antennis subclauatis nigra, plantis albis.  
*Musca leucopa*. Linn. *Syst. Nat.* 2. 983. 23.  
 Habitat in Americae meridionalis insulis. *Dom. Lund.*  
 Magnus. Caput mihi deest. Corpus nigrum abdominis segmentis margine laterali albicantibus. Primum segmentum subtus rubrum. Pedes apice albi.
- segnis*. 36. S. antennis setariis nudus, thorace aeneo, abdomine ferrugineo, apice nigro. *Syst. Ent.* 772. 47.  
*Musca segnis*. Linn. *Syst. Nat.* 2. 988. 57. *Fn. Sv.* 1823.  
*Musca oblonga*, thorace aeneo, abdomine antennis flauo posterius nigro. *Geoff. Inf.* 2. 526. 62.  
*Musca* antennis setariis tomentosa, corpore elongato depresso, thorace aeneo, abdomine rufo extremitatibus

CLASS. VIII. ANTEIATA. Syrphus. 429

tibus nigro, femoribus posticis clauatis spinosis, *De-  
geer Inf.* 6. 121. 12. tab. 7. fig. 10.

Habitat in Europae nemoribus.

Variat thorace pedibusque nigris et tibiis albo annu-  
latis.

Varietatem ex Italia misit *D. Allioni* abdomine antice  
ferrugineo nigro maculato.

37. *S. antennis setariis tomentosus, thorace subaeneo, fe-  
moribus posticis clauatis. Syst. Ent.* 772. 48. *femora-  
tus.*

*Musca femorata* antennis setariis tomentosa, thorace  
subaeneo, femoribus posticis clauatis dentatis. *Linn.*

*Syst. Nat.* 2. 988. 58. *Fn. Sv.* 1824.

Habitat in Europa.

Abdomen mox nigrum, mox pube ferruginea.

38. *S. antennis setariis nudus niger, thorace lincis latera- conopseus.  
libus, abdomine fasciis tribus flavis. Syst. Ent.* 768. 26.

*Musca conopsoidea* antennis clauatis mucronatis basi  
coadunatis, abdomine cingulis tribus flavis. *Linn.*

*Syst. Nat.* 2. 982. 21. *Fn. Sv.* 1790.

*Reaum. Inf.* 4. tab. 33. fig. 12. 13.

Habitat in Europae nemoribus.

39. *S. antennis setariis elongatis cylindricis niger, fronte ichneumonea  
thoracis postico pedibusque ferrugineis. Syst. Ent.* 768. 27. *ichneumonea.*

*Musca ichneumonea* antennis setariis elongatis cylin-  
dricis, thorace pedibusque longiusculis ferrugineis,  
abdomine cinereo. *Linn. Syst. Nat.* 2. 986. 42. *Fn.  
Sv.* 1809.

Habitat in Europa.

40. *S. antennis setariis elongatis niger, pedibus rufis. sphegus.  
Syst. Ent.* 768. 28. \*

Habitat in Anglia. *Mus. Dom. Banks.*

41. *S. antennis setariis nudus ater, alis nigris, margine cylindri-  
tenuiori albo maculato. cus.*

Habitat in Americae meridionalis insulis. *Dom. Vahl.*

Caput orbiculatum oculis magnis connatis nigris, fron-  
te flava. Thorax brevis, ater, immaculatus. Ab-  
domen

430 CLASS. VIII. ANTLIATA. Syrphus

domen elongatum, cylindricum, atrum. Alae nigrae margine tenuiori maculis magnis albidis.

- elongatus.* 42. S. antennis setariis nudus, thorace aeneo, abdomine clauato nigro, fasciis duabus flauis. *Syst. Ent.* 768. 29. \*

Habitat in Daniae floribus.

- podagricus.* 43. S. antennis setariis nudus, thorace aeneo, abdomine bifasciato, femoribusque posticis clauatis. *Syst. Ent.* 768. 30. \*

Habitat in Daniae floribus.

- crabroniformis.* 44. S. antennis setariis tomentosus, thorace variegato, abdomine flauo, segmento primo secundoque apice fuscis. *Syst. Ent.* 768. 31. \*

Habitat in Madera Insula. *Mus. Dom. Banks.*

- diopthalmus.* 45. S. antennis setariis nudus, thorace maculato, abdomine cingulis sex flauis, femoribus posticis dentatis. *Syst. Ent.* 769. 32. \*

*Musca diopthalma.* Linn. *Syst. Nat.* 2. 986. 43. *Fn. Sv.* 1810.

Habitat in Europae floribus.

- vespiformis.* 46. S. antennis setariis nudus, thorace maculato, abdomine cingulis sex flauis. *Syst. Ent.* 769. 33. \*

*Musca vespiformis.* Linn. *Syst. Nat.* 2. 986. 44. *Fn. Sv.* 1811.

*Musca* antennis setariis tomentosa nigra, pedibus antennisque flauis, alis fuluo variegatis, abdomine cingulis luteis alternatim latis. *Degeer Inf.* 6. 122. 13. tab. 7. fig. 13.

Habitat in Europae floribus.

- festuus.* 47. S. antennis setariis nudus, thorace lineis lateralibus, abdomine cingulis quatuor flauis interruptis. *Syst. Ent.* 769. 34.

*Musca festiua.* Linn. *Syst. Nat.* 2. 986. 45. *Fn. Sv.* 1811.

*Musca* nigra flauo variegata, alis fuscis, antennis capite breuioribus. *Geoff. Inf.* 2. 505. 27.

*Musca citro fasciata* antennis setariis tomentosa nigra, thorace lineis binis lateralibus scutelloque luteis, abdomine

**CLASS. VIII. ANTLIATA. Syrphus. 431**

domine cingulis quatuor luteis interruptis. *Deger*  
*Inf.* 6. 118. 9.

*Conops festiva. Scop. carn.* 964.

Habitat in Europae floribus frequens.

48. S. antennis setariis tomentosus, thorace lineato, abdomine aeneo, alis maculatis. *Syst. Ent.* 769. 35. \* **ruficornis.**  
 Habitat in Daniae floribus.

49. S. antennis setariis tomentosus obscure aeneus, antennis tibiisque flavescentibus. **flavicornis.**  
 Habitat in Germania. *Dom. de Hattorf.*  
 Affinis S. ruficorni. Alae immaculatae.

50. S. antennis setariis niger immaculatus, alis albis macula fusca. *Syst. Ent.* 770. 36. \* **lugubris.**  
 Habitat in Daniae floribus.

51. S. antennis setariis tomentosus niger, abdomine atro sepulcrali. **sepulcralis.**  
*Syst. Ent.* 772. 49.  
*Musca sepulcralis. Linn. Syst. Nat.* 2. 991. 73. *Fn. Sv.* 1835.  
 Habitat in Sueciae floribus.  
 Oculi maculati.

52. S. antennis setariis aeneus nitens, antennis nigris. **metallina.**  
*Gen. Inf. Mant.* 307. \*  
 Habitat in Germaniae floribus.

53. S. antennis setariis subtomentosus niger, abdomine glaucius. basi pellucido, cingulisque duobus albidis interruptis. **glaucius.**  
*Syst. Ent.* 770. 37. \*  
*Musca glaucia. Linn. Syst. Nat.* 2. 986. 47. *Fn. Sv.* 1813.  
*Conops glaucius. Scop. carn.* 966.  
 Habitat in Europae floribus.

54. S. antennis setariis subtomentosus ater, abdominis primo segmento maculis duabus pellucidis. **noctiluca.**  
*Syst. Ent.* 770. 38.  
*Musca noctiluca. Linn. Syst. Nat.* 2. 986. 48. *Fn. Sv.* 1814.  
 Habitat in Europae floribus.

432 CLASS. VIII. ANTLIATA. Syrphus.

*gibbosus*. 55. S. antennis setariis subtomentosus fuscus, abdomine subglobofo atro, cingulis quatuor albis, squamis halterum buccatis. *Syst. Ent.* 770. 39.

*Musca gibbosa*. Linn. *Syst. Nat.* 2. 987. 49. Fn. Sv. 1815.

Habitat in Europae floribus.

*globosus*. 56. S. antennis setariis subtomentosus, thorace antice ferrugineo, abdomine subglobofo pallido, apice nigro. *Syst. Ent.* 770. 40.\*

Habitat in Angliae floribus.

*Ribesii*. 57. S. antennis setariis nudiusculus, thorace immaculato, abdomine cingulis quatuor flavis, primo interrupto. *Syst. Ent.* 770. 41.

*Musca Ribesii*. Linn. *Syst. Nat.* 2. 987. 50. Fn. Sv. 1816.

*Musca* abdomine ovato nigro, lunularum pari cingulisque tribus flauescentibus. *Geoff. Inf.* 2. 511. 45.

*Musca* antennis setariis tomentosa, thorace aeneo pilis flauescentibus, abdomine nigro, fasciis quatuor transversis flavis, prima interrupta. *Degeer Inf.* 6. 103. 4. tab. 6. fig. 8.

*Goed. Inf.* 1. tab. 41.

Habitat inter Aphides Ribis.

*Pyrastris*. 58. S. antennis setariis niger nudiusculus, thorace immaculato, abdomine bis tribus lunulis albis recurvatis. *Syst. Ent.* 771. 42.

*Musca Pyrastris*. Linn. *Syst. Nat.* 2. 987. 51. Fn. Sv. 1817.

*Musca* thorace nigro viridi, abdomine atro ovato, tribus paribus lunularum albicantium. *Geoff. Inf.* 2. 516. 46.

*Musca Rosae* antennis setariis tomentosa, thorace aeneo, abdomine nigro, tribus paribus lunularum albicantium. *Degeer Inf.* 6. 108. 5. tab. 6. fig. 18.

*Scop. carn.* 931.

*Frisch. Inf.* 11. tab. 22. fig. 1.

*Reaum. Inf.* 3. tab. 31. fig. 9.

*Albin. Inf.* tab. 66.

*Goed. Inf.* 1. tab. 47.

**CLASS VIII. ANTLIATA. Syrphus 439**

*Blank. Inf. tab. 10. fig. P.*

*Sulz. Inf. tab. 20. fig. 132.*

*Merian. Europ. tab. 89.*

Habitat inter Aphides Pyri.

59. *S. antennalis* setariis nudus niger, thorace maculato, abdomine cingulis quatuor scutelloque flavis. *Syst. Ent. 771. 43.* *Mentha-firi.*

*Musca Mentha-firi. Linn. Syst. Nat. 2. 987. 53. Fn. Sv. 1819.*

*Musca* thorace nigro viridi, apice flauo, abdomine nigro, fasciis luteis transuersis aequalibus. *Geoff. Inf. 2. 514. 41.*

*Musca* antennis setariis tomentosa nigra, corpore elongato depresso, thorace luteo maculato, abdomine cingulis quatuor, scutello pedibusque flavis. *Degeer Inf. 6. 119. 10.*

Habitat inter Aphides larua, at Syrphus in floribus melilugus.

60. *S. antennalis* setariis tomentosus, thorace subaeneo, abdomine atro, fasciis tribus albidis, secunda interrupta. *Thyma-firi.*

Habitat in Germaniae floribus. *Dom. de Hattorf.*

Statura et summa affinitas *S. Mentha-firi*. Frons flaua antennis nigris. Thorax tomentosus, subaeneus, scutello flauo. — Abdomen atrum fasciis tribus glaucis, secunda late interrupta. Alae immaculatae. Pedes flauo femoribus basi nigris.

61. *S. antennalis* setariis nudus, thorace subaeneo immaculato, abdomine maculis octo lutescentibus. *Syst. Ent. 771. 44.* *mellinus.*

*Musca mellina. Linn. Syst. Nat. 2. 988. 55. Fn. Sv. 1821.*

*Musca* thorace nigro viridi, apice flauo, abdomine oblongo, paribus tribus trigonorum lutescentium. *Geoff. Inf. 2. 515. 42.*

*Musca* antennis setariis tomentosa, thorace aeneo, abdomine nigro, maculis octo tetragonis flauo testaceis. *Degeer Inf. 6. 117. 8.*

*Fabr. Sp. Inf. T. II.*

E e

*Musca*



434 CLASS. VIII. ANTLIATA. Syrphus.

*Musca mellina.* Scop. carn. 934.  
Habitat inter Europae Aphides.

*scriptus.* 62. S. antennis fetariis nudus, thorace lineis abdomineque lineari fasciis flavis. *Syst. Ent.* 772. 45.

*Musca scripta.* Linn. *Syst. Nat.* 2. 987. 54. *Fa. Sv.* 1820.

*Reaum. Inf.* 4. tab. 10. fig. 2. 3.

*Roef. Inf.* 2. *Musc.* tab. 6.

Habitat inter Aphides Larua, in floribus Syrphus.

*albimanus.* 63. S. antennis fetariis tomentosus, thorace aeneo, abdomine elongato atro, punctis quatuor albis.

Habitat in Angliae hortis. *Mus. Dom. Banks.*

Statura et magnitudo omnino S. scripti. Os aeneum linea media eleuata. Thorax cum scutello tomentosus, aeneus. Abdomen elongatum, holosericeum, atrum punctis quatuor albis. Pedes fusci tibiis anticis albis.

*dimidiatus.* 64. S. antennis fetariis nudus ater, alis basi nigris.

Habitat in Americae meridionalis insulis. *Mus. Dom. Teats.*

Paruus. Thorax fuscus vix pubescens. Abdomen glabrum, atrum, nitidum. Alae albae a basi ad medium nigrae. Pedes nigri.

*pipiens.* 65. S. antennis fetariis nudus, abdomine vtrinque albo maculato, femoribus posticis clauatis dentatis. *Syst. Ent.* 772. 46.

*Musca pipiens.* Linn. *Syst. Nat.* 2. 988. 56. *Fa. Sv.* 1822.

*Musca oblonga,* femoribus posticis maioribus. *Gross. Inf.* 2. 519. 49.

*Musca* antennis fetariis tomentosa nigra, corpore elongato cylindrico, abdomine maculis sex albidis, femoribus posticis clauatis, dentatis. *Degeer Inf.* 4. 120. 11. tab. 7. fig. 8.

*Conops pipiens.* Scop. carn. 969.

Habitat in Europae floribus.

176. M V S C A.

\* Antennis seta plumata.

1. *M. antennis plumatis fusca, abdomine pellucido, cin- inanis.*  
gulis tribus nigris. *Syst. Ent.* 773. 1.  
*Musca inanis* antennis plumatis pilosa flavescent, abdo-  
mine pellucido, cingulis duobus nigris. *Linn. Syst.*  
*Nat.* 2. 989. 61. *Fn. Sv.* 1825.  
*Musca apiuora* antennis plumatis pilosa, capite flavo,  
thorace nigro, abdomine fasciis flavis nigrisque, alis  
flavescentibus fusco maculatis. *Degeer Inf.* 6. 56. 3.  
tab. 3. fig. 4.  
*Nemotelus niger*, abdomine niueo, fasciis duabus ni-  
gris. *Geoff. Inf.* 2. 543. 1. tab. 18. fig. 4.  
*Harris Inf. angl.* tab. 10. fig. 2.  
*Reaum. Inf.* 4. tab. 33. fig. 15.  
*Schaeff. Icon.* tab. 36. fig. 7. 8.  
*Habitat in Europae nemoribus.*
2. *M. antennis plumatis nigra, abdominis segmento primo pellucens.*  
pellucido, albo. *Syst. Ent.* 773. 2. *Linn. Syst. Nat.*  
2. 989. 62. *Fn. Sv.* 1826.  
*Musca* antennis plumatis pilosa nigra, fronte flava, ab-  
dominis segmento secundo albo pellucido, alis ma-  
cula venisque fuscis. *Degeer Inf.* 6. 53. 1. tab. 3.  
fig. 1.  
*Voluccella* abdomine antice albo, postice nigro, alis al-  
bis, nigra macula. *Geoff. Inf.* 2. 540. 1. tab. 18. fig. 3.  
*Harris Inf. angl.* tab. 10. fig. 4.  
*Raj. Inf.* 271. 1.  
*Sulz. Inf.* tab. 20. fig. 133.  
*Habitat in Europae umbrosis.*
3. *M. antennis plumatis pilosa nigra, fronte aurea, alis meridiana.*  
basi ferrugineis. *Syst. Ent.* 773. 3. *Linn. Syst. Nat.* na.  
2. 989. 63. *Fn. Sv.* 1827.  
*Musca* antennis plumatis pilosa nigra, fronte flava, alis  
basi flavo testaceis. *Degeer Inf.* 6. 55. 2.  
*Musca meridiana.* *Scop. carn.* 1827.  
*Reaum. Inf.* 4. tab. 26. fig. 10.

*Harris Inf. angl. tab. 9. fig. 9.*

*Schaeff. Icon. tab. 108. fig. 7.*

*Musca nigra*, alis basi ferrugineis. *Geoff. Inf. 2. 495. 4.*

Habitat in Europae nemoribus frequens.

*carnaria. 4. M. antennis plumatis pilosa nigra, thorace lineis pallidioribus, abdomine nitido tessellato. Syst. Ent. 773.*

*4. Linn. Syst. Nat. 2. 990. 68. Fa. Sv. 1832.*

*Musca nigra*, abdomine nitido tessellato, thorace lineolis pallidioribus longitudinalibus, ano fulvo. *Geoff. Inf. 2. 527. 65.*

*Musca vivipara maior* antennis plumatis pilosa cinerea, oculis rubris, thorace lineis tribus nigris, abdomine maculis nigris cinereis tessellato, maior. *Degur*

*Inf. 6. 63. 8. tab. 3. fig. 5.*

*Musca carnaria vulgaris. Raj. Inf. 270. 1.*

*Musca subvariegata. Scop. carn. 869.*

*Aldr. Inf. 348. tab. 2. fig. 16.*

*Frisch. Inf. 7. tab. 14.*

*Reaum. Inf. 4. tab. 29. fig. 4 — 6.*

*Roef. Inf. 2. Musc. tab. 9. fig. 10.*

Habitat in Europae, Americae cadaueribus.

*lardaria. 5. M. antennis plumatis nigra, thorace abdomineque tessellatis.*

Habitat in Germania. *Dom. de Hattorf.*

Minor *M. carnariae*, cui simillima. Thorax et abdomen concolora obscure tessellata. Abdomen magis globosum, obtusum. Pedes nigri.

*utilis. 6. M. antennis plumatis, thorace lineato, abdomine tessellato, ano rufo.*

Habitat in Americae meridionalis insulis. *Dom. Vahl.*

Maior *M. domesticae*. Os albidum antennis nigris. Thorax pilosus, cinereus, nigro lineatus. Abdomen conicum, pilosum, tessellatum, ano siue ultimo segmento rufo.

*domestica. 7. M. antennis plumatis, thorace lineato, abdomine tessellato subtus basi pallido. Syst. Ent. 774. 5.*

*Musca domestica* antennis plumatis pilosa nigra, thorace lineis quinque obsoletis, abdomine nitidulo tessellato,

fellato, oculis fuscis. *Linn. Syst. Nat.* 2. 990. 69. *Fn. Sv.* 1833.

*Musca nigra*, abdomine nitidulo, tessellato, thorace lineolis pallidioribus longitudinalibus, ano concolore. *Geoff. Inf.* 2. 528. 66.

*Musca domestica maior* antennis plumatis pilosa, thorace cinereo nigro lineis obscuris, abdomine nigro fusco, maculis nigris, subtus pallide flavo fusco, alarum basi fulva. *Degeer Inf.* 6. 72. 10. tab. 4. fig. 5. 6.

*Musca bipennis maior*, thorace glabro striato, abdomine supino nigro, toto maculis albis punctato. *Raj. Inf.* 270. 2.

*Scop. carn.* 872.

*Aldr. Inf.* tab. 2. fig. 23.

*Toblot. micr.* 1. pp. tab. 5.

Habitat Larva in stercore equino, Musca in domibus.

8. *M. antennis plumatis atra*, thoracis lateribus canis nigro punctatis. *Syst. Ent.* 774. 6. \* *americana.*  
Habitat in America meridionali.

9. *M. antennis plumatis pilosa*, thorace cinereo aeneo, abdomine subaureo, ano retuso. *Syst. Ent.* 775. 7. \* *retusa.*  
Habitat in noua Hollandia. *Mus. Dom. Banks.*

10. *M. antennis plumatis viridi nitens*, pedibus nigris. *Caesar.*  
*Syst. Ent.* 775. 8. *Linn. Syst. Nat.* 2. 989. 64. *Fn. Sv.* 1828.

*Musca* thorace abdomineque viridi nitente, pedibus nigris. *Geoff. Inf.* 2. 522. 53.

*Musca bipennis*, thorace et abdomine superne coeruleo colore pulcro nitente, capite nigro. *Raj. Inf.* 272. 1.

*Degeer Inf.* 6. 61. 6.

*Scop. carn.* 871.

*Mouff. Inf.* tab. 49.

*Merian. Europ.* 1. tab. 49.

*Reaum. Inf.* 4. tab. 8. fig. 1.

— — — tab. 19. fig. 8.

438 CLASS. VIII. ANTLIATA. Musca.

*β. Musca caesarina. Scop. carn. 873.*

Habitat in Europae cadaueribus.

*β. Triplo minor, et simillima, vix distinguenda.*

*cornicina. 11. M. antennis plumatis, thorace cupreo nitido, abdomine viridi aeneo, pedibus nigris.*

Habitat in Italia. *D. Allioni.*

Vix satis distincta a *M. Caesare*. Labium lateribus argento nitet. Thorax cupreus, immaculatus.

*cadaveri. 12. M. antennis plumatis nitens, thorace coeruleo, abdomine viridi. Syst. Ent. 775. 9. Linn. Syst. Nat. 2. 989. 65. Fn. Sv. 1829.*

*Musca* thorace coeruleo nitente, abdomine viridi nitente. *Geoff. Inf. 2. 524. 57.*

*Musca viridi coerulea* antennis plumatis pilosa, thorace viridi coeruleo nitente, abdomine coeruleo violaceo, pedibus nigris. *Degeer Inf. 6. 62. 7.*

*Goed. Inf. 1. tab. 54.*

Habitat in Europae cadaueribus.

*putrida. 13. M. antennis plumatis viridi nitens, abdomine globoso cyaneo. Syst. Ent. 775. 10.\**

Habitat in America. *Mus. Tottianum.*

*fygia. 14. M. antennis plumatis pilosa, thorace coerulescente, abdomine obscure aeneo, pedibus ferrugineis.*

Habitat in Insula Terra noua Americae. *Mus. Dom. Banks.*

Paullo minor *M. mortuorum*. Caput fuscum ore subferrugineo. Thorax pilosus, coerulescens, immaculatus. Abdomen subrotundum, obscure aeneum. Pedes ferruginei plantis nigris.

*mortuorum. 15. M. antennis plumatis, thorace nigro, abdomine viridi aeneo, pedibus nigris. Syst. Ent. 775. 11. Linn.*

*Syst. Nat. 2. 989. 66. Fn. Sv. 1830.*

*Musca carnaria coerulea* antennis plumatis pilosa, thorace nigro, abdomine coeruleo nitente fasciis nigris. *Degeer Inf. 6. 57. 4.*

Habitat in Europae cadaueribus.

Varietatem duplo minorem ex Italia misit *Allioni*.

16. *M. antennis plumatis*, thorace aeneo albo punctato, *leonina*.  
abdomine coeruleo, ano bipunctato. *Syst. Ent.* 776.

12. \*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

17. *M. antennis plumatis*, thorace nigro, abdomine coeruleo nitente, fronte fulua. *Syst. Ent.* 776. 13. *Linn. Sys. Nat.* 2. 989. 67. *Fn. Sv.* 1831.

*Musca* thorace nigro, abdomine coerulescente. *Geoff.*

*Inf.* 2. 524. 59.

*Musca carnaria*. *Scop. carn.* 868.

*Musca chrysocephala* antennis plumatis pilosa, capite aureo flauo, thorace nigro violaceo, abdomine viridi coeruleo nitente. *Degeer Inf.* 6. 69. 5.

*Goed. Inf.* 1. tab. 53.

*Raj. Inf.* 271.

*Merian. Eurap.* tab. 49.

*Reaum. Inf.* 4. tab. 24. fig. 13 — 15.

Habitat in Europae cadaueribus.

18. *M. antennis plumatis*, thorace abdomineque cyaneis, *cyaneis*.  
pedibus nigris.

Habitat ad Cap. bon. sp.

Magnitudo et statura praecedentis. Corpus cyaneum squama halterum lactea. Pedes nigri femoribus anticis cyaneis.

19. *M. antennis plumatis*, thorace cupreo nigro lineato, *macellaria*.  
abdomine aeneo, pedibus nigris. *Syst. Ent.* 775. 14. \*

Habitat in America meridionali.

20. *M. antennis plumatis*, thorace lineato, abdomine sub-ferrugineo nigro maculato. *Syst. Ent.* 776. 15. \*

*Musca maculata* antennis plumatis pilosa nigra, thorace nigro lineato, abdomine atro maculato, ano bipunctato. *Linn. Sys. Nat.* 2. 990. 70.

*Musca* antennis plumatis pilosa nigro cinerea maculis nigris, thorace lineis quatuor nigris. *Degeer Inf.* 6. 84. 13. tab. 3. fig. 22.

Habitat in Europa boreali.

*cornuta*. 21. *M. antennis plumatis nigra, abdomine testaceo, lineis dorsali nigra.*

*Musca autumnalis antennis plumatis pilosa nigra, abdomine flavo testaceo, fascia longitudinali dorsali nigra. Degeer Inf. 6. 83. 12.*

Habitat in Germania. *Dom. de Hattorf.* In Italia. *D. Allioni.*

Magnitudo *M. vulpinae*. Caput nigrum orbita oculorum antice argentea, postice testacea. Thorax pilosus, niger, vix lineatus. Abdomen testaceum albo vario situ nitidum linea dorsali nigra, quae basi apiceque extenditur, subtus testaceum ano nigro. Pedes nigri.

*testacea*. 22. *M. antennis plumatis cinerea, scutello abdominae pedibusque testaceis.*

Habitat in Germania. *Dom. de Hattorf.*

Statura praecedentis. Alae albae immaculatae. Antennae nigrae brevissime plumatae.

*augur*. 23. *M. antennis plumatis cinerea, abdomine coerulescente, lateribus testaceo diaphanis. Syst. Ent. 777. 16.\**

Habitat in nova Hollandia. *Mus. Dom. Banks.*

*camina*. 24. *M. antennis plumatis setaria cinerea, thorace punctis, abdomine strigis nigris.*

Habitat in Anglia. *Mus. Dom. Banks.*

Media. Thorax valde pilosus, cinereus punctis quatuor distinctis dorsalibus nigris. Abdomen pilosum, obconicum, cinereum segmentorum marginibus nigris. Pedes testacei.

*plebeja*. 25. *M. antennis plumatis nigra, pedibus rufis.*

Habitat in Germania, Kiliae.

Magnitudo *M. rusticae*. Caput nigrum, certo situ griseo nitidum. Thorax pilosus, niger, lineola media basios cinerea. Scutellum testaceum. Abdomen pilosum, atrum. Pedes rufi plantis nigris.

*rustica*. 26. *M. antennis plumatis grisea, femoribus rufis. Syst. Ent. 777. 17.\**

Habitat Hafniae.

**CLASS. VIII. ANTLIATA. Musca. 443**

17. *M. antennis plumatis setaria pallide testacea, alis punctis duobus costaque fuscis.* *elata.*

Habitat Kiliae.

Affinis *M. cucullariae*. Os vesiculosum, album. Antennae rufae seta plumata nigra. Thorax pilosus pallide testaceus siue griseus. Abdomen et pedes dilutiora, testacea. Alae albae punctis duobus posteriore maiore transverso, fuscis. Costa alarum nigra.

**\*\* Antennis feta nuda.**

18. *M. antennis setariis nigra, abdominis lateribus testaceo diaphanis.* *Syst. Ent. 777. 18. Linn. Syst. Nat. 2. 961. 74. Fn. Sv. 1836.* *fera.*

*Musca* nigra, abdomine hemisphaerico luteo, fascia longitudinali nigra. *Geoff. Inf. 2. 509. 33.*

*Musca rotundata.* *Harris Inf. angl. tab. 9. fig. 2.*

Habitat in Europa frequens.

19. *M. antennis setariis pilosa, thorace fusco, abdomine atro, lateribus basi testaceis.* *lurida.*

Habitat in Germania. *Dom. de Hattorf.*

Differt manifeste a praecedente. Antennae ferrugineae claua nigra. Os albidum. Thorax pilosus, fuscus scutello flavescente. Abdomen atrum basi lateribus testaceo diaphanis, quae strigis duabus cinguntur.

29. *M. antennis setariis rufa, abdomine fasciis duabus aureis.* *Syst. Ent. 777. 19.\** *bifasciata.*

Habitat in America meridionali. *Dom. v. Rohr.*

30. *M. antennis setariis nigra, alis basi ferrugineis.* *Syst. Ent. 777. 20. Linn. Syst. Nat. 2. 991. 75. Fn. Sv. 1837.* *grossa.*

*Musca* atra basi alarum ferruginea. *Geoff. Inf. 2. 495. 24.*

*Musca* antennis setariis pilosa nigra, capite flavo, oculis fuscis, alis basi ferrugineis. *Degeer Inf. 6. 21. 1. tab. 1. fig. 1.*

*Harris Inf. angl. tab. 9. fig. 1.*

*Schaeff. Icon. tab. 108. fig. 6.*



442 CLASS. VIII. ANTLIATA. Musca.

- Reaum. Inf.* 4. tab. 26. fig. 10.  
Habitat in Europae stercore bouino.
- hystrix.* 31. *M. antennis setariis pilosissima atra, ore albicante. Syst. Ent.* 777. 21.\*  
*Musca pilosa. Drury Inf.* 1. tab. 45. fig. 7.  
Habitat in America meridionali.
- tremula.* 32. *M. antennis setariis pilosa atra nitens, alarum basi, squama halteribusque ferrugineis. Linn. Syst. Nat.* 2. 991. 77.  
*Scop. carn.* 885.  
Habitat in Europa boreali.  
Minor *M. fera* magisque oblonga abdomine nitido.
- rotundata.* 33. *M. antennis setariis, thorace sublineato, abdomine subrotundo ferrugineo, linea longitudinali punctorum nigrorum. Syst. Ent.* 778. 22.  
*Musca rotundata* antennis setariis pilosa, thorace sublineato, abdomine subrotundo flavo, maculis longitudinalibus fuscis confertis. *Linn. Syst. Nat.* 2. 992. 76. *Fn. Sv.* 1838.  
*Musca nigra*, abdomine hemisphaerico rufo punctorum nigrorum ordine longitudinali. *Geoff. Inf.* 2. 509. 32.  
*Musca* antennis setariis subglabra, capite magno albedo, thorace nigro lateribus albo, abdomine subrotundo rufo, maculis quatuor nigris. *Degeer Inf.* 6. 29. 8. tab. 1. fig. 11.  
*Scop. carn.* 883.  
*Harris Inf. angl.* tab. 9. fig. 5. 6.  
Habitat in Europa frequens.
- laruarum.* 34. *M. antennis setariis nigricans, scutelli apice subtestaceo, abdomine tessellato. Syst. Ent.* 778. 23. *Linn. Syst. Nat.* 2. 992. 78. *Fn. Sv.* 1839.  
*Musca maior larvarum* antennis setariis pilosa nigra, thorace lineis longitudinalibus atris, abdomine maculis cinereis maior. *Degeer Inf.* 6. 24. 3. tab. 1. fig. 7.  
*Scop. carn.* 888.  
*Ann. mir. Inf.* tab. 6.  
*Degeer Inf.* 1. tab. 11. fig. 23.

Habitat

CLASS. VIII. ANTLIATA. *Musca*, 443

Habitat in Lepidopterorum Larvis, in radicibus Brassicae oleraceae, unde radix strumosa et capita laxa.

Conf. *Musca minor larvarum*. *Degeer Inf.* 6. 25. 4.  
Vix distincta videtur.

35. *M. antennis setariis nigra*, abdomine cinereo nigro *radicum*. fasciato. *Syst. Ent.* 878. 24.

*Musca radicum* antennis setariis pilosa, abdomine cinereo, linea dorsali cingulisque quatuor nigris. *Linn. Syst. Nat.* 2. 992. 79. *Fn. Sv.* 1839.

Habitat in Raphani radicibus.

36. *M. antennis setariis nigra*, abdomine cylindrico, *brassica-* segmento secundo tertioque rufis. *Syst. Ent.* 877. 25. *ria*.

*Musca cylindrica* antennis setariis pilosa cinereo nigra, abdomine cylindrico elongato medio rufo. *Degeer Inf.* 6. 30. 9. tab. 1. fig. 12.

Habitat in Brassicae radicibus.

37. *M. antennis setariis nigra*, abdominis lateribus basi *lateralis*. sanguineis. *Syst. Ent.* 778. 26. \*

*Musca rufa maculata* antennis setariis pilosa nigra, abdominis lateribus macula magna ferruginea, alis fuscis postice hyalinis. *Degeer Inf.* 6. 28. 7. tab. 1. fig. 9.

Habitat in Germania, Kiliac.

38. *M. antennis setariis nigricans*, abdominis segmento *canicula-* secundo tertioque latere diaphanis. *Syst. Ent.* 779. *ris*. 27. *Linn. Syst. Nat.* 2. 992. 80. *Fn. Sv.* 1841.

Habitat in Europa, sub canicula vesperi et meridie sub arborum coma choreas agens.

39. *M. antennis setariis nigra*, abdomine supra glabro sub- *coemete-* violaceo. *Syst. Ent.* 779. 28. *Linn. Syst. Nat.* 2. *riorum*. 992. 82. *Fn. Sv.* 1842.

Habitat in Europa.

40. *M. antennis setariis cinerea*, thorace maculis quinque *pluvialis*. nigris, abdomine maculis obsoletis. *Syst. Ent.* 779. 29. *Linn. Syst. Nat.* 2. 992. 83. *Fn. Sv.* 1844.

*Musca*

444 CLASS. VIII. ANTLIATA. Musca.

*Musca cinerea* thorace maculis quinque nigris, abdomine maculis tridentatis. *Gross. Inf.* 2. 529.

*Musca antennis setariis pilosa cinerea*, thorace maculis rotundatis, abdomineque maculis nouem nigris trigonis. *Degeer Inf.* 6. 27. 6.

*Scop. carn.* 891.

Habitat in Europa ante pluuiam chœreas agens.

*subul-* 41. *M. antennis setariis tota atra, alis hyalinis, femoribus*  
*tans.* posticis saltatoriis. *Syst. Ent.* 779. 30. *Linna. Syst.*  
*Nat.* 2. 993. 84.

Habitat in Europa boreali, saliendo volitans.

*tigrina.* 42. *M. antennis setariis pilosa cinerea, femoribus apice*  
tibiisque ferrugineis. *Syst. Ent.* 779. 31. \*  
Habitat in Anglia victitans rapina congenerum.

*chorea.* 43. *M. antennis setariis atra, alis totis albis.*

Habitat in Germania. *Dom. de Hattorf.*

Affinis sequenti. Corpus totum atrum alis solis omnino albis.

*roralis.* 44. *M. antennis setariis aterrima, alis apice subalbicanibus.* *Syst. Ent.* 779. 32. \*

*Musca roralis* antennis setariis pilosa, aterrima. *Linna.*  
*Syst. Nat.* 2. 993. 85. *Fn. Sv.* 1246.

*Musca roralis.* *Scop. carn.* 890.

Habitat in Europa.

*fenestra-* 45. *M. antennis setariis glabra atra, abdomine supra ru-*  
*lis.* gofo, strigis albis, alis fuscis. *Syst. Ent.* 780. 33.

*Musca fenestralis* antennis filatis glabra atra, abdomine  
supra rugoso, strigis albis, pedibus ferrugineis. *Linna.*  
*Syst. Nat.* 2. 981. 14. *Fn. Sv.* 1245.

*Nemotelus fenestralis* niger antennis cylindricis muticis,  
abdomine oblongo, fasciis quatuor albis, alis nigro fuscis. *Degeer Inf.* 6. 189. 11.

Habitat in Europae fenestris.

Parua. Pedes nigri postici elongati. Strigae abdominis albae certo abdominis situ disparent.

*medita-* 46. *M. antennis setariis pilosa obscura, abdomine cin-*  
*bunda.* reo, punctis quatuor fuscis.

Habi-

Habitat in Italia. *D. Allioni. Hafniae. Dom. Lund.*  
Media. Os albicat. Thorax pilosus, obscurus, im-  
maculatus. Abdomen ovatum, cinereum punctis  
quatuor dorsalibus fuscis. Pedes nigri tibiis rufe-  
scentibus.

47. *M. antennis setariis cinerea, alarum costa pilis brevif- serratorum*  
simis serrata, abdomine pedibusque ferrugineis. *Syst.*  
*Ent. 780. 34. Linn. Syst. Nat. 2. 993. 86. Fn. Sv.*  
*1847.*

*Musca latrinarum* antennis setariis pilosa cinerea, ab-  
domine elongato conico rufo, pedibus griseo flavis,  
alis abdomine duplo longioribus. *Degeer Inf. 6. 31.*  
*10. tab. 1. fig. 15.*

Habitat in Europae latrinis.

48. *M. antennis setariis nigra, abdomine pallidiore, ocu- cellaris.*  
lis ferrugineis. *Syst. Ent. 780. 35. Linn. Syst. Nat.*  
*2. 992. 87. Fn. Sv. 1848.*

*Musca ferrugineo fusca subpilosa, oculis ferrugineis,*  
alis trineruiis. *Geoff. Inf. 2. 537. 85.*

*Musca oenopota. Scop. carn. 905.*

*Raj. Inf. 261.*

*Reaum. Inf. 5. tab. 8. fig. 7. 11. 12.*

Habitat in Europae cellis vinariis.

49. *M. antennis setariis atra, abdomine cinerascete, alis meteori-*  
basi subflavis. *Syst. Ent. 780. 36. Linn. Syst. Nat. ca.*  
*2. 993. 88. Fn. Sv. 1849.*

*Musca vaccarum* antennis plumatis pilosa, thorace ni-  
gro, abdomine griseo hirsutissimo, alis basi fuscis,

*Degeer Inf. 6. 85. 14. tab. 5. fig. 1.?*

Habitat in Europa, ingruente pluvia choreas agens.

50. *M. antennis setariis nigra, halteribus plantis posticis Frit.*  
abdomineque virescenti pallidis. *Syst. Ent. 780. 37.*  
*Linn. Syst. Nat. 2. 994. 90. Fn. Sv. 1851.*

Habitat in Hordei glumis, grana destruens.

51. *M. antennis setariis atra, alarum costa nigra, alis fer- putris,*  
rugineis. *Syst. Ent. 780. 38. Linn. Syst. Nat. 2. 993.*  
*89. Fn. Sv. 1850.*

*Scop.*

*Scop. carn.* 904.

*Goed. Inf.* 1. tab. 73.

*Merian. Europ.* tab. 43. fig. 89.

*Frösch. Inf.* 1. tab. 7.

Habitat in fœtibus, casco pinguedineque alta.

*cupraria.* 52. *M.* antennis setariis glauca aenea, thorace viridi, abdomine oblongo cupreo, halteribus nudis. *Syst. Ent.* 781. 39. *Linn. Syst. Nat.* 2. 994. 92. *Fn. Sv.* 1853.

*Musca* oblonga, capite viridi, thorace aeneo, abdomine cupreo, alis macula fusca. *Geoff. Inf.* 2. 535. 61.

*Nemotelus cuprarius* antennis clauatis apice setigeris, abdomine oblongo depresso violaceo cupreo, thorace viridi aurato, alis macula fusca. *Deger Inf.* 6. 200. 16. tab. 12. fig. 4.

*Reaum. Inf.* 4. tab. 22. fig. 7. 8.

*Scop. carn.* 911.

Habitat in Europae floribus.

*polita.* 53. *M.* antennis setariis nitida, thorace coerulesco, abdomine aeneo, halteribus nudis. *Syst. Ent.* 781. 40. \*

*Musca polita* antennis setariis glabra aenea, capite viridi, thorace abdomineque coerulesco, halteribus nudis. *Linn. Syst. Nat.* 2. 894. 93. *Fn. Sv.* 1854.

*Nemotelus auratus* antennis clauatis apice setigeris, capite nudo, abdomine breui viridi aureo nitido, femoribus nigris, tibiis testaceis, alisque hyalinis. *Deger Inf.* 6. 202. 18.

Habitat in Europa boreali frequens.

*viduata.* 54. *M.* antennis setariis nigro aenea, abdomine aeneo, squamis halterum ciliatis. *Syst. Ent.* 781. 41. *Linn. Syst. Nat.* 2. 994. 94. *Fn. Sv.* 1853.

Habitat in Europa boreali.

*pubera.* 55. *M.* antennis setariis nigra, abdominis ultimo segmento incuruato, pube vtrinque reflexa. *Syst. Ent.* 781. 42. *Linn. Syst. Nat.* 2. 994. 95. *Fn. Sv.* 1855.

Habitat in Europa boreali.

*fasciata.* 56. *M.* antennis setariis testacea, abdomine nigro, alis apice fuscis, fascia alba, pedibus elongatis. *Syst. Ent.* 781. 43. \*

Habitat

**CLASS. VIII. ANTLIATA. Musca. 447**

Habitat in America meridionali. *Mus. Tottianum.*

Variat capite thoraceque nigris.

57. *M. antennis setariis liuida, fronte rubra, pedibus elongatis testaceis, geniculis nigris. Syst. Ent. 781. 44. petronella.*

*Musca Petronella* antennis setariis pilosa glabra liuida, fronte rubra, cauda incuruata, pedibus elongatis testaceis. *Linn. Syst. Nat. 2. 994. 96. Fn. Sv. 1856. It. Westrogoth. 55.*

Habitat in Europa, supra aquas quietas cursitans.

58. *M. antennis setariis atra, tibiis pallidis, vertice procrustata.*

*minulo. Syst. Ent. 782. 45.\**

Habitat in Anglia, capt. d. 20. Maii.

59. *M. antennis setariis nigra, pedibus elongatis luteis, corrigiofemoribus nigro annulo. Syst. Ent. 782. 46.\* Linn. lata.*

*Syst. Nat. 2. 995. 100.*

Habitat frequens Hafniae ad fossas paludosas.

60. *M. antennis setariis atra, pedibus elongatis pallidis, cursitans.*

*alis incumbentibus albis. Syst. Ent. 782. 47.\**

Habitat in Angliae arboribus, mares facie Cimicis celeberrime cursitans vix volitans, capt. d. 13. Maii.

61. *M. antennis setariis atra, alis incumbentibus albis, fasciis duabus nigris. It. Norwag. d. 11. Iul.\* cimicoides.*

Habitat in Norwagiae arboribus detruncatis, capt. d. 11. Iul.

62. *M. antennis setariis aeneo viridis, cauda vncinata, pedibus elongatis liuidis. Syst. Ent. 782. 49. Linn. vngulata.*

*Syst. Nat. 2. 995. 97. Fn. Sv. 1858.*

*Musca* thorace abdomineque viridi nitente, pedibus ferrugineis, alis trineruiis. *Geoff. Inf. 2. 523. 56.*

*Nemotelus aeneus* antennis clauatis apice setigeris, corpore breui, conico viridi aeneo, oculis aureis, pedibus pallide flauis. *Degeer Inf. 6. 194. 15. tab. 11. fig. 19. 20.*

Habitat ad Europae aquas frequens.

63. *M. antennis setariis aeneo viridis, cauda vncinata, alis nobilitata.*

*macula nigra apice alba. Syst. Ent. 782. 49. Linn. ta.*

*Syst. Nat. 2. 995. 98.*

*Musca*

448 CLASS.VIII. ANTLIATA. Musca.

*Musca* thorace abdomineque viridi sericeo, pedibus pallidis, alarum medio fusco. *Geoff. Inf.* 2. 523. 55.  
Habitat in Europa.

*aequifris.* 64. *M.* antennis setariis aeneo viridis, abdomine nigro fasciato, cauda vncinata, alis maculatis. *Syst. Ent.* 782. 50. \*

Habitat in Brasilia. *Mus. Dom. Banks.*

*notata.* 65. *M.* antennis setariis pilosa, thorace fusco lineato, abdomine aeneo, alis fusco nebulosis, puncto atro.

Habitat in Angliae floribus. *Mus. Dom. Teats.*

Statura omnino praecedentium. Caput nigrum. Thorax fuscus lineis quatuor virescentibus. Abdomen breue, obtusum, ouatum, aeneum. Pedes elongati, nigri. Alae fusco nebulosae fere lineatae basi ad eostam parum testaceae et versus apicem puncto distincto atro.

*arrogans.* 66. *M.* antennis setariis nigra, alis fuscis, fasciis tribus albis. *Syst. Ent.* 783. 51. *Linn. Syst. Nat.* 2. 995. 99. *Fn. Sv.* 185.

Habitat in Europa.

*longicornis.* 67. *M.* antennis setariis elongatis pilosa, aeneo nitens, abdomine obscuriore. *Syst. Ent.* 783. 52. \*

Habitat in America.

Variat pedibus obscuris et pallidis.

*rostrata.* 68. *M.* antennis setariis aenea, abdomine nigro fasciato, ore corneo prominulo atro. *Syst. Ent.* 783. 53. \*

Habitat Hafniae.

*4fasciata.* 69. *M.* antennis setariis pallida, abdomine fasciis quatuor atris, alis albis.

Habitat in Germania. *Dom. de Hattorf.*

Corpus statura omnino *M.* rostratae, at paullo minus, pallidum fasciis quatuor abdominis atris. Alae albae, immaculatae. Pedes elongati, pallidi.

*diaphana.* 70. *M.* antennis setariis, thorace aeneo, abdomine cinereo, primo secundoque segmento lateribus diaphanis. *Syst. Ent.* 783. 54. \*

Habitat in Lipsiae nemoribus.

CLASS. VIII. ANTLIATA. Musca. 449

71. *M. antennis setariis, thoracæ griseo, abdomine ferrugineo, alis punctis tribus.* *Syst. Ent.* 783. 55.

*Musca cucularia* antennis setariis pilosa, thorace plumbeo, oculis abdomineque ferrugineis, pedibus pallidis. *Linn. Syst. Nat.* 2. 995. 103.

Habitat in Europa.

Antennarum seta simplex, niuea.

72. *M. antennis setariis rufo ferruginea, alis puncto obscuriore.* *Syst. Ent.* 784. 56. *Linn. Syst. Nat.* 2. 996. 104. *Fn. Sv.* 1860. *scybalaria.*

*Musca scybalaria.* *Scop. carn.* 896.

Habitat in Europae stercore.

73. *M. antennis setariis grisea hirta, alis puncto obscuro.* *Syst. Ent.* 784. 57. *Linn. Syst. Nat.* 2. 996. 105. *Fn. Sv.* 1861. *stercoraria.*

*Musca hirsuta cinerea, alis puncto obscuriore.* *Geoff. Inf.* 2. 530. 69.

*Musca antennis plumatis pilosa hirsutissima griseo flavia, fronte rufa, alis puncto fusco.* *Degeer Inf.* 6. 88. 17.

*Reaum. Inf.* 4. tab. 27. fig. 1 — 3.

Habitat in Europae stercore.

74. *M. antennis setariis obscure plumbea, alis reticulatis, costa atra.* *Syst. Ent.* 784. 59. *marginata.*

Habitat in Saxonia.

75. *M. antennis subplumatis liuens glabra, alis puncto fimetaria obscuriore.* *Syst. Ent.* 784. 58. *Linn. Syst. Nat.* 2. 996. 106. *Fn. Sv.* 1862.

*Scop. carn.* 897.

Habitat in Europae fimetis.

76. *M. antennis setariis cinerea, capite ferrugineo, punctis duobus nigris, alarum costa nigra.* *rufifrons.*

Habitat in Italia. *D. Allioni.*

Statura omnino *M. marginatae*. Caput rufum ore vesiculoso albo punctisque duobus verticis nigris. Antennae rufae margine omni nigro setaeque porrectae albae. Thorax et abdomen obscure cinerea, subla-

*Fabr. Sp. Inf. T. II.*

F f

neata.



450. CLASS. VIII. ANTLIATA. Musca.

neata. Alae albae punctis nigris vix reticulatae costata nigra. Pedes ferruginei.

*reticulata*. 77. *M. antennis setariis pilosa cinerea, alis obsolete reticulatis, costa albo nigroque punctata.*

Habitat in Italia. *D. Allioni.*

Magnitudo *M. sinetariae*. Caput cum antennis rufum ore niueo. Thorax, abdomen, pedes cinerea, immaculata. Alae punctis nigris fere reticulatae. Costa punctis albis nigrisque variegata.

*pennipes*. 78. *M. antennis setariis pilosa, abdomine rufo, alis nigris, macula ferruginea margineque postico albo.*

Habitat in America boreali. *Mus. Dom. Blomfield.*

Magnitudo *M. domesticae*. Caput nigrum ore albo. Thorax niger antice lineis strigaeque flauis. Abdomen ouatum, rufum, immaculatum. Alae nigrae macula magna bascos oblonga ferruginea margineque postico albo. Pedes nigri tibiis posticis pennatis.

*parietina*. 79. *M. antennis setariis cinerea, alis fuscis albo punctatis maculatisque, fronte testacea. Syst. Ent. 784. 60.*

*Linn. Syst. Nat. 2. 996. 107. Fn. Sv. 1863.*

Habitat in Europae floribus.

*umbrarum*. 80. *M. antennis setariis cinerea, abdomine nigro fasciato, alis fuscis albo maculatis. Syst. Ent. 784. 61.*

*Musca umbrarum* antennis setariis, alis fuscis albo maculatis, fronte niuea. *Linn. Syst. Nat. 2. 996. 107. Fn. Sv. 1864.*

Habitat in Europae floribus.

*vibrans*. 81. *M. antennis setariis, alis hyalinis apice nigris, capite rubro. Syst. Ent. 784: 62. Linn. Syst. Nat. 2. 996. 112. Fn. Sv. 1867.*

*Musca* alis albis apice nigris. *Geoff. Inf. 2. 424. 4.*

*Musca* antennis setariis pilosa, coerulesco nigra, nitida, capite rubro, alarum apice puncto fusco. *Degm. Inf. 6. 32. 11. tab. 1. fig. 19.*

*Scop. carn. 939.*

Habitat in Europae arboretis alas continuo vibrans.

82. *M. antennis setariis, alis apice puncto laterali nigro, abdomine cylindrico. Syst. Ent. 785. 63. cynipsea.*

*Musca cynipsea* antennis setariis, alis apice puncto nigro, abdominis primo articulo obouato. *Linn. Syst. Nat. 2. 997. 113. Fn. Sv. 1868.*

*Musca* antennis setariis pilosa nigro aenea, nitidissima, capite nigro, alis versus apicem puncto nigro. *Degeer Inf. 6. 33. 12.*

*Scop. carn. 947.*

Habitat in Europae hortis.

Odorem fragrantem spargunt.

83. *M. antennis setariis atra, alis nigris apice albis. Syst. grossifica-Ent. 785. 64. Linn. Syst. Nat. 2. 996. 109. Fn. Sv. tionis. 1865.*

*Musca nigra*, alis nigris, apice macula rotunda alba. *Geoff. Inf. 2. 493. 1.*

Habitat in Europae frutetis.

84. *M. antennis setariis flauescens, abdomine atro, alis punctis duobus nigris. Syst. Ent. 785. 65. Linn. Syst. combina- ta. Nat. 2. 997. 114.*

Habitat in Europae frutetis.

85. *M. antennis setariis cinerea, abdomine atro, alis albis, punctis duobus arcuque apicis fuscis. Syst. Ent. stylata. 785. 66.\**

Habitat in Angliae nemoribus, capt. d. 23. Iun.

86. *M. antennis setariis testacea, alis maculis duabus arcu- arcuata. cuque apicis nigris.*

Habitat Kiliæ.

Affinis *M. stylatae*. Caput testaceum fronte ferruginea.

Thorax pilosus, pallide testaceus. Abdomen magis obscurum stylo ani breui, crasso, obtuso. Alae magnae obscuro maculatae maculisque duabus magnis in medio nigris. Apex niger puncto paruo albo.

87. *M. antennis setariis cinerea, ano pedibusque ferrugineis, alis maculatis. Syst. Ent. 785. 67.\* umbellatarum.*

*Musca Hgoscyanii* antennis setariis, alis vnguiculatis albis, fusco maculatis. *Linn. Syst. Nat. 2. 998. 20. Fn. Sv. 1873. ?*

*Musca Leontodontis* antennis setariis pilosa viridi grisea, oculis viridibus, antennis pedibusque flauo testaceis, alis albis fusco maculatis. *Degeer Inf.* 6. 46. 17. tab. 2. fig. 17. 18.?

Habitat in Angliae floribus umbellatis.

*graminum.* 88. *M. antennis setariis cinerea, alis fuscis, maculis albis obsoletis. Syst. Ent.* 785. 68. \*

Habitat in pratis graminosis Lipsiae.

*dorsalis.* 89. *M. antennis setariis, alis punctatis, thorace cinereo, abdomine rufo, linea dorsali nigra. Syst. Ent.* 786. 69. \*

Habitat in Bohemiae pratis.

*feminatio- nis.* 90. *M. antennis setariis, alis atris cinereo punctatis, abdomine basi subtus flauo. Syst. Ent.* 786. 70. \*

Habitat in pratis Lipsiensibus.

*picta.* 91. *M. antennis setariis fusca, alis atris, maculis vtrinque duabus oppositis punctisque duobus intermediis albis.*

Habitat in America boreali. *Mus. Dom. Blomfield.*

Statura praecedentium. Caput et thorax ferrugineo fusca. Abdomen nigrum, immaculatum. Alae magnae basi valde attenuatae, atrae basi parum ferrugineae. In medio ad vtrumque marginem maculae duae oblongae, acutae approximatae, albae et inter has puncta duo alba. Pedes picei.

*flaua.* 92. *M. antennis setariis flaua, antennis apice puncto nigro. Syst. Ent.* 786. 71.

*Musca flaua* antennis setariis flaua nuda, oculis viridissimis. *Linn. Syst. Nat.* 2. 997. 115. *Fn. Sv.* 1869.

*Musca flaua, oculis nigris. Geoff. Inf.* 2. 496. 8.

*Musca* antennis setariis pilosa tota flaua, oculis viridi aureis. *Degeer Inf.* 6. 34. 13.

Habitat in Europae floribus.

Duplo maiorem, at vix distinctam ex Italia misit *D. Allioni.*

CLASS. VIII. ANTLIATA. Musca. 433

93. *M. antennis setariis flava, abdomine punctis nigris* *apuncta-*  
quatuor. *Syst. Ent.* 786. 72. *Linn. Syst. Nat.* 2. 997. *ta.*  
116.

Habitat in Europa boreali.

94. *M. antennis setariis, alis vngiculatis albis, puncto* *aestuan-*  
centrali nigricante. *Syst. Ent.* 786. 73. *Linn. Syst.*  
*Nat.* 2. 997. 117. *Fn. Sv.* 1870.

Habitat in Europa boreali.

95. *M. antennis setariis subtus flava, supra nigra, lineis tho-* *lineata.*  
racis scutelloque flavis.

Habitat in Germaniae floribus. *Dom. de Hattorf.*

Parva: Caput flavum vertice nigro. Thorax flavus  
dorso nigro lineis quatuor scutelloque flavis. Abdo-  
men supra nigrum, subtus flavum. Alae immacu-  
latae.

96. *M. antennis setariis, alis albis, thorace virescente, ab-* *Serratu-*  
domine cinereo, lineis quatuor punctorum nigro-  
rum. *Syst. Ent.* 786. 74. *Linn. Syst. Nat.* 2. 997.  
118. *Fn. Sv.* 1871. *lae.*

Habitat in Europae Carduis.

97. *M. antennis setariis, alis albis, margine maculisque* *germina-*  
nigris. *Syst. Ent.* 786. 75. *Linn. Syst. Nat.* 2. 998. *tionis.*  
122. *Fn. Sv.* 1874.

*Scop. carn.* 940.

Habitat primo vere sub arborum foliis.

98. *M. antennis setariis, alis albis, puncto terminali fa-* *Vrticae.*  
scis tribus distinctis fuscis. *Syst. Ent.* 787. 76.  
*Linn. Syst. Nat.* 2. 998. 123. *Fn. Sv.* 1875.

Habitat in Europae Vrtica.

99. *M. antennis setariis, alis albis, fasciis fuscis inaequa-* *Cerasi.*  
libus, posticis exterius connexis. *Syst. Ent.* 787. 77.  
*Linn. Syst. Nat.* 2. 998. 124. *Fn. Sv.* 1878.

*Musca* antennis setariis pilosa rufa, pedibus flavo te-  
staceis, oculis viridibus, scutello luteo, alis albis,  
fasciis inaequalibus flexuosis fuscis. *Degeer Inf.* 6:  
50. 19.

454 CLASS. VIII. ANTLIATA. Musca.

*Blank. Inf.* tab. 16. fig. L.

*Red. riuent.* 264.

*Reaum. Inf.* 2. tab. 38. fig. 22. 23.

Habitat in Ceraforum nucleis.

*Cardui.* 100. M. antennis setariis nigra, alis albis, fascia flexuosa fusca.

*Musca Cardui* antennis setariis, alis albis, linea geminata fusca S litterae figura, oculis viridibus. *Lin.*

*Syst. Nat.* 2. 998. 126. *Fn. Sv.* 1876.

*Musca* antennis setariis pilosa nigra, capite antennisque rufis, alis albis, fascia flexuosa fusca. *Degeer Inf.* 6. 49. 18.

*Musca* alis albis linea vndulata fusca sigmoidica. *Geoff. Inf.* 2. 496. 8.

*Goed. Inf.* 1. tab. 50.

*Blank. Inf.* tab. 16. fig. T.

*Reaum. Inf.* 3. tab. 45. fig. 12 — 14.

Habitat in Cardui floribus.

*Syngenesiae.* 101. M. antennis setariis atra, alis albis, fasciis tribus abbreviatis maculaque apicis nigris.

Habitat in Germaniae floribus. *Dom. de Hattorf.*

Corpus paruum totum atrum. Alae albae, hyalinae fasciis tribus nigris, quae tamen marginem tenuiorem haud attingunt, bascos vero ad costam inferne extenditur. Ala terminatur macula atra.

*Tussilaginis.* 102. M. antennis setariis flava, alis albis, fasciis quatuor fuscis. *Syst. Ent.* 787. 78.\*

Habitat in Daniae floribus.

*Solstitialis.* 103. M. antennis setariis, alis albis, fasciis quatuor connexis nigris, scutello flavo. *Syst. Ent.* 787. 79. *Lin.*

*Syst. Nat.* 2. 999. 127. *Fn. Sv.* 1879.

*Musca* alis vnguiculatis albis, fasciis tribus fuscis, thoracis apice flavo. *Geoff. Inf.* 2. 499. 14.

*Musca Arctii* antennis setariis pilosa viridi flava, oculis aurcis, alis fasciis quaternis transuersalibus pallide fuscis. *Degeer Inf.* 6. 42. 16. tab. 2. fig. 10. 11.

Habitat in Cardui floribus.

**CLASS. VIII. ANTLIATA. Tabanus.** 455

104. *M. antennis setariis ferruginea, scutello flavo, alis Onopordinis. variegatis. Syst. Ent. 787. 80.\**

Habitat in Dania & Carduis.

105. *M. antennis setariis, alis fuscis albo variis, margine Hieracii. crassiori maculis tribus nigris, pupilla alba. Syst. Ent. 787. 81.\**

Habitat in Angliae floribus compositis.

**177. T A B A N V S.**

1. *T. oculis fuscescentibus, haustello longitudine corporis. Linn. Syst. Nat. 2. 999. 1. Mus. Lud. Vlr. 421. rostratus.*

Habitat ad Cap. bon. sp.

Hauustellum valde porrectum forte proprii generis. Abdominis segmenta apice cinereo pubescentia.

2. *T. oculis fuscescentibus, abdomine atro, margine fuluo pubescente, haustello corpore dimidio brevior. haustellatus.*

Habitat in Hispania. *Mus. Dom. Banks.*

Hauustellum setis quinque huius generis videtur corpore brevius. Thorax pubescens fuscus. Abdomen atrum nitidum margine fuluo pubescente.

3. *T. oculis virefcentibus, abdominis dorso maculis albis trigonis longitudinalibus. Syst. Ent. 788. 1. Linn. Syst. Nat. 2. 1000. 4. Fn. Sv. 1882. bovinus.*

*Tabanus* thorace cinereo, abdomine flauescente segmentis singulis triangulo albo. *Geoff. Inf. 2. 459. 1.*

*Tabanus* fuscus, antennis lunulatis, oculis viridibus, tibiis flavis, abdomine lineis transuersis maculisque trigonis albide flavis. *Degter Inf. 6. 219. 1. tab. 12. fig. 10. 11.*

*Tabanus tropicus. Harris Inf. angl. tab. 7. fig. 2.*

*Reaum. Inf. 4. tab. 17. fig. 8.*

*Schaeff. Elem. tab. 122.*

*Scop. carn. 1006.*

Habitat in Europa pecoribus infestus.

4. *T. oculis virefcentibus, thorace lineato, abdomine obscure coerulescente, antennis rufis. lineatus.*

Habitat in America. *Dom. Blackburn.*

456 CLASS. VIII. ANTLIATA. Tabanus.

Magnitudo *T. bouini*. Os tomento cinereo tectum.  
Antennae rufae. Thorax fuscus ferrugineo lineatus.  
Abdomen obscure coerulescens. Alae fuscae. Pedes  
picei.

*autumnalis*. 4. *T. alis hyalinis, abdomine fusco, ordine triplici albi-  
do maculoso. Syst. Ent. 788. 2. Linn. Syst. Nat. 2.*  
1000. 5. *Fn. Sv. 1883.*

*Harris Inf. angl. tab. 7. fig. 4.*

*Scop. carn. 1007.*

Habitat in Europa.

*fasciatus*. 6. *T. oculis viridibus, corpore brunneo, alis fascia fusca.*  
*Syst. Ent. 788. 3. \**

Habitat in Sierra Leon Africae. *Dom. Drury.*

*calens*. 7. *T. oculis viridibus, abdominis dorso linea alba, an-  
tennis rubris. Syst. Ent. 788. 4. \* Linn. Syst. Nat.*  
2. 1000. 6.

*Tabanus giganteus* fuscus, antennis lunulatis rufis,  
oculis virescentibus, scutello rufo, macula nigra, pe-  
dibus ferrugineis. *\*Degeer Inf. 6. 226. 1. tab. 30.*  
fig. 1.

Habitat in America meridionali.

*tarandinus*. 8. *T. oculis viridibus, abdominis segmentis margine la-  
teis, pedibus rufis. Syst. Ent. 788. 5. Linn. Syst.*  
*Nat. 2. 1000. 7. Fn. Sv. 1884.*

Habitat in Europa boreali, Tarandorum cornua tend-  
la pungens, vulnerans, deformans.

*feruens*. 9. *T. oculis viridibus, abdomine antennisque luteis, ca-  
pite thoraceque fuscis. Syst. Ent. 789. 6. Linn. Syst.*  
*Nat. 2. 1000. 9.*

*Tabanus* capite antennisque lunulatis flavis, thorace  
fusco lateribus flavo, abdomine flavo, apice alisque  
fuscis. *Degeer Inf. 6. 228. 4. tab. 30. fig. 4.*

Habitat in America meridionali.

*exaestuans*. 10. *T. oculis viridibus, abdominis segmentis margine al-  
bis, tibiis candidis. Syst. Ent. 789. 7. Linn. Syst.*  
*Nat. 2. 1000. 8.*

*Tabanus*

**CLASS. VIII. ANTLIATA. Tabanus. 457**

- Tabanus fuscus*, antennis lunulatis, thorace griseo, abdomine lineis transversis griseis, tibiis albis apice nigris, alis hyalinis. *Degeer Inf.* 6. 229. 8. tab. 30. fig. 5.  
Habitat Surinami.
11. *T. oculis fasciatis*, alis albis, costa fusca, antennis ruficor-  
fis. *Syst. Ent.* 789. 8.\*  
*Tabanus plumbeus*. *Drury Inf.* 1. tab. 44. fig. 2.  
Habitat in America. *Dom. Drury.*
12. *T. oculis fasciatis niger*, abdominis primo segmento *pellucens*,  
coerulefcente, tibiis candidis. *Gen. Inf. Mant.* 308.\*  
Habitat in Germania. *Dom. de Hattorf.*
13. *T. abdomine coerulefcente*, alis atris. *Syst. Ent. atratus*.  
789. 9.\*  
Habitat in America.
14. *T. oculis fascia purpurea*, corpore cinereo. *Syst. Ent. Bromius*.  
789. 10. *Linn. Syst. Nat.* 2. 1001. 12. *Fn. Su.* 1885.  
*Tabanus maculatus* antennis lunulatis, oculis viridi-  
bus, fascia pupurafcente vnica, corpore griseo, ci-  
nereo nigroque variegato. *Degeer Inf.* 6. 221. 3.  
*Schaeff. Icon.* tab. 130. fig. 6.  
*Seop. carn.* 1011.  
Habitat in Europa boreali frequens.  
Oculorum colores cum vita pereunt, aqua calida iterum  
reuiuifcunt.
15. *T. oculis fuscis cinereis*, abdomine obscuro, basi pal-  
lido pellucido. *italicus*.  
Habitat in Italia. *D. Allioni.*  
Corpus paruum statura *T. Bromii*, cinereum oculis ma-  
gnis fuscis. Abdomen obscurum basi f. primo segmen-  
to pallido pellucido lineaque dorsali cinerea. Pedes  
pallide ferruginei, apice nigri.
16. *T. oculis* - - - corpore liuido, antennis viridibus, *mexica-*  
*alis virefcentibus*, punctis anafomofantibus. *Linn. mus.*  
*Syst. Nat.* 2. 1000. 10.



938 CLASS. VIII. ANTLIATA. Tabanus.

*Tabanus oliuaceus* viridi oliuaceus, antennis lunulatis, oculis flauo aureis, pedibus viridibus, alis viriscentibus, punctis costarum fuscis. *Degeer Inf.* 6. 230. 6. tab. 30. fig. 6.  
Habitat Surinami.

*rusticus*. 17. *T. oculis glaucis cinereus, fronte punctis duobus atris.* *Syst. Ent.* 789. 11.

*Tabanus rusticus* oculis glaucis, corpore cinereo pubescente. *Linn. Syst. Nat.* 2. 1000. 11.

*Tabanus cinereus*, tibiis fulvis. *Geoff. Inf.* 2. 462. 7.  
Habitat in Europa.

*paganus*. 18. *T. oculis antice viridibus, fasciis tribus fulvis, abdomine vtrinque ferrugineo maculato.* *Syst. Ent.* 789. 12. \*

*Tabanus bouinus.* *Harris Inf. angl.* tab. 7. fig. 1.  
Habitat in Anglia, capt. d. 28. Iun.

*occidentalis*. 19. *T. oculis fasciis binis fuscis, corpore fusco, abdomine lineis tribus fulvis.* *Syst. Ent.* 790. 13. *Linn. Syst. Nat.* 2. 1001. 13.

*Tabanus fuscus*, antennis lunulatis, abdomine lineis tribus fulvis, pedibus anticis nigris, macula albida, reliquis antennisque fulvis. *Degeer Inf.* 6. 228. 3. tab. 30. fig. 3.  
Habitat Surinami.

*tropicus*. 20. *T. oculis fasciis ternis purpurascentibus, abdominis lateribus ferrugineis.* *Syst. Ent.* 790. 14. *Linn. Syst. Nat.* 2. 1001. 14. *Fn. Sv.* 1886.

*Tabanus fuscus*, antennis lunulatis, oculis viridibus, fasciis ternis purpureis, pedibus fuscis, abdominis lateribus macula magna rufa. *Degeer Inf.* 6. 220. 2. tab. 12. fig. 15.

*Tabanus sanguisorba.* *Harris Inf. angl.* tab. 7. fig. 3.  
*Schaeff. Icon.* tab. 131. fig. 4.  
*Scop. carn.* 1008.

Habitat in Europa, pecoribus molestus.

**CLASS. VIII. ANTLIATA. Tabanidæ. 459**

21. *T. oculis fasciis tribus purpurascentibus, abdomine borealis.*  
atro, segmentorum marginibus albidis. *It. Norwag.*  
d. 8. Jul. \*

Habitat in Norwagia rarius.

22. *T. oculis fasciis quaternis fuscis, tibiis albis. Syst. Ent. antarcti-*  
790. 15. *Linn. Syst. Nat. 2. 1001. 15.* *cus.*

Habitat in America calidiore.

23. *T. oculis fasciis quaternis vndatis, alis fusco puncta- pluvialis.*  
tis. *Syst. Ent. 790. 16. Linn. Syst. Nat. 2. 1001. 16.*  
*Fn. Sv. 1887.*

*Tabanus fuscus, alis cinereis, punctis numerosissimis*  
albis. *Geoff. Inf. 2. 461. 5.*

*Tabanus antennis subulatis, oculis viridibus, fasciis*  
quinis vndatis purpureis, corpore fusco cinereo, alis  
fusco maculatis. *Degeer Inf. 6. 223. 5. tab. 13. fig. 1. 2.*

*Harris Inf. angl. tab. 7. fig. 8.*

*Reaum. Inf. 4. tab. 18. fig. 1.*

*Schaeff. Icon. tab. 85. fig. 8. 9.*

*Scop. carn. 1012.*

Habitat in Europa.

24. *T. oculis maculatis, alis dimidiato fuscis, macula al- longicor-*  
ba, antennis longioribus. *Syst. Ent. 790. 17. \** *nis.*

Habitat in Brasilia. *Mus. Dom. Banks.*

25. *T. oculis - - - ferrugineus, abdomine vtrinque bidenta-*  
maculis duabus flavis, scutello bidentato. *Gen. Inf. tus.*  
*Mant. 308. \**

Habitat in Austria. *D. Schulz.*

26. *T. oculis - - - fuscus, abdomine ferrugineo, basi bispino-*  
nigro, scutello bispinoso. *Gen. Inf. Mant. 308. \** *fus.*

Habitat Goettingae. *D. Schulz.*

27. *T. oculis viridibus nigro punctatis, alis maculatis. coecu-*  
*Syst. Ent. 790. 18. Linn. Syst. Nat. 2. 1001. 17. Fn. tiens.*  
*Sv. 1888.*

*Tabanus fuscus, abdominis lateribus pedibusque flavis,*  
alis maculis fuscis. *Geoff. Inf. 2. 463. 8.*

*Taba-*

460 CLASS. VIII. ANTLIATA. Rhingia.

*Tabanus* antennis fubulatis, oculis aureis purpureo punctatis, abdomine maculis trigonis flauis, alis maculis tribus fuscis. *Degeer Inf.* 6. 224. 6. tab. 11. fig. 3.

*Tabanus nubilofus.* *Harris Inf. angl.* tab. 7. fig. 5.

*Musca bipennis pulcra*, alis maculis amplis albis piftis. *Ruf. Inf.* 272.

*Schaeff. Icon.* tab. 8. fig. 1.

*Scop. carn.* 1014.

Habitat in Europa.

*lugubris.* 28. *T.* oculis corporeque nigris, alis fuscis, macula alba. *Syst. Ent.* 791. 19. *Linn. Syst. Nat.* 2. 1001. 18. *Fa. Sv.* 1889.

Habitat in Europa.

178. RHINGIA.

*rostrata.* 1. *Rhingia*, *Syst. Ent.* 791. 1. *Scop. carn.* 972.

*Conops rostrata* antennis setariis nigra, abdomine ovato flauo laeui, rostro conico. *Linn. Syst. Nat.* 2. 1004. 1.

*Musca rostrata* antennis setariis tomentofa, abdomine ovato breui flauo, thorace nigro, capite rostratoconico. *Degeer Inf.* 6. 130. 19. tab. 7. fig. 21. 22.

Habitat in Europa frequens, pecoribus moleſta.

179. ASILVS.

*groffus.* 1. *A. hirsutus niger*, thorace abdominisque baſi cineris. *Syst. Ent.* 791. 1. \*

Habitat in America. *Muf. Dom. Banks.*

*maurus.* 2. *A. ſubferrugineus*, peſtoris lateribus thoracisque dorſo lineis tribus nigris. *Syst. Ent.* 792. 2. *Linn. Syst. Nat.* 2. 1006. 1.

Habitat in Africa.

*algirus.* 3. *A. corpore toto ferrugineo.* *Syst. Ent.* 792. 3. *Linn. Syst. Nat.* 2. 1006. 2.

Habitat in Africa.

4. *A. fronte thorace pedibusque ferrugineis, alis flavis barbarus*  
apice margineque tenuiori nigris. *Syst. Ent.* 792. 4.  
*Linn. Syst. Nat.* 2. 1007. 3.  
Habitat in Africa.

5. *A. abdomine tomentoso antice segmentis tribus nigris crabroni-*  
postice flavo inflexo. *Syst. Ent.* 2. 792. 5. *Linn. Syst. formis.*  
*Nat.* 2. 1007. 4. *Fn. Sv.* 1908.

*Afilus ferrugineus* abdominis articulis prioribus atris,  
posterioribus quatuor flavis. *Geoff. Inf.* 2. 468. 3.  
tab. 17. fig. 3.

*Afilus subhirsutus*, antennis fetigeris, abdomine antice  
nigro postice flavo fulvo. *Degeer Inf.* 6. 244. 7.  
tab. 14. fig. 3.

*Musca maxima crabroniformis.* *Raj. Inf.* 267.

*Erax crabroniformis.* *Scop. carn.* 974.

*Frisch. Inf.* 3. tab. 8.

*Reaum. Inf.* 4. tab. 8. fig. 3.

*Schaeff. Elem.* tab. 13.

— *Icon.* tab. 8. fig. 15.

Habitat in Europa frequens.

6. *A. hirsutus ater, thorace basi albo.* *Gen. Inf. Mant. ephippi-*  
308. \* *um.*

*Afilus dorsalis hirsutus niger*, antennis muticis, thora-  
ce postice villis viridi flavis. *Degeer Inf.* 6. 239. 2.  
tab. 13. fig. 9.

Habitat in Germania. *Dom. de Hattorf.*

7. *A. hirsutus niger, abdomine postice albo.* *Syst. Ent. gibbosus.*  
793. 6. *Linn. Syst. Nat.* 2. 1007. 6. *Fn. Sv.* 1909.

*Afilus Bombylius hirsutus niger*, antennis muticis, ab-  
domine postice albo griseo, capite villis griseis. *De-*  
*geer Inf.* 6. 238. 1. tab. 13. fig. 6. 7.

*Oestrus bouinum*, cauda alba. *Linn. Flor. Lapp.* 360.

*Schaeff. Icon.* tab. 8. fig. 11.

Habitat in Europa boreali.

8. *A. hirtus niger, abdomine basi fasciis duabus niueis fasciatus.*  
*Syst. Ent.* 793. 7. \*

Habitat in Sierra Leon Africae. *Dom. Drury.*

462 CLASS. VIII. ANTLIATA. *Afilus*.

*aeftuans*. 9. *A. cinereus*, vltimis abdominis tribus segmentis albis.  
*Syst. Ent.* 793. 8. *Linn. Syst. Nat.* 2. 1007. 5. *Amoen.*  
*acad.* 6. 413. 95.

Habitat in America boreali.

*ater*. 10. *A. hirsutus ater*, barba albida. *Syst. Ent.* 793. 9. *Linn.*  
*Syst. Nat.* 2. 1007. 7. *Fn. Sv.* 1910.

*Afilus totus niger subhirsutus, alis atris. Geoff. Inf.* 2.  
469. 5.

*Crux prostratus. Scop. carn.* 973.

Habitat in Europa.

*diadema*. 11. *A. ater*, alis nigris, fronte alba.

Habitat in Italia. *D. Allioni.*

Statura omnino *A. atri*, a quo tamen differt fronte sub  
antennis alba, barba vero nigra alisque totis inprimis  
tamen versus marginem crassiore nigra.

*flauus*. 12. *A. hirsutus niger*, thorace postice cinereo, abdomine  
supra hirsuto fuluo. *Syst. Ent.* 793. 10.

*Afilus flauus niger hirsutus, thorace postice flauo, ab-*  
*domine supra fuluo, plantis ferrugineis. Linn. Syst.*  
*Nat.* 2. 1007. 8. *Fn. Sv.* 1911.

*Afilus hirsutus niger, antennis muticis, thorace villis*  
*albidis, abdomine ovato villis flauo rufis splendidibus.*  
*Degeer Inf.* 6. 240. 3. tab. 13. fig. 10.

*Schaeff. Icon.* tab. 51. fig. 2.

*Linn. It. Gottl.* 327.

Habitat in Europa boreali.

Barba capitis albida.

*violaceus*. 13. *A. hirsutus ater*, abdomine violaceo. *Gen. Inf. Mant.*  
308. \*

Habitat in Germania. *Dom. de Hattorf.*

*gilvus*. 14. *A. niger*, abdomine supra fuluo. *Syst. Ent.* 793. 11.

*Afilus gilvus abdomine pubescente nigro, supra rufo.*  
*Linn. Syst. Nat.* 2. 1007. 9. *Fn. Sv.* 1912.

*Afilus niger, abdominis segmentis tribus a tergo rufis.*  
*Geoff. Inf.* 2. 468. 4.

*Afilus rufus hirsutus niger, antennis muticis, alis ni-*  
*gricantibus, abdomine supra villis rufis splendidibus.*  
*Degeer Inf.* 6. 241. 4. tab. 13. fig. 15.

*Schaeff.*

*Schaeff. Icon. tab. 78. fig. 6.*

Habitat in Europa.

Abdominis segmenta omnia excepto primo et ultimo dorso fulva.

15. *A. hirtus*, thorace cinereo pubescente; abdomine atro, punctis tribus marginalibus albis maculisque duabus dorsalibus rufis.

Habitat in Italia. *D. Allioni.*

Magnitudo et statura praecedentis. Caput fronte barbaque cinereis. Antennae nigrae. Thorax niger dorso pube cinerascens. Abdomen atrum segmento 4. 5. macula magna dorsali fulva. Puncta tria alba utrinque in margine. Pedes atr. Alae fusco ferruginae.

16. *A. halteribus* abdominisque incisuris flavis, femoribus nigris. *Syst. Ent. 793. 12. Linn. Syst. Nat. 2. 1008. 10. Fw. Sv. 1913.*

*Afilus* subhirtus niger, antennis muticis, alis fuscis, halteribus flavis, abdominis incisuris villosis flavescens. *Degeer Inf. 6. 242. 5. tab. 14. fig. 1.*

*Schaeff. Elem. tab. 23. fig. 1.*

Habitat in Europa boreali.

17. *A. corpore* cinereo immaculato. *Syst. Ent. 793. 13. \* plumbeus.*  
Habitat in nova Hollandia. *Mus. Dom. Banks.*

18. *A. hirtus* cinereus, ano pedibusque nigris. *Syst. Ent. 794. 14. forcipatus.*

*Afilus forcipatus* hirtus subcinereus, lateribus flavis.

*Linn. Syst. Nat. 2. 1008. 13. Fw. Sv. 1914.*

*Afilus* cinereus hirtus. *Geoff. Inf. 2. 473. 16.*

*Afilus cinereus* subhirtus, griseo fuscoque nigricans, antennis setigeris, thorace fascia longitudinali nigra, pedibus fuscis, abdomine elongato cylindrico. *Degeer Inf. 6. 246. 8. tab. 14. fig. 9.*

*Erax forcipatus. Scop. carn. 975.*

*Frisch. Inf. 3. 35. tab. 17.*

Habitat in Europa, in nova Hollandia. *Mus. Dom. Banks.*

464 CLASS. VIII. ANTLIATA. Afilus.

- germanicus*. 19. *A. hirtus niger, tibiis rufis, alis fuscis basi albidis*.  
*Syst. Ent.* 794. 15. \* *Linn. Syst. Nat.* 2. 1008. 12.  
*Schaeff. Icon.* tab. 48. fig. 9. 10.  
 Habitat in Germania.
- rufipes*. 20. *A. fuscus, abdomine cinereo, fascia apiceque nigra,*  
*tibiis rufis.* *Syst. Ent.* 794. 16. \*  
 Habitat in America meridionali.  
 Simillimum, at abdomine obscuriore misit *Dom. Latr.*
- maculatus*. 21. *A. cinereus, abdominis segmentis macula quadrata atra,*  
*pedibus piccis, plantis nigra.* *Syst. Ent.* 794. 17. \*  
 Habitat in India orientali. *D. Koenig.*
- marginellus*. 22. *A. cinereo fuscus, pedibus rufis apice nigris, alis mar-*  
*gine atro.*  
 Habitat in Insulae Americae St. Croix. *Dom. Vahl.*  
 Statura praecedentium. Corpus obscure cinereum, frontis barba albidula. Pedes omnes rufi apice nigri. Alae albiae margine crassiori inprimis in medio atro.
- annulatus*. 23. *A. cinereus, abdomine apice nigro, femoribus testaceis,*  
*annulo nigro.* *Syst. Ent.* 794. 18. \*  
 Habitat in India orientali. *D. Koenig.*
- stylatus*. 24. *A. hirtus cinereus, thorace abdomineque nigro maculatis,*  
*tibiis rufis, ano stylato.* *Syst. Ent.* 795. 19. \*  
 Habitat in America meridionali.
- cingulatus*. 25. *A. cinereus, tibiis rufis nigro annulatis.*  
 Habitat in Italia. *D. Allioni.*  
 Corpus minus *A. forcipato cinereum*, immaculatum.  
 Pedes nigri femoribus rufis annulis duobus nigris.
- tipuloides*. 26. *A. cinereus nudus, pedibus ferrugineis, plantis nigris.*  
*Syst. Ent.* 795. 20.  
*Afilus tipuloides* cinereus nudus, thoracis lineis dorsalibus tribus nigris. *Linn. Syst. Nat.* 2. 1008. 14.  
*Fn. Sv.* 1915.  
*Afilus* lividus, thoracis lineis dorsalibus tribus nigris.  
*Geoff. Inf.* 2. 474. 17.  
 Habitat in Europa.

27. *A. niger*, thorace fugaci aureo maculato, abdomine ~~testaceo~~,  
vtrunque maculis quinque albis. *Syst. Ent.* 795. 21.

*Linn. Syst. Nat.* 2. 1008. 11.

*Afilus tertius*. *Schaeff. Icon.* tab. 8. fig. 13.

Habitat in Germania.

28. *A. testaceus*, thorace abdomineque nigro lineatis, alis *lineatus*.  
fuscis, macula media testacea.

Habitat in Insula St. Cruz Americes. *Dom. Vahl*.

Corpus magnitudine et statura *A. oelandici*. Caput testaceum oculis rostrisque apice nigris. Thorax testaceus lineis tribus dorsalibus nigris lateralibus brevioribus. Abdomen cylindricum, incurvum testaceum maculis dorsalibus nigris. Alae fuscae macula magna media testacea. Pedes testacei.

29. *A. cinereus*, abdomine atro, segmentorum marginibus *cinctus*.  
albis.

Habitat in Germania. *Dom. de Hattorf*.

Paruus. Caput et thorax obscure cinerea halteribus flavis. Abdomen atrum nitidum segmentorum marginibus albis. Pedes nigri.

30. *A. fuscus*, abdomine alisque cyaneis. *cyaneus*.

Habitat ad Cap. bon. sp. *Mus. Dom. Banks*.

Magnitudo *A. oelandici*. Caput et thorax fusca, obscura, minime nitida. Abdomen et alae nitida, cyanea, immaculata.

31. *A. ater nudus*, pedibus halteribusque ferrugineis. *Syst. oelandici*.  
*Ent.* 795. 22. *Linn. Syst. Nat.* 2. 1008. 15. *Fn. Sv.* *cus*.  
1916.

*Afilus niger* glaber, femoribus halteribusque ferrugineis, alis nigris. *Geoff. Inf.* 2. 270. 8.

*Afilus cylindricus* glaber niger, antennis setigeris, abdomine elongato cylindrico, apice clauato, pedibus flavis, alis corpore brevioribus. *Degeer Inf.* 6. 249.  
10. tab. 14. fig. 13.

Habitat in Europa.

Variat alis nigris et hyalinis et interdum pedibus nigris.



466 CLASS.VIII. ANTLIATA. Conops.

- comopoides.* 32. A. thorace nigro flauo maculato, abdomine atro, fuscis rufis. *Syst. Ent.* 795. 23. \*  
Habitat in noua Hollandia. *Mus. Dom. Banks.*
- apliciformis.* 33. A. ater glaber, femoribus posticis longitudine abdominis. *Syst. Ent.* 796. 24. \*  
Habitat in Anglia.

180. C O N O P S.

- vesicularis.* 1. C. nigricans, occipite vesiculari, abdomine flauescente basi nigro. *Syst. Ent.* 796. 1.  
*Conops vesicularis* antennis clauatis mucronatis subferrugineis subconnexis, occipite vesiculari, abdomine glauco flauescente. *Linn. Syst. Nat.* 2. 1005. 4. *Fn. Sv.* 1903.  
*Afilus* antennis capite longioribus clauatis acuminatis nigro rufoque variis glaber, segmentis abdominalibus omnibus margine flauis, alis fuscis, margine albo. *Geoff. Inf.* 2. 472. 13.  
Habitat in Europae nemoribus.
- aculeata.* 2. C. atra abdominis incisuris thoracisque punctis duobus anticis flauis. *Syst. Ent.* 796. 2. *Linn. Syst. Nat.* 2. 1005. 6. *Fn. Sv.* 1906.  
*Conops 4fasciata* nigra, antennis nigris, abdomine fuscis quatuor transuersis apiceque flauis, pedibus rufis. *Degeer Inf.* 6. 261. 1. tab. 15. fig. 1.  
Habitat in Europa boreali.
- rufipes.* 3. C. atra, abdomine basi ferrugineo segmentorumque marginibus albis, pedibus ferrugineis.  
Habitat in Germania. *Dom. de Hattorf.*  
Statura C. aculeatae. Frontis flaua antennis fuscis. Thorax ater, immaculatus antice vtrinque prominet. Halteres flauis. Abdomen clauatum basi ferrugineum apice nigrum segmentorum marginibus albis. Pedes ferruginei. Alae costae fuscae.
- macrocephala.* 4. C. nigra, abdomine segmentis quatuor margine flauis, antennis pedibusque rufis. *Syst. Ent.* 797. 3.

CLASS. VIII. ANTLIATA. Stomoxys. 467

*Conops macrocephala* antennis clauatis mucronatis, luteis, abdomine subcylindrico glabro, segmentis quatuor margine flavescentibus. *Linn. Syst. Nat.* 2. 1005. 5. *Fn. Sv.* 1902.

*Conops rufus*, antennis pedibusque ferrugineis, abdomine fasciis fuscis flauisque, alis fusco maculatis. *Dögeer Inf.* 6. 264. 3.

*Asilus* antennis capite longioribus clauatis, acuminatis nigro rufoque variis, glaber, segmentis abdominalibus secundo et tertio margine flauis, alis fusco nebulosis. *Geoff. Inf.* 2. 471. 12.  
*Vdm. Diss.* 100.

Habitat in Europae nemoribus.

5. *C. nigra* glabra, abdomine cylindrico, segmentis tri-*flauipes*.  
bus margine flauis. *Syst. Ent.* 797. 4. *Linn. Syst.*  
*Nat.* 2. 1005. 7. *Fn. Sv.* 1904.  
Habitat in Europa boreali.

181. S T O M O X Y S.

1. *S. atra*, thorace antice ferrugineo piloso, alis atris al-*morio*.  
bo maculatis. *Syst. Ent.* 797. 1. \*  
Habitat in Brasilia. *Mus. Dom. Banks.*

2. *S. antennis* plumatis grisea, abdomine supra fusco, ma-*parasita*.  
culis baseos testaceo diaphanis.  
Habitat - - - *Mus. Dom. Teats.*  
Magna. Os albidum rostro porrecto nigro. Antennae  
valde plumatae. Thorax griseus, sublineatus. Ab-  
domen supra fuscum maculis primi secundique segmen-  
ti dorsalibus testaceo diaphanis, subtus albidum. Pe-  
des albi.

3. *S. antennis* plumatis grisea pilosa, abdominis lateribus *Siberita*.  
pallido diaphanis. *Syst. Ent.* 798. 2. \*  
Habitat Hafniae.

4. *S. antennis* subplumatis grisea, pedibus atris. *Syst. Ent.* *calci-*  
798. 3. *trans.*  
*Stomoxys.* *Geoff. Inf.* 2. 539. 1. tab. 18. fig. 2.

468 CLASS. VIII. ANTLIATA. Myops.

*Conops calcitrans* antennis subplumatis cinerea glabra ouata. *Linn. Syst. Nat.* 2. 1004. 2. *Fn. Sv.* 1900.

*Musca pungens* antennis plumatis pilosa griseo cinerea nigro maculata, rostro longo corneo porrecto. *De geër Inf.* 6. 78. 11. tab. 4. fig. 12.

*Empis calcitrans.* *Scop. carn.* 1001.

*Sulz. Inf.* tab. 21. fig. 138.

Habitat ad pedes Boum, vnde eorum continuus calcitratus: ingruente pluuiâ frenâe pungit,  
Simillima *M. domesticâ*, at rostrum omnino huius generis.

*irritans.* 5. *S.* cinerea subpilosa, abdomine nigro maculato. *Syst. Ent.* 798. 4.

*Conops irritans* cinerea subpilosa, abdomine nigromaculato. *Linn. Syst. Nat.* 2. 1004. 3. *Fn. Sv.* 1901.

Habitat in Europâ pecorum dorso copiose insidens.

182. M Y O P A.

*ferruginea.* 1. *M. ferruginea*, abdomine cylindrico incuruo, fronte latescente. *Syst. Ent.* 798. 1.

*Conops ferruginea.* *Linn. Syst. Nat.* 2. 1005. 8. *Fn. Sv.* 1907.

*Sicus ferrugineus.* *Scop. carn.* 1004.

*Astilus* antennis capite breuioribus clauatis setosis nigro rufoque varius glaber, alis nigris, oris aculeo in medio incuruato. *Geoff. Inf.* 2. 473. 14.

Habitat in Europâ nemoribus.

*testacea.* 2. *M. ferruginea*, thoracis dorso nigro.

*Conops testacea* antennis setariis testacea, abdomine subouato hamoso, facie vesiculari alba, alis hyalinis. *Linn. Syst. Nat.* 2. 1006. 11.

*Stomoxoides.* *Schaeff. Elem.* tab. 120.

Habitat in Germania.

Præcedente maior. Os vesiculosum, flauescens. Corpus ferrugineum thoracis dorso nigro. Halteres flau.

CLASS. VIII. ANTLIATA. Culex. 469

3. M. abdomine cylindrico incurvato, corpore atro. *Syst. Ent. atra.*  
*Ent.* 799. 2. \*  
Habitat in Daniae nemoribus.
4. M. testacea, abdomine hamoso albo fasciato. *Syst. Ent. cincta.*  
799. 3. \*  
Habitat in India orientali. *D. Koenig.*
5. M. ferruginea, abdomine hamoso griseo, facie vesicu- *buccata.*  
lari alba, alis nebulosis. *Syst. Ent.* 799. 4.  
*Conops buccata.* *Linn. Syst. Nat.* 2. 1006. 12. *Fn. Sv.*  
1905.  
*Sicus buccatus.* *Scop. carn.* 1005.  
Habitat in Europae nemoribus.

183. C V L E X.

1. C. cinereus, abdomine annulis fuscis octo. *Syst. Ent. pipiens.*  
799. 1. *Linn. Syst. Nat.* 2. 1002. 1. *Fn. Sv.* 1890.  
*Culex communis* nigro fuscus, abdomine annulis cince-  
reis. *Degeer Inf.* 6. 316. 1.  
*Geoff. Inf.* 2. 579. tab. 19. fig. 4.  
*Linn. Flor. Lappon.* 363. 364.  
*Blank. Inf.* tab. 16. fig. A. D.  
*Reaum. Inf.* 4. tab. 43. 44.  
*Swammerd. Bibl.* tab. 31. fig. 4 — 8.  
— — — tab. 32. fig. 1 — 5.  
*Sulz. Inf.* tab. 121. fig. a.  
*Roef. Inf.* 3. tab. 15.  
*Schaeff. Elem.* tab. 54.  
*Ledermüll. microgr.* tab. 79. 85.  
*Ioblot. micr.* 1. pp. 2. tab. 13. fig. A, E. H. I. L.  
Habitat in Europae aquis paludosis. In australiori-  
bus maior, magis venenata.  
Insectum pipiens continuo susurro puncturaque mole-  
rissimum, aues inprimis aquaticas sustentat.  
Mas antennis pectinatis vix pungit.
2. C. fuscus, thorace sublineato. *Syst. Ent.* 800. 2. *bifurca-*  
*Culex bifurcatus* fuscus rostro bifurco. *Linn. Syst. tus.*  
*Nat.* 2. 1002. 3. *Fn. Sv.* 189.

470, CLASS. VIII. ANTLIATA. Culex.

*Raj. Inf.* 74. 4.

*Reaum. Inf.* 4. tab. 40. fig. 1. 2.

*Sulz. Inf.* 21. fig. 136.

Habitat in Europae paludosis.

*lutescens.* 3. *C. flavus*, alis hyalinis, costa flavescente. *Syst. Ent.* 800. 3. \*

Habitat in Daniae paludosis.

*pulicaris.* 4. *C. fuscus*, alis albis, maculis tribus obscuris. *Syst. Ent.* 800. 4. *Linn. Syst. Nat.* 2. 1003. 4. *Fn. Sv.* 1892.

*Culex* alis maculis tribus obscuris, antennis apice bifurcis. *Geoff. Inf.* 2. 579. 2.

*Linn. Flor. Lapp.* 365.

*Derh. physicoth.* l. 1. c. 11. fig. 5. 6.

Habitat in Europa, America.

Insectum reptans, mordens punctum fuscum reliquens.

*reptans.* 5. *C. niger*, alis hyalinis, pedibus nigris, annulo alba. *Syst. Ent.* 800. 5. *Linn. Syst. Nat.* 2. 1003. 5. *Fn. Sv.* 1893.

Habitat in Europa boreali frequentissimus; tempore sereno vespertino reptatu molestissimus.

*morio.* 6. *C. ater*, alis albis, femoribus posticis clauatis serratis. *Syst. Ent.* 800. 6. \*

Habitat in Anglia, capt. d. 13. Maii.

*equinus.* 7. *C. abdomine fusco, fronte alba.* *Syst. Ent.* 801. 7. *Linn. Syst. Nat.* 2. 1003. 6. *Fn. Sv.* 1894. *Linn. Flor. Lapp.* 359.

Habitat in Europa boreali inter crines equorum cursitans.

184. E M P I S.

1. E. nigra, alis subrotundis fulco ferrugineis, *Syst. Ent. borealis.*  
801. 1. *Linn. Syst. Nat.* 2. 1003. 1. *Fn. Sv.* 1895.

*Empis nigra*, alis maximis ouatis obscure fuscis, pedibus rufis nigrisque. *Degeer Inf.* 6. 255. 2. tab. 14. fig. 17.

Habitat in Europa boreali tempore vespertino sereno choreas acreas instituens.

2. E. nigra, pedibus posticis elongatis pennatis. *Syst. Ent. pennipes.*  
801. 2.

*Empis pennipes* antennis filatis nigra, pedibus posticis longis, alterius sexus pennatis. *Linn. Syst. Nat.* 2. 1003. 2. *Fn. Sv.* 1896.

*Afilus pennipes.* *Scop. carn.* 994.

*Sulz. Inf.* tab. 21. fig. 137.

Habitat in Europa frequens in Geranii syluatici et Cardaminis pratensis floribus.

Ex Italia varietatem duplo maiorem, vix tamen distinctam, accepi.

3. E. cinerea, alis oblongis, macula costali nigra, cauda forcipata, appendiculata. *Syst. Ent.* 801. 3.

*Empis forcipata* antennis filatis cinerea, alis oblongis, cauda appendiculata. *Linn. Syst. Nat.* 2. 1004. 4. *Fn. Sv.* 1898.

Habitat in Europa boreali.

4. E. nigra, tarsis anticis incrassato ouatis. *Gen. Inf. maura.*  
*Mant.* 308. \*

Habitat Kiliae, Hamburgi supra aquas stagnantes choreas innumeras instituens.

*Femina?* pedibus simplicibus.

5. E. livida, thorace lineato, alis basi pedibusque ferrugineis. *Syst. Ent.* 801. 4.

472 CLASS. VIII. ANTLIATA. Bombylius.

*Empis livida* antennis filatis livida, thorace lineis tribus nigris. *Linn. Syst. Nat.* 2. 1003. 3. *Fn. Sv.* 1897.

*Empis griseo fusca*, thorace lineis tribus longitudinalibus nigris, pedibus rufis, alis hyalinis. *Degeer Inf.* 6. 254. 1. tab. 14. fig. 14.

Habitat in Europa in floribus Heraclei sphondylii frequens.

*cinerea.* 6. *E. cinerea*, thorace immaculato pedibus pallidis, alis apice fusciscentibus. *Syst. Ent.* 802. 5. \*

Habitat in Sueciae floribus umbellatis.

*maculata.* 7. *E. cinerea*, rostro abdominis lateribus pedibusque testaceis, alis maculatis.

Habitat in Italia. *D. Allioni.*

Statura omnino *E. cinerea*. Caput fuscum rostro basi compresso, testaceo. Thorax cinereus, sublineatus, Abdomen cinereum linea laterali testacea. Alae albae fusco maculae. Pedes testacei ungulis nigris.

*stercorea.* 8. *E. testacea*, linea dorsali nigra, alis reticulatis. *Syst. Ent.* 802. 6. *Linn. Syst. Nat.* 2. 1004. 5. *Fn. Sv.* 1899.

Habitat in Europa frequens in floribus umbellatis.

185. B O M B Y L I V S.

*maior.* 1. *B. alis dimidiato nigris sinuatis.* *Syst. Ent.* 802. 1. *Linn. Syst. Nat.* 2. 1009. 1. *Fn. Sv.* 1918.

*Bombylius variegatus* niger, villis griseis, alis dimidiato fuscis et hyalinis. *Degeer Inf.* 6. 268. 1. tab. 15. fig. 10.

*Afilus lanigerus*, alarum basi fusca. *Geoff. Inf.* 2. 466. 1.

*Reaum. Inf.* 4. tab. 8. fig. 11. 12. 13.

*Mouff. Inf.* 64. fig. 5.

*Scop. carn.* 1018.

**CLASS. VIII. ANTLIATA. Bombylius. 473**

*Raj. Inf.* 273.

*Schaeff. Icon.* tab. 79. fig. 5.

*Huffnag. Inf.* tab. 8. fig. 1.

Habitat in Europa mel florum volitando hauriens.

2. B. alis dimidiato nigris aequalibus.

*aequalis.*

Habitat in America boreali. *Mus. Dom. Teats.*

Differt sane a praecedente duplo minor. Corpus cinereo hirtum. Alae albae margine crassiori late nigro, at colore haud sinuato, sed recto, aequali.

3. B. alis fusco punctatis, corpore flauescente postice albo. *Syst. Ent.* 802. 2. *Linn. Syst. Nat.* 2. 1009. 2. 1919. *medius.*

*Bombylius punctatus* niger villis fulvis, alis fusco punctatis. *Degeer Inf.* 6. 269. 2. tab. 15. fig. 12.

*Petiv. Gazoph.* tab. 36. fig. 5.

*Schaeff. Elem.* tab. 27. fig. 1.

— *Icon.* tab. 78. fig. 3.

*Scop. carn.* 1019.

Habitat primo vere in Europae aruis.

4. B. alis nigro maculatis, corpore cinerascente postice nigro. *Syst. Ent.* 802. 3. *Linn. Syst. Nat.* 2. 1009. 3. *capensis.*  
*Mus. Lud. Vlk.* 423.

Habitat in Africa.

5. B. alis immaculatis. *Syst. Ent.* 803. 4. *Linn. Syst. Nat.* 2. 1009. 4. *minor.*  
*Fn. Sv.* 1920.

*Scop. carn.* 1020.

*Petiv. Gazoph.* tab. 42. fig. 9.

*Reaum. Inf.* 4. tab. 8. fig. 11 — 13.

*Schaeff. Icon.* tab. 46. fig. 9.

Habitat in Europae aruis.

6. B. alis basi feminigris ater, abdomine albo maculato. *Syst. Ent.* 803. 5. *Linn. Syst. Nat.* 2. 1010. 5. *ater.*



Habitat in Italia. *D. Allioni.*

Statura et magnitudo omnino praecedentis, at in  
latus. Alae basi atrae, apice fuscae.

*pygmaeus.* 9. B. alis dimidiato punctisque nigris, thorace ful  
apiceque albo.

Habitat in America boreali. *Mus. Dom. Feats.*

Minutus. Caput nigrum. Thorax villosus, fuscu  
gine antico et postico albis. Abdomen hirtu  
rugineum. Alae secundum costam dimidiato  
et praeterea puncta nigra sparsa. Pedes ferri

## 186. HIPPOBOSCA.

*equina.* 1. H. alis obtusis, thorace albo variegato, pedibus  
dactylis. *Syst. Ent.* 803. L. *Linn. Syst. Nat.* 2  
1. *Fn. Sv.* 1921.

*Hippobosca* pedibus tetradactylis, alis cruciatis.  
*Inf.* 2. 547. 1.

*Hippobosca* fusca, alis magnis fuscis, vnguibus f  
cibus. *Degeer Inf.* 6. 275. 1. tab. 16. fig. 1.

*Scop. carn.* 1022.

*Sulz. Inf.* tab. 21. fig. 141.

*Reaum. Inf.* 6. tab. 48.

*Schaeff. Inf.* tab. 170. fig. 2.

**CLASS. VIII. ANTLIATA. Pycnogonum. 475**

2. H. alis obtusis, thorace unicolore. *Syst. Ent.* 804. 2. *auicularia*.  
*Linn. Syst. Nat.* 2. 1010. 2. *Fn. Sv.* 1922.  
*Hippobosca* viridis, alis magnis hyalinis, vnguibus appendiculatis. *Degeer Inf.* 6. 285. 2. tab. 16. fig. 21. 22.  
Habitat in Auiibus, Hirundinibus.
  
3. H. alis subulatis, pedibus hexadactylis. *Syst. Ent.* 804. *Hirundinis*.  
3. *Linn. Syst. Nat.* 2. 1010. 3. *Fn. Sv.* 1923.  
*Hippobosca* pedibus hexadactylis, alis diuaticis. *Geoff.*  
*Inf.* 2. 547. 2.  
*Scop. carn.* 1023.  
*Reaum. Inf.* 4. tab. 11. fig. 1 — 5.  
*Schaeff. Elem.* tab. 70.  
— — *Icon.* tab. 53. fig. 1. 2.  
Habitat in Hirundinibus.
  
4. H. alis nullis. *Syst. Ent.* 804. 4. *Linn. Syst. Nat.* 2. *ouina*.  
1011. 4. *Fn. Sv.* 1923.  
*It. Westrogoth.* 59.  
*Frisch. Inf.* 5. tab. 18.  
Habitat inter ouium lanam.

**187. PYCNOGONVM.**

1. P. palpis duobus, corpore ouato. *It. Norwag. d.* 3. *Balaenarum*.  
Aug. \*
  
- Pediculus Balaenarum* abdomine dilatato muricato, rostro porrecto subulato. *Syst. Ent.* 810. 35.  
*Pycnogonum.* *Brünnich. Inf.* tab. 1. fig. 17.  
*Phalangium Balaenarum* abdomine dilatato muricato, rostro subulato. *Linn. Syst. Nat.* 2. 1028. 6.  
*Phalangium littorale.* *Stroem. Sundm.* 209. tab. 1. fig. 17.  
*Pediculus Ceti.* *Baister. subsc.* 2. 146. tab. 12. fig. 3.  
Habitat in Oceano Norwagico sub lapidibus.

*Phalangium gygys* corpore minuto cylindrico  
meris tuberculatis, pedibus longissimis. *Lin.*

*Nat.* 2. 1027. 1.

*Phalangium marinum.* *Stroem. Sandm.* 208. 1  
fig. 16.

Habitat in mari Norwagico, capt. d. 20. Aug.

Animal tardum Mytilorum testas penetrat, exhan-

## 188. P E D I C V L V S.

*humanus.* 1. P. abdomine lobato cinereo. *Syst. Ent.* 804. 1.

*Syst. Nat.* 2. 1016. 1. *Fn. Sv.* 1939.

*Red. Exper.* tab. 18.

- - *Oper.* tab. 25.

*Swammerd. Bibl.* tab. 1. fig. 3 — 6.

*Alb. Aran.* tab. 42.

*Bonann. microgr.* fig. 55.

*Baker. microsc.* tab. 13. fig. 4.

*Schaeff. Elem.* tab. 95.

*Schrank. Beytr.* 112. 1.

*Sulz. Inf.* tab. 22. fig. 145.

*Ledermüll. microsc.* 45. tab. 21.

Habitat in capite et vestimentis humanis. In pueri  
lois frequentissimus morbos auertens.

Varietas capitis durior, coloratior, vestimentorum  
xior, magis cinerea.

**CLASS. VIII. ANTLIATA. Pediculus. 477**

*Raj. Inf. 8.*

*Red. Exper. tab. 19. fig. 1.*

Hospitatur in hominis immundi pube, superciliis, pel-  
litur oleo Tabaci, mercurialibus.

3. P. abdomine orbiculato, linea alba, scutello trilobo, *ricinoidis*.  
rostro albo. *Syst. Ent. 805. 3. Linn. Syst. Nat.*  
2. 1017. 3.

Habitat in America pedes obambulantium intrans, san-  
guinem hauriens, in iis oua deponens vlcera cacoe-  
thica caussans.

4. P. capite porrecto acuminato ferrugineus, abdomine *Cameli*.  
ouato. *Syst. Ent. 805. 4.*

*Pediculus Cameli. Linn. Syst. Nat. 2. 1017. 6.*

*Red. Exper. tab. 22.*

- - *Oper. tab. 25.*

*Albin. aran. 76. tab. 51.*

Habitat in Africae Camelis.

5. P. Cerui. *Syst. Ent. 805. 5. Linn. Syst. Nat. 2. 1017. 7. Cerui.*  
*Fn. Sv. 1944.*

*Frisch. Inf. 12. tab. 5.*

*Red. Exper. tab. 5.*

Habitat in Ceruo Elapho, Dama.

6. P. flauescens, striis fuscis obscuris, abdomine tubercu- *Bufali.*  
lis decem conicis marginalibus. *Degeer Inf. 7. 68.*  
3. tab. 1. fig. 11. 12.

Habitat in Bufalis Cap. bon. sp.

Pediculo humano paullo minor.

7. P. abdomine lineis transuersis 6to ferrugineis. *Syst. Bonis.*  
*Ent. 805. 6. Linn. Syst. Nat. 2. 1017. 9. Fn. Sv.*  
1946.

Habitat in Europae Bobus, Pellitur decocto Ledi et  
Sedi acris.

478 CLASS. VIII. ANTLIATA. *Pediculus*.

*Vituli*. 8. *P. abdomine acuminato plumbeo. Syst. Ent.* 805. 7.  
*Linn. Syst. Nat.* 2. 1018. 10. *Fn. Sv.* 1947.  
 Habitat in Europae Bobus.

*Equi*. 9. *P. Equi caballi. Syst. Ent.* 806. 8. *Linn. Syst. Nat.* 2.  
 1018. 11. *Fn. Sv.* 1948.  
 Habitat in Europae Equis.

*Afini*. 10. *P. capite porrecto obtuso, abdomine ovato fusco stria-*  
*to. Syst. Ent.* 806. 9.  
*Pediculus Afini. Linn. Syst. Nat.* 2. 1018. 12.  
*Albin. aran.* 76. tab. 51.  
*Red. Exper.* tab. 22. fig. 1.  
 - - *Oper.* tab. 24.  
 Habitat in Europae australioris Afinis.

*Vulturis*. 11. *P. oblongus, abdomine fusco, lineis duabus flavesce-*  
*ntibus. Syst. Ent.* 806. 10. \*  
 Habitat in Indiae orientalis Vulturibus.

*Tinnunculi*. 12. *P. capite sagittato, postice vtrinque mucronato. Syst.*  
*Ent.* 806. 11.  
*Pediculus Tinnunculi. Linn. Syst. Nat.* 2. 1018. 13. *Fn.*  
*Sv.* 1949.  
*Alb. aran.* tab. 43.  
*Red. Exper.* tab. 13.  
*Frisch. Inf.* 11. 24. tab. 24.  
 Habitat in Falconibus, Tinnunculis.

*Buteonis*. 13. *P. abdomine marginato, segmentis vtrinque punctis*  
*duobus impressis. Gen. Inf. Mant* 309. \*  
*Pediculus maximus. Scop. carn.* 1036.  
 Habitat in Falcone Buteone.

*Strigis*. 14. *P. abdomine ovato albo, margine pedibusque rubris.*  
*Syst. Ent.* 806. 12.  
*Pediculus haematopus. Scop. carn.* 1035.  
 Habitat in Strigibus, Falconibus.

CLASS. VIII. ANTLIATA. Pediculus. 479

15. P. abdomine ovato, margine striato. *Syst. Ent.* 806. *Corui.*

13. *Linn. Syst. Nat.* 2. 1018. 14. *Fn. Sv.* 1050.

*Pediculus* albo nigroque varius, abdomine ovato oblongo vtrinque incisuris nigris. *Geoff. Inf.* 2. 600. 8.

*Pediculus Cornicis* albidus, fasciis dorsalibus transversis fuscis, antennis brevissimis ad apicem capitis. *Degeer Inf.* 7. 76. 3. tab. 4. fig. 11.

*Red. Exper.* tab. 16. fig. 2.

- - *Oper.* tab. 14.

Habitat in Coruis.

16. P. abdomine ovato pallido, maculis lateralibus nigris, *Cornicis*. pupilla alba. *Syst. Ent.* 807. 14.

*Pediculus ocellatus. Scop. carn.* 1038.

Habitat in Cornice.

17. P. thorace angustato, abdomine ovato pallido diaphano. *Syst. Ent.* 807. 15.

*Pediculus infausti. Linn. Syst. Nat.* 2. 1018. 15. *Fn. Sv.* 1951.

Habitat in Coruo infausto.

18. P. capite obtuso fusco, punctis quatuor nigris. *Syst. Picae. Ent.* 807. 16.

*Pediculus Picae. Linn. Syst. Nat.* 2. 1018. 16.

*Alb. Aran.* 76. tab. 45.

*Red. Exper.* tab. 5.

Habitat in Coruo Pica.

19. P. albus, abdomine linea dorsali fusca. *Gen. Inf. Mant.* *Orioli.* 309. \*

*Pediculus dolichocephalus. Scop. carn.* 1039.

Habitat in Coracia Oriolo.

20. P. abdomine oblongo albido fusco fasciato. *Syst. Ent.* *Cuculi.* 807. 17.

*Pediculus fasciatus. Scop. carn.* 1040.

Habitat in Cuculo canoro.

480 CLASS. VIII. ANTLIATA. Pediculus.

- Cygni.** 21. P. capite emarginato, abdomine ovato, margine nigro striato. *Syst. Ent.* 307. 13.  
*Pediculus Cygni.* Linn. *Syst. Nat.* 2. 1018. 17.  
*Albin. aran.* 76. tab. 58.  
*Red. Oper.* tab. 20.  
Habitat in Cygno.
- Anseris.** 22. P. filiformis pallidus, margine nigro punctato. *Syst. Ent.* 307. 19.  
*Pediculus anatis anseris.* Linn. *Syst. Nat.* 2. 1018. 18.  
*Fn. Sv.* 1952.  
*Alb. aran.* 76. tab. 58.  
*Red. Exper.* tab. 10.  
Habitat in Anseribus feris mansuetisque. Pellitur pere.
- Mergi.** 23. P. albidus, capite flauescente, corpore elongato.  
*Ricinus Mergi ferrati.* *Deger Inf.* 7. 78. 5. tab. 4. fig. 13.  
Habitat in Mergis ferratis.
- Procellariae.** 24. P. elongatus filiformis fuscus, pedibus pallidis. *Syst. Ent.* 308. 20.  
Habitat in Brasiliae Procellariis. *Mus. Dom. Banks.*
- Diomedae.** 25. P. capite obtuso, albus, abdominis lateribus nigris. *Syst. Ent.* 308. 21.\*  
Habitat in Brasiliae Diomedeis. *Mus. Dom. Banks.*
- Sternae.** 26. P. capite trigono, abdomine ovato pallido, dorso longitudinaliter nigricante. *Syst. Ent.* 308. 22.  
*Pediculus Sternae.* Linn. *Syst. Nat.* 2. 1019. 21. *Fn. Sv.* 1953.  
*Ricinus Lari* griseo albidus, oculis maculisque capitis nigris pedibus crassioribus. *Deger Inf.* 7. 77. 4. tab. 4. fig. 12.

**CLASS. VIII. ANTLIATA. Pediculus. 481**

*Red. Exper.* tab. 9. fig. 2.

Habitat in Sternis, Laris.

27. *P. abdomine subclauato pallido, spiraculis niueis. Gruis.*  
*Syst. Ent.* 809. 23.

*Pediculus Gruis. Linn. Syst. Nat.* 2. 1019. 24. *Fn. Sv.* 1954.

*Red. Exper.* tab. 3.

- - *Oper.* tab. 19.

*Frisch. Inf.* 5. 15. tab. 4.

Habitat in Gruibus.

28. *P. elongatus filiformis, abdomine albo, lateribus nigro punctatis. Ciconiae.*  
*Syst. Ent.* 808. 24.

*Pediculus Ciconiae. Linn. Syst. Nat.* 2. 1019. 25. *Fn. Sv.* 1955.

*Frisch. Inf.* 8. 9. tab. 6.

Habitat in Ciconiis.

29. *P. oblongus fuscus, capite trigono canaliculato. Systr. Recurui-  
Ent.* 808. 25. *rostrae.*

*Pediculus recuruirostrae. Linn. Syst. Nat.* 2. 1019. 28.  
*Fn. Sv.* 1956.

Habitat in Recuruirostris.

30. *P. glaucus, thorace angustissimo, abdomine ovato, incisuris pallidis. Haema-  
Syst. Ent.* 809. 26. *topi.*

*Pediculus Haematopi. Linn. Syst. Nat.* 2. 1019. 29.  
*Fn. Sv.* 1957.

Habitat in Haematopo Ostralega.

31. *P. capite globoso maximo, corpore pallido fusco-que striato. Pauonis.  
Syst. Ent.* 809. 27. *Linn. Syst. Nat.* 2. 1019. 30.

*Geoff. Inf.* 2. 600. 9.

*Albin. aran.* 76. tab. 47.

*Fabr. Sp. Inf. T. II.*

H h

*Frisch.*



482 CLASS. VIII. ANTLIATA. Pediculus.

*Frisch. Inf.* tab. 3. fig. 6.

*Red. Exper.* tab. 15.

- - *Oper.* tab. 23.

Habitat in Pauonibus.

*Meleagris* 32. *P.* capite trigono obtuso, abdomine ovato griseo. *Syst. Ent.* 809. 28.

*Pediculus Meleagridis.* *Linn. Syst. Nat.* 2. 1020. 3. *Fn. Sv.* 1958.

*Frisch. Inf.* 8. tab. 4.

Habitat in Gallopauonibus.

*Gallinae.* 33. *P.* thorace capiteque vtrunque mucronatis. *Syst. Ent.* 809. 29. *Linn. Syst. Nat.* 2. 1020. 32. *Fn. Sv.* 1959.

*Geoff. Inf.* 2. 601. 11.

*Schrank. Beytr.* 114. 3.

Habitat in Gallinis domesticis.

Pellitur Pipere.

*Phasian.* 34. *P.* capite ovato magno, abdomine globoso obtuso. *Gen. Inf. Mant.* 310. \*

*Red. Oper.* 1.

Habitat in Phasiano Motmot.

*Caponis.* 35. *P.* abdominis margine nigro. *Syst. Ent.* 809. 30. *Linn. Syst. Nat.* 2. 1020. 33. *Fn. Sv.* 1960.

*Pulex Capi.* *Red. Exper.* tab. 16. fig. 1.

*Pediculus Galli.* *Frisch. Inf.* 11. tab. 24.

*Geoff. Inf.* 2. 601. 10.

Habitat in Gallina domestica.

*Columbae.* 36. *P.* corpore filiformi ferrugineo postice clauato. *Syst. Ent.* 809. 31.

*Pediculus Columbae.* *Linn. Syst. Nat.* 1020. 36.

CLASS. VIII. ANTLIATA. Pediculus. 483

*Pediculus oblongus*, filiformis albicans, corporis lateribus vtrinque ferrugineis. *Geoff. Inf.* 2. 599. 7.

*Red. Exper.* tab. 2. fig. 1.

*Schränk. Beytr.* 114. 4. tab. 5. fig. 3.

*Alb. aran.* tab. 43.

Habitat in Columbis domesticis.

37. *P. pallide fuscus*, abdomine albescente, maculis lateribus fuscis, capite triangulo. *Emberizae.*

*Ricinus Emberizae. Degeer Inf.* 7. 74. 2. tab. 4. fig. 9.

*Pediculus subflaescens*, abdomine ovato medio diaphano, macula fusca, lateribus punctato ferrugineis. *Geoff. Inf.* 2. 599. 6.

Habitat in Europae Emberizis.

38. *P. corpore ovato*, cauda quadriseta. *Syst. Ent.* 809. *Pari.*  
\* 32. *Linn. Syst. Nat.* 2. 1020. 37.

*Acarus ovatus* pallidus, ano pilis quatuor corporis longitudine, pedibus anticis quatuor crassioribus, omnibus physopodibus. *Schränk. Beytr.* 38. 17. tab. 1. fig. 36.

Habitat in Europae Paris.

Pedes octo numerat *Schränk.* Forte potius *Acarus*.

39. *P. capite cordato rufo*, abdomine albo basi attenuato. *Motacillae.*  
*Gen. Inf. Mant.* 310.\*

*Pediculus albiuentris. Scop. carn.* 1051.

Habitat in Motacilla Troglodyte.

40. *P. pallescens*, abdomine albo nigro maculato. *Syst. Hirundinis.*  
*Ent.* 810. 33.

*Pediculus Hirundinis. Linn. Syst. Nat.* 2. 1020. 38.

*Fn. Sv.* 1963.

Habitat in Hirundine apode.

484 CLASS. VIII. ANTLIATA. Acarus.

*Gryllotalpa*. 41. *P. sanguineus*, pedibus albis. *Gen. Inf. Mantiss. pae.* 310.\*

*Pediculus rostratus*. *Scop. carn.* 1052.

Habitat in *Acheta Gryllotalpa*.

*Apis*. 42. *P. filiformis*, ferrugineus. *Syst. Ent.* 810. 34.

*Pediculus Apis*. *Linn. Syst. Nat.* 2. 1020. 40.

*Frisch. Inf.* 8. tab. 16.

Habitat in Europae Apibus.

189. A C A R V S.

*Elephantinus*. 1. *A. orbicularis depressus liuidus*, macula bascos ouata nigra. *Syst. Ent.* 810. 1. *Linn. Syst. Nat.* 2. 1022. 1.

*Acarus ouatus* macula bascos nigra triloba vtringue fulcis tribus longitudinalibus. *Schrank. Beytr.* 124. 29. tab. 6. fig. 14. 15.

Habitat in India.

Conf. *Acarus grossus*. *Pall. spic. Zool.* 9. 43. tab. 4. fig. 12. An distinctus?

*Ricinus*. 2. *A. globoso ouatus*, macula bascos rotunda, antennis clauatis. *Syst. Ent.* 810. 2. *Linn. Syst. Nat.* 2. 1023. 7. *Fn. Sv.* 1967.

*Acarus liuidus*, antennis breuibus subclauatis, abdomine antice macula ouata fusca nitente. *Geoff. Inf.* 2. 621. 1.

*Acarus ricinoides violaceo niger*, capite pedibusque fulcis, corpore globoso ouato. *Degeer Inf.* 7. 94. 4. tab. 5. fig. 16. 17.

*Ricinus caninus*. *Raj. Inf.* 10.

*Scop. carn.* 1057.

*Aldrov. Inf.* 550.

Habitat frequens in Bobus, canibus.

3. *A. obouatus planus*, macula bascos obouata. *Gen. Inf. Reduuius.*  
*Mant.* 310. *Linn. Syst. Nat.* 2. 1022. 3. *Fn. Sv.* 1966.  
*It. Gottl.* 62. 126.

*Acarus obouatus planus*, corpore antice clypeo nigro rotundo, pedibus nigris. *Degeer Inf.* 7. 101. 5.  
 tab. 6. fig. 1. 2.

*Pediculus ouinus.* *Raj. Inf.* 9.

*Frisch. Inf.* 6. tab. 19.

Habitat in Bobus, canibus.

4. *A. ouatus fuscus*, antennis pedibusque pallidis. *pallipes.*  
 Habitat Kiliae. *Dom. Kerstens.*

Magnitudo fere *A. Ricini*. Corpus totum ouatum, depressum, planum, fuscum pilis rarioribus pallidis adpersum, antennis pedibusque pallidis.

5. *A. ouatus planus niger*, abdomine ferrugineo, antennis clauatis. *It. Norwag.* d. 9. Aug. \* *sanguis-*  
*gus.*

Habitat in hominibus, animalibus Norwagiae.

6. *A. obouatus niger*, margine albo. *Syst. Ent.* 811. 3. *aegypti-*  
*Linn. Syst. Nat.* 2. 1022. 2. *Mus. Lud. Vlr.* *us.*  
 425.

Habitat in Aegypto.

7. *A. orbiculatus ater*, lateribus vndato albis, puncto *vndatur.*  
 nigro. *Syst. Ent.* 811. 4. \*

Habitat in noua Hollandia. *Mus. Dom. Banks.*

8. *A. ouatus fuscus*, margine incrassato. *Syst. Ent. Lipsiensis.*  
 811. 5. \*

Habitat Lipsiae.

486 CLASS. VIII. ANTLIATA. Acarus.

*Iguanae.* 9. *A. ouatus planus aureo maculatus, abdominis margine striato subcrenato. Syst. Ent. 811. 6.\**

Adhaesit mordax sacco gulari Lacertae Iguanae ex America in spiritu vini missae.

*lineatus.* 10. *A. ouatus ferrugineus, lineis duabus vndatis albis. Syst. Ent. 811. 7.\**

Habitat in America.

*aureolatus.* 11. *A. obouatus fuscus, lineolis duabus maculaque postica palmata viridi aureis. Gen. Inf. Mant. 310.\**

*Acarus aureolatus. Pall. spicil. Zool. fasc. 9. p. 41. tab. 3. fig. 10.*

Habitat in America.

*Indus.* 12. *A. ovalis ferrugineus, macula bascos ouata atra. Syst. Ent. 812. 8. Linn. Syst. Nat. 2. 1022. 4.*

Habitat in India.

*americana-nus.* 13. *A. obouatus rubicundus, scutello geniculisque pedum albis. Syst. Ent. 812. 9. Linn. Syst. Nat. 2. 1022. 5.*

*Acarus Nigua ouatus planus ruber macula dorsali alba, geniculis pedum albidis. Degeer Inf. 7. 153. 1. tab. 37. fig. 9. 10.*

*Kalm. Act. Holm. 1754. 9. 10.*

Habitat in Americae Bobus, Equis.

*crassipes.* 14. *A. pedibus secundi paris crassissimis. Syst. Ent. 812. 10. Linn. Syst. Nat. 2. 1023. 8. Fn. Sv. 1969.*

*Acarus pedibus secundis crassissimis, corpore toto testaceo. Schrank. Beytr. 123. 23. tab. 6. fig. 4.*

Habitat

**CLASS. VIII. ANTLIATA. Acarus. 487**

Habitat in Europae terra, gregarius, agilis in hortorum terra praegnanse frequens.

15. A. thorace angulato cruciato, pedibus vnguiculatis *Vespertilio* corpore longioribus. *Syst. Ent.* 812. 11. *Linn. Syst. Nat.* 2. 1023. 9.

*Pediculus Vespertilionis.* *Linn. Fn. Sv.* 1941.

*Scop. carn.* 1058.

*Frisch. Inf.* 7. tab. 7.

Habitat in Vespertilione murino.

In plano incedere haud valet.

16. A. femoribus tertiis crassissimis. *Syst. Ent.* 812. 12. *passerinus.* *Linn. Syst. Nat.* 2. 1023. 10. *Fn. Sv.* 1970.

*Acarus vespertilionis fuscus*, ouatus pedibus pallidis. *Geoff. Inf.* 2. 627. 14.

*Acarus pedibus tertii paris magnis et crassissimis.* *Degeer Inf.* 7. 109. 7. tab. 6. fig. 12.

*Pediculus Vespertilionis.* *Baeker. micr.* tab. 15.

*Degeer Aët. Holm.* 1740. 341. tab. 1. fig. 2.

*Red. viv.* tab. 2. fig. 4.

*Schränk. Beytr.* 7. 3.

Habitat in Passeribus variis.

17. A. flauus, pedibus primis longissimis motatoriis. *motatorius.* *Syst. Ent.* 812. 13. *Linn. Syst. Nat.* 2. 1023. 11. *Fn. Sv.* 1971.

*Acarus rufus* pedibus primi paris longissimis, abdomine postice crenato. *Schränk. Beytr.* 8. 4. tab. 1. fig. 8. 9.

Habitat in fungis agilis, pedibus anticis vibrans.

*coleoptra*-18 A. ater, lateribus angulato acutis.

tus.

*Acarus coleoptratus* ater, lateribus nigro subcoleoptratis. *Linn. Syst. Nat.* 2. 1023. 13. *Fn. Sv.* 1973.

*Acarus marginatus* oustus crustaceus fusco castaneus nitidus, corporis lateribus margine plano angulato.

*Degeer Inf.* 7. 133. 20. tab. 8. fig. 6.

*Acarus ater*, lateribus abdominis antrorsum acutis. *Geoff. Inf.* 2. 626. 12.

Habitat in Europae arboribus sub cortice.

*Aphidioi*-19. A. ruber, pedibus primis longissimis cursoriis, abdomine postice bicorni. *Syst. Ent.* 813. 14. *Linn. Syst.*

*Nat.* 2. 1023. 12. *Fn. Sv.* 1972.

*Acarus petrarum* ruber, pedibus anticis longitudine corporis. *Geoff. Inf.* 2. 625. 10.

*Acarus saltatorius* ano bicaudato. *Schrank. Beytr.* tab. 1. fig. 33. 34.

Habitat in ligno putrescente.

*talaris*. 20. A. rubicundo hyalinus, abdomine vtrinque macula fusca. *Syst. Ent.* 813. 15. *Linn. Syst. Nat.* 2. 1023. 14. *Fn. Sv.* 1974.

*Acarus fuscus autumnalis.* *Geoff. Inf.* 2. 626. 11.

*Acarus oblongus* albouirescens s. flauescens, pedibus omnibus aequalibus. *Degeer Inf.* 7. 128. 18. tab. 7. fig. 20. 21. 22. 23.

*Acarus ellipitoideus*, pedibus secundi tertiique parvis distantibus. *Schrank. Beytr.* 35. 14. tab. 1. fig. 31. 32.

Habitat in Europae plantis inclusis tela ducens filis parallelis, quibus plantas hybernaculorum suffocat. In Tiliae foliis autumnno frequens.

21. *A. albidus*, femoribus capiteque ferrugineis, abdomine fetoso. *Syst. Ent.* 8.3. 16. *Linn. Syst. Nat.* 2. 1024. 15. *Fn. Sv.* 1947. *Siro.*

*Acarus Farinae* oblongus albus, capite rufescente, pedibus conicis crassissimis aequalibus. *Degeer Inf.* 7. 97. 3. tab. 5. fig. 15.

*Acarus lateribus sublobatis*, abdomine bifeto, fetis corpore brevioribus, pedibus quatuor posticis gracilibus. *Schrank. Beytr.* 121. 22. tab. 6. fig. 2.

*Blank. Inf.* tab. 14. fig. A. B.

*Ledermüll. micr.* 68. tab. 33. fig. 2.

*Bonann. micr.* fig. 112.

*Lewwenh. epist.* 77. tab. 370. fig. 9. 10.

*Rivin. prurit.* 18. fig. D. E.

*E. N. C.* dec. 2. ann. 10. app. 34.

Habitat in Farina, Caseo diutius asseruatis.

Conf. *Acarus domesticus* albus maculis binis fuscis, corpore ovato medio coarctato, pilis longissimis, pedibus aequalibus. *Degeer Inf.* 7. 789. 1. tab. 5. fig. 1 — 4. vix satis distinctus.

22. *A. albus*, pedibus rufescentibus, posticis quatuor seta longissima. *Scabiei.*

*Acarus Scabiei* subrotundus albus, pedibus rufescentibus brevibus, posticis quatuor seta longissima, plantis quatuor anticis fistulatis capitulo terminatis. *Degeer Inf.* 7. 94. 2. tab. 5. fig. 12. 13.

*Acarus humanus subcutaneus.* *Geoff. Inf.* 2. 622. 2.

*Baker. microsc.* 193. tab. 13. fig. 2. a. b. c.

*At. angl. abbr.* 5. 197. 283. tab. 9. fig. 127. 128.

*Bonann. microsc.* 113.

*Rivin. prur.* 18. fig. A. B.



490 CLASS. VIII. ANTLIATA. Acarus.

Habitat in Vlceribus Scabiei cutem hominis rugas sequens, penetrat, titillationem et vesiculam excitat. Causam nec symptoma morbi esse, euincunt obseruata, analogia cum gallis, contagium, cura. Praecedente multo minor et distinctus.

Conf. *Acarus* exulcerans pedibus longissimis setaceis, anticis duobus breuibus. *Syst. Ent.* 814. 19. *Linn. Syst. Nat.* 2. 1024. 18. *Fn. Sv.* 1976. Forte eiusdem speciei.

*lactis.* 23. *A.* abdomine ouato obtuso, postice setis quatuor longitudine corporis declinatis. *Syst. Ent.* 813. 17. *Linn. Syst. Nat.* 2. 1024. 16.

Habitat in cremore lactis diu conseruato et in lactis vasis immundis.

*Dysenteriae.* 24. *A.* pedibus setis duabus, abdominis ouati postica setis quatuor longitudine corporis horizontalibus. *Syst. Ent.* 814. 18. *Linn. Syst. Nat.* 2. 1024. 17. *Amoen. acad.* 5. 97.

Habitat in doliis Cereuisiae Ilgneis, impuris.

*geniculatus.* 25. *A.* niger, femorum geniculis subglobosis. *Syst. Ent.* 814. 20. *Linn. Syst. Nat.* 2. 1025. 19. *Fn. Sv.* 1977.

*Acarus* femoribus omnibus clauatis. *Schrank. Beytr.* 126. 26. tab. 6. fig. 9. 10.

Habitat in arborum ramis emortuis.

*baccarum.* 26. *A.* abdomine distento rubro, lateribus obscurioribus. *Syst. Ent.* 814. 21. *Linn. Syst. Nat.* 2. 1025. 23. *Fn. Sv.* 1980.

Schaeff.

**CLASS. VIII. ANTLIATA. Acarus; 491**

*Schaeff. Icon.* tab. 27. fig. 1.

*Scop. carn.* 1074.

*Schrank. Beytr.* 34. 13. tab. 1. fig. 30.

Habitat in Baccis Ribris rubri, Grossulariae aliisque.

27. *A. subrotundus albidus, abdomine rufo. It. Norwag. Zofterae.*  
d. 29. Iul. \*

Habitat in Fucis Oceani Norwagici.

28. *A. abdomine rufo, pedibus posticis longissimis filiformibus. Syst. Ent.* 814. 22. *Linn. Syst. Nat.* 2. 1025. *Muscarm.*  
24. *Fn. Sv.* 1981.

*Degeer Inf.* 7. 114. 10. tab. 7. fig. 1.

*Geoff. Inf.* 2. 624. 6.

Habitat in Europae Muscis.

29. *A. abdomine rubro vtrunque punctis binis coccineis. Gymno-*  
*Syst. Ent.* 814. 23. *Linn. Syst. Nat.* 2. 1026. 26. *pterorum.*  
*Fn. Sv.* 1982.

*Acarus Libellulae globosus ruber, pedibus brevissimis, stigmate dorsali concauo. Degeer Inf.* 7. 119.  
14. tab. 7. fig. 9.

*Reaum. Inf.* 5. tab. 38. fig. 1 — 3.

*Geoff. Inf.* 2. 625. 5.

Habitat in Vespis, Apibus, Afilis.

30. *A. ouatus rufus, ano albicante. Syst. Ent.* 814. 24. *Coleo-*  
*Linn. Syst. Nat.* 2. 1026. 27. *Fn. Sv.* 1973. *ptirato-*

*Acarus infectorum rufus, ano albicante. Geoff. Inf.*  
2. 623. 4. *rum.*

*Acarus*

492 CLASS. VIII. ANTLIATA. Acarus.

*Acarus fucorum* pallide fuscus flaescens macula dorsali triangulari obscura, pedibus anticis longioribus.  
*Degeer Inf.* 7. 112. 9. tab. 7. fig. 15.

*Blank. Inf.* tab. 14. fig. H.

*Frisch. Inf.* 4. tab. 10.

*Schrank. Beytr.* 10. 6. tab. 1. fig. 13. 14.

*Reaum. Inf.* 6. tab. 4. fig. 13. 14.

*Roef. Inf.* 4. tab. 1. fig. 10—15.

*Schaeff. Icon.* tab. 27. fig. 2.

Habitat in Scarabacis, Nicrophoris.

*Phalangii.* 31. *A. ouatus ruber*, rostro porrecto, pedibus longioribus.

*Degeer Inf.* 117. 12. tab. 7. fig. 5.

Habitat frequens in Phalangis Araneisque, arcte pedibus aliisque corporis partibus adhaerens.

*Phaëtontis.* 32. *A. ouatus postice acuminatus*, pedibus apice fasciculatis. *Syst. Ent.* 815. 25. \*

Habitat in Phaëtonte erubescente Oceani australis Fig. pict. *Mus. Dom. Banks.*

*rupestris.* 33. *A. fuscus*, linea dorsali bicolore. *Syst. Ent.* 815. 26.  
*Linn. Syst. Nat.* 2. 1026. 28. *Fn. Sv.* 1984.

*Schrank. Beytr.* 9. 5. tab. 1. fig. 11. 12.

Habitat sub Europae lapidibus frequens.

*longicornis.* 34. *A. ruber*, antennis bifidis rostro longioribus. *Syst. Ent.* 815. 27. *Linn. Syst. Nat.* 2. 1026. 29. *Fn. Sv.* 1985.

*Chelifer totus ruber*, antennis extremo bifetis. *Geoff. Inf.* 2. 618. 2. tab. 20. fig. 5.

Habitat in Europae rupibus.

35. *A. ouatus ruber*, rostro porrecto subulato. *Syst. Ent. littoralis.* 815. 28.

*Acarus fusco rufus*, pedibus sanguineis. *Linn. Syst. Nat.* 2. 1026. 30. *Fn. Sv.* 1986.

Habitat ad littora in petris.

Corpus glabrum, ovatum. Antennae filiformes quadriarticulatae. Caput et thorax sanguinea. Abdomen valde obtusum magis obscurum vtrinque punctis aliquot impressis.

36. *A. pallidus*, lineis duabus flexuosis nigris, pedibus *Fucorum*, posticis brevissimis incurvis. *It. Norwag.* d. 23. Iul.

Habitat in Oceani Norwagici Fucis.

37. *A. fusco rufescens*, abdomine subgloboso glabro im- *Fun-*  
maculato. *Syst. Ent.* 815. 29. *Linn. Syst. Nat.* 2. *rum.*  
1026. 31. *Fn. Sv.* 1987.

*Acarus* abdomine glabriusculo sub incessu globoso.  
*Schrank. Beytr.* 127. 27. tab. 6. fig. 11, 12.

Habitat gregarius in Fungis.

38. *A. subglobosus atro coerulescens.* *Syst. Ent.* 816. 30. *Tremel-*  
*Linn. Syst. Nat.* 2. 1026. 32. *Fn. Sv.* 1991. *lae.*

Habitat in Tremella iuniperina.

494 CLASS. VIII. ANTLIATA. Acarus.

*Salicinus*. 39. *A. ruber*, linea dorsali duplici fusca, antice bifurca.  
*Syst. Ent.* 816. 31. *Linn. Syst. Nat.* 2. 1027. 34. *Fn.*  
*Sv.* 1989.

Habitat in *Salicibus* *agilissimus*.

*croceus*. 40. *A. fulvus*, thoracis lateribus puncto rufo. *Syst.*  
*Ent.* 816. 32. *Linn. Syst. Nat.* 2. 1027. 35. *Fn.*  
*Sv.* 1990.

Habitat in *Salicis* *Rosa monstrosa*.

